

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great
Resource to assist you with your Studies 😊


Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



SA EXAM
PAPERS



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

SENIOR CERTIFICATE EXAMINATIONS/ NATIONAL SENIOR CERTIFICATE EXAMINATIONS

PHYSICAL SCIENCES: CHEMISRTY P2

2021

MARKS: 150

TIME: 3 hours

This question paper consists of 15 pages and 4 data sheets.



INSTRUCTIONS AND INFORMATION

1. Write your centre number and examination number in the appropriate spaces on the ANSWER BOOK.
2. This question paper consists of TEN questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two subquestions, e.g. between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. You may use appropriate mathematical instruments.
8. Show ALL formulae and substitutions in ALL calculations.
9. Round off your FINAL numerical answers to a minimum of TWO decimal places.
10. Give brief motivations, discussions, etc. where required.
11. You are advised to use the attached DATA SHEETS.
12. Write neatly and legibly.




QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Choose the answer and write only the letter (A–D) next to the question numbers (1.1 to 1.10) in the ANSWER BOOK, e.g. 1.11 E.

- 1.1 Which ONE of the following compounds has hydrogen bonds between molecules?
- A Pentanal
B Pentan-2-one
C Pentanoic acid
D Methyl butanoate (2)
- 1.2 To which homologous series does a compound with molecular formula $C_6H_{12}O_2$ belong?
- A Ketones
B Alcohols
C Aldehydes
D Carboxylic acids (2)
- 1.3 Which functional groups are involved in the formation of esters?
- A Formyl and carbonyl
B Hydroxyl and carbonyl
C Hydroxyl and carboxyl
D Carbonyl and carboxyl (2)
- 1.4 The equation below represents a reaction at equilibrium.
- $$\underset{\text{yellow}}{2CrO_4^{2-}(\text{aq})} + 2H^+(\text{aq}) \rightleftharpoons \underset{\text{orange}}{Cr_2O_7^{2-}(\text{aq})} + H_2O(l)$$
- Which ONE of the following will change the colour of the mixture from yellow to orange?
- A Addition of sodium hydroxide pellets
B Addition of concentrated hydrochloric acid
C Increase in pressure at constant temperature
D Decrease in pressure at constant temperature (2)

- 1.5 Consider the potential energy graph for the reaction shown below.




The activation energy for the FORWARD reaction in terms of P , Q and R is:

- A Q
- B $R - P$
- C $Q - R$
- D $Q - P$

(2)

- 1.6 A reaction reaches equilibrium in a closed container according to the following balanced equation:





Which ONE of the following changes will INCREASE the value of the equilibrium constant?

- A Removing $\text{NH}_3(\text{g})$
- B Heating the container
- C Cooling the container
- D Increasing the volume of the container

(2)

- 1.7 Sulphuric acid ionises in water according to the following equations:



Consider the following statements regarding the ionisation above:

- I:** $\text{H}_2\text{O}(\ell)$ acts as a base in both reactions.
II: $\text{HSO}_4^-(\text{aq})$ acts as an ampholyte.
III: $\text{SO}_4^{2-}(\text{aq})$ is the conjugate base of H_2SO_4 .

Which of the statements above is/are TRUE?

- A I only
 B I and II
 C I and III
 D I, II and III (2)

- 1.8 Which ONE of the following reactions, when used in a voltaic cell, will give a positive reading on the voltmeter?

- A $\text{Mg}^{2+}(\text{aq}) + \text{Zn}(\text{s}) \rightarrow \text{Mg}(\text{s}) + \text{Zn}^{2+}(\text{aq})$
 B $\text{Cu}(\text{s}) + 2\text{Ag}^+(\text{aq}) \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{Ag}(\text{s})$
 C $\text{Co}^{2+}(\text{aq}) + \text{Sn}^{2+}(\text{aq}) \rightarrow \text{Co}(\text{s}) + \text{Sn}^{4+}(\text{aq})$
 D $3\text{Ni}^{2+}(\text{aq}) + 2\text{Fe}(\text{s}) \rightarrow 3\text{Ni}(\text{s}) + 2\text{Fe}^{3+}(\text{aq})$ (2)

- 1.9 Which ONE of the following statements is CORRECT for an ELECTROLYTIC CELL?

- A The anode is the positive electrode.
 B The cathode is the positive electrode.
 C Oxidation takes place at the cathode.
 D Reduction takes place at the anode. (2)

- 1.10 Which ONE of the following shows the industrial processes in which AMMONIA is a reactant and a product respectively?

	REACTANT	PRODUCT
A	Ostwald	Contact
B	Ostwald	Haber
C	Contact	Haber
D	Contact	Ostwald

(2)
[20]

QUESTION 2 (Start on a new page.)

The letters **A** to **F** in the table below represent six organic compounds.

A	Methanoic acid	B	Pentanal
C	$C_{10}H_{22}$	D	$ \begin{array}{c} \text{Br} \\ \\ \text{CH}-\text{CH}_2-\text{CH}_3 \\ \\ \text{CH}_3(\text{CH}_2)_2\text{CH}-\text{CH}_2 \\ \\ \text{Br} \end{array} $
E	$ \begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{O}-\text{H} \\ \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{C}-\text{C}-\text{H} \\ \quad \quad \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \end{array} $	F	$ \begin{array}{ccccccc} & \text{H} & & \text{H} & & \text{H} & & \text{O} & & \text{H} \\ & & & & & & & & & \\ \text{H} & -\text{C} & - & \text{C} & - & \text{C} & - & \text{C} & - & \text{C} & - \text{H} \\ & & & & & & & & & \\ & \text{H} & & \text{H} & & \text{H} & & & & \text{H} \end{array} $

- 2.1 Write down the LETTER(S) that represent(s) the following:
- 2.1.1 A ketone (1)
- 2.1.2 TWO compounds that are FUNCTIONAL ISOMERS (1)
- 2.1.3 A hydrocarbon (1)
- 2.2 For compound **D**, write down the:
- 2.2.1 Homologous series to which it belongs (1)
- 2.2.2 IUPAC name (3)
- 2.3 Consider compound **F**.
- Write down the IUPAC name of its:
- 2.3.1 POSITIONAL isomer (2)
- 2.3.2 CHAIN isomer (2)
- 2.4 During the reaction of compound **A** with compound **E** in the presence of an acid catalyst, two products are formed.
- For the ORGANIC product formed, write down the:
- 2.4.1 IUPAC name (2)
- 2.4.2 STRUCTURAL FORMULA of its FUNCTIONAL GROUP (1)

- 2.5 Compound **C** ($\text{C}_{10}\text{H}_{22}$) reacts at high temperatures and pressures to form a three-carbon alkene **P** and an alkane **Q**, as shown below.



Write down the:

- | | | |
|-------|---|-----|
| 2.5.1 | Type of reaction that takes place | (1) |
| 2.5.2 | Molecular formula of compound Q | (2) |
| 2.5.3 | STRUCTURAL FORMULA of compound P | (2) |
- [19]**




QUESTION 3 (Start on a new page.)

Learners use compounds **A**, **B** and **C** to investigate one of the factors that influences the VAPOUR PRESSURE of organic compounds.

A	Butan-1-ol
B	Butan-2-one
C	Propanoic acid

- 3.1 Define the term *vapour pressure*. (2)
- 3.2 Write down the independent variable for this investigation. (1)
- 3.3 Which compound, **A** or **B**, has the higher vapour pressure? (1)
- 3.4 Fully explain the answer to QUESTION 3.3. Include the TYPES OF INTERMOLECULAR FORCES in your explanation. (4)
- 3.5 The graph below represents the relationship between vapour pressure and temperature for compound **A** at sea level. **X** and **Y** represent different temperatures.




- 3.5.1 Write down the term for the temperature represented by **X**. (1)
- 3.5.2 State the phase of compound **A** at temperature **Y**. Choose from GAS, LIQUID or SOLID. (1)
- 3.5.3 Redraw the graph above in your ANSWER BOOK. On the same set of axes, sketch the curve that will be obtained for compound **C**. Clearly label curve **A** and curve **C**. (2)

[12]

QUESTION 4 (Start on a new page.)

4.1 The flow diagram below shows the conversion of propane to propan-2-ol.



4.1.1 State ONE reaction condition for **Step 1**. (1)

4.1.2 Write down the NAME or FORMULA of the INORGANIC product formed in **Step 1**. (1)

4.1.3 Name the TYPE of substitution reaction represented by **Step 2**. (1)

4.1.4 Write down the NAME or FORMULA of the INORGANIC reagent needed in **Step 2**. (1)

4.1.5 Write down the IUPAC name of compound X. (2)

4.2 Ethane can be prepared from chloroethane ($\text{CH}_3\text{CH}_2\text{Cl}$) by a TWO-STEP process. You are supplied with the following chemicals:

H_2	HCl	Cl_2	H_2O	Pt	Ethanol	concentrated H_2SO_4	concentrated NaOH
--------------	--------------	---------------	----------------------	----	---------	--------------------------------------	-------------------


Select chemicals in the table above that can be used for this preparation.

Using CONDENSED structural formulae, write down a balanced equation for EACH reaction. Indicate the reaction conditions for EACH reaction. (8)


[14]

QUESTION 5 (Start on a new page.)

Two experiments, **I** and **II**, are conducted to investigate one of the factors that affects the rate of the reaction of aluminium carbonate, $\text{Al}_2(\text{CO}_3)_3$, with EXCESS hydrochloric acid, HCl . The balanced equation for the reaction is:



The apparatus used is shown below.



The reaction conditions used for each experiment are as follows:

Experiment I:

100 cm^3 of $1,5 \text{ mol} \cdot \text{dm}^{-3}$ $\text{HCl}(\text{aq})$ reacts with $0,016 \text{ mol}$ $\text{Al}_2(\text{CO}_3)_3$ granules at 25°C

Experiment II:


50 cm^3 of $2 \text{ mol} \cdot \text{dm}^{-3}$ $\text{HCl}(\text{aq})$ reacts with $0,016 \text{ mol}$ $\text{Al}_2(\text{CO}_3)_3$ granules at 25°C

- 5.1 Define the term *rate of a reaction*. (2)
- 5.2 Using the experimental setup above, state the measurements that must be made to determine the rate of this reaction. (2)
- 5.3 Use the collision theory to explain how the average reaction rate in **Experiment I** differs from the average reaction rate in **Experiment II**. (3)
- 5.4 The average rate of the reaction in **Experiment II** during the first 2,5 minutes is $4,4 \times 10^{-3} \text{ mol} \cdot \text{min}^{-1}$.
Calculate the number of moles of $\text{Al}_2(\text{CO}_3)_3$ that remains in the flask after 2,5 minutes. (3)
- 5.5 Calculate the maximum volume of $\text{CO}_2(\text{g})$ that can be prepared at 25°C in **Experiment I**. Take molar gas volume at 25°C as $24\,000 \text{ cm}^3 \cdot \text{mol}^{-1}$. (3)


[13]

QUESTION 6 (Start on a new page.)

Pure hydrogen iodide gas, HI(g) , of concentration $1 \text{ mol}\cdot\text{dm}^{-3}$, is sealed in a 500 cm^3 container at temperature T . The reaction reaches equilibrium according to the following balanced equation:




- 6.1 Define the term *chemical equilibrium*. (2)
- 6.2 The graph below shows how the concentrations of the reactant and products vary with time during the reaction.



- 6.2.1 Which ONE of the curves, X or Y, represents the changes in the concentration of the products? Give a reason for the answer. (2)
- 6.2.2 How does the rate of the forward reaction compare to that of the reverse reaction at $t = 4$ minutes? Choose from HIGHER THAN, LOWER THAN or EQUAL TO. (1)
- 6.3 The equilibrium constant, K_c , for the reaction is 0,04 at temperature T . Calculate the number of moles of iodine, $\text{I}_2\text{(g)}$, present at time $t = 6$ minutes. (9)

- 6.4 The graph below shows how the rates of the forward and reverse reactions change with time.



The temperature of the container is increased at $t = 10$ minutes.



- 6.4.1 Which reaction(s) show(s) an increase in rate at $t = 10$ minutes? Choose from FORWARD, REVERSE or BOTH FORWARD AND REVERSE. (1)
- 6.4.2 Is the heat of reaction (ΔH) for this reaction POSITIVE or NEGATIVE? Fully explain the answer. (4)
- [19]**

QUESTION 7 (Start on a new page.)

Learners prepare a solution of known concentration by dissolving 2 g pure sodium hydroxide crystals, NaOH, in water in a 250 cm³ volumetric flask.

- 7.1 Write down the term for the underlined phrase. (1)
- 7.2 Calculate the: (4)
- 7.2.1 Concentration of the sodium hydroxide solution (4)
- 7.2.2 pH of the solution (4)

The learners now react 1,5 g of pure CaCO₃ with 50 cm³ dilute HCl of unknown concentration. The EXCESS HCl is neutralised with 25 cm³ of the NaOH solution that they prepared. The balanced equations for the reactions are:



- 7.3 Calculate the initial concentration of the dilute HCl(aq). (8)
- [17]**



QUESTION 8 (Start on a new page.)

- 8.1 When a piece of sodium metal (Na) is added to water in a test tube, hydrogen gas is released. When phenolphthalein indicator is added to the test tube, the solution turns pink.

8.1.1 Define the term *reduction* in terms of electron transfer. (2)

8.1.2 Write down the reduction half-reaction. (2)


8.1.3 Write down the balanced equation for the reaction that takes place (3)

8.1.4 Give a reason why the solution turns pink. (1)

When a piece of copper is added to water in a test tube, no reaction is observed.

8.1.5 Refer to the relative strengths of the REDUCING AGENTS to explain why no reaction is observed. (3)

- 8.2 Consider the cell notation below.



8.2.1 What does the single line (|) in the cell notation above represent? (1)

8.2.2 State the energy conversion that takes place in this cell. (1)



8.2.3 Calculate the initial emf of the cell under standard conditions. (4)

[17]



QUESTION 9 (Start on a new page.)





The diagrams below show two electrochemical cells in which carbon electrodes are used. In cell **A**, concentrated copper (II) chloride solution is used and in cell **B**, liquid aluminium oxide is used.

**Cell A****Cell B**

- 9.1 What type of electrochemical cell, ELECTROLYTIC or GALVANIC, is shown above? Give a reason for the answer. (2)
- 9.2 Write down the:
- 9.2.1 Half-reaction that takes place at the anode of cell **A** (2)
- 9.2.2 Half-reaction that takes place at the cathode of cell **B** (2)
- 9.2.3 NAME or FORMULA of the product formed at the cathode of cell **A** (1)
- 9.3 Give a reason why the mass of electrode **Y** decreases after a while. (1)
- [8]**

QUESTION 10 (Start on a new page.)

- 10.1 The incomplete equations below show the four steps involved in the industrial preparation of sulphuric acid (H_2SO_4). **A** and **B** represent two compounds.



Write down the NAME or FORMULA of:

10.1.1 Compound **A** (1)

10.1.2 Compound **B** (1)

10.1.3 The catalyst used in **Step II** (1)

The sulphuric acid formed in **Step IV** is used to prepare ammonium sulphate.

10.1.4 Write down a balanced equation for this reaction. (3)

- 10.2 The diagram below shows a bag of fertiliser.



10.2.1 Write down the meaning of NPK. (1)

10.2.2 The bag contains 4 kg of ammonium nitrate, NH_4NO_3 , which is the only source of nitrogen. Calculate the mass of the fertiliser in the bag. (4)
[11]

TOTAL: 150

**DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)**

**GEGEWENS VIR FISIESE WETENSKAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)**

TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESE KONSTANTES

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure <i>Standaarddruk</i>	p^θ	$1,013 \times 10^5 \text{ Pa}$
Molar gas volume at STP <i>Molêre gasvolume by STD</i>	V_m	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard temperature <i>Standaardtemperatuur</i>	T^θ	273 K
Charge on electron <i>Lading op elektron</i>	e	$-1,6 \times 10^{-19} \text{ C}$
Avogadro's constant <i>Avogadro-konstante</i>	N_A	$6,02 \times 10^{23} \text{ mol}^{-1}$

TABLE 2: FORMULAE/TABEL 2: FORMULES

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ or/of $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$
$\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$	$\text{pH} = -\log[\text{H}_3\text{O}^+]$
$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14} \text{ at/by } 298 \text{ K}$	
$E_{\text{cell}}^\theta = E_{\text{cathode}}^\theta - E_{\text{anode}}^\theta / E_{\text{sel}}^\theta = E_{\text{katode}}^\theta - E_{\text{anode}}^\theta$ or/of $E_{\text{cell}}^\theta = E_{\text{reduction}}^\theta - E_{\text{oxidation}}^\theta / E_{\text{sel}}^\theta = E_{\text{reduksie}}^\theta - E_{\text{oksidasie}}^\theta$ or/of $E_{\text{cell}}^\theta = E_{\text{oxidisingagent}}^\theta - E_{\text{reducingagent}}^\theta / E_{\text{sel}}^\theta = E_{\text{oksideermiddel}}^\theta - E_{\text{reduseermiddel}}^\theta$	

TABLE 3: THE PERIODIC TABLE OF ELEMENTS
TABEL 3: DIE PERIODIEKE TABEL VAN ELEMENTE

KEY/SLEUTEL																	
Atomic number Atoomgetal												Atomic number Atoomgetal					
Electronegativity Elektronegativiteit												Symbol Simbool					
Approximate relative atomic mass Benaderde relatieve atoommassa												Approximate relative atomic mass Benaderde relatieve atoommassa					
29 Cu 63,5												29 Cu 63,5					
1 H 1,0												1 H 1,0					
3 Li 6,9												3 Li 6,9					
11 Na 23												11 Na 23					
19 K 39												19 K 39					
37 Rb 86												37 Rb 86					
55 Cs 133												55 Cs 133					
87 Fr 226												87 Fr 226					
21 Sc 45												21 Sc 45					
22 Ti 48												22 Ti 48					
23 V 51												23 V 51					
24 Cr 52												24 Cr 52					
25 Mn 55												25 Mn 55					
26 Fe 56												26 Fe 56					
27 Co 59												27 Co 59					
28 Ni 59												28 Ni 59					
29 Cu 63,5												29 Cu 63,5					
30 Zn 65												30 Zn 65					
31 Ga 70												31 Ga 70					
32 Ge 73												32 Ge 73					
33 As 75												33 As 75					
34 Se 79												34 Se 79					
35 Br 80												35 Br 80					
36 Kr 84												36 Kr 84					
37 Rb 86												37 Rb 86					
38 Sr 88												38 Sr 88					
39 Y 89												39 Y 89					
40 Zr 91												40 Zr 91					
41 Nb 92												41 Nb 92					
42 Mo 96												42 Mo 96					
43 Tc 101												43 Tc 101					
44 Ru 103												44 Ru 103					
45 Rh 106												45 Rh 106					
46 Pd 108												46 Pd 108					
47 Ag 108												47 Ag 108					
48 Cd 112												48 Cd 112					
49 In 115												49 In 115					
50 Sn 119												50 Sn 119					
51 Sb 122												51 Sb 122					
52 Te 128												52 Te 128					
53 I 127												53 I 127					
54 Xe 131												54 Xe 131					
55 Cs 133												55 Cs 133					
56 Ba 137												56 Ba 137					
57 La 139												57 La 139					
58 Ce 140												58 Ce 140					
59 Pr 141												59 Pr 141					
60 Nd 144												60 Nd 144					
61 Pm 147												61 Pm 147					
62 Sm 150												62 Sm 150					
63 Eu 152												63 Eu 152					
64 Gd 157												64 Gd 157					
65 Tb 159												65 Tb 159					
66 Dy 163												66 Dy 163					
67 Ho 165												67 Ho 165					
68 Er 167												68 Er 167					
69 Tm 169												69 Tm 169					
70 Yb 173												70 Yb 173					
71 Lu 175												71 Lu 175					
72 Hf 179												72 Hf 179					
73 Ta 181												73 Ta 181					
74 W 184												74 W 184					
75 Re 186												75 Re 186					
76 Os 190												76 Os 190					
77 Ir 192												77 Ir 192					
78 Pt 195												78 Pt 195					
79 Au 197												79 Au 197					
80 Hg 201												80 Hg 201					
81 Tl 204												81 Tl 204					
82 Pb 207												82 Pb 207					
83 Bi 209												83 Bi 209					
84 Po 209												84 Po 209					
85 At 210												85 At 210					
86 Rn 222												86 Rn 222					
87 Fr 223												87 Fr 223					
88 Ra 226												88 Ra 226					
89 Ac 227												89 Ac 227					
90 Th 232												90 Th 232					
91 Pa 231												91 Pa 231					
92 U 238												92 U 238					
93 Np 237												93 Np 237					
94 Pu 242												94 Pu 242					
95 Am 243												95 Am 243					
96 Cm 247												96 Cm 247					
97 Bk 247												97 Bk 247					
98 Cf 251												98 Cf 251					
99 Es 252												99 Es 252					
100 Fm 257												100 Fm 257					
101 Md 258												101 Md 258					
102 No 259												102 No 259					
103 Lr 262												103 Lr 262					

KEY/SLEUTEL

Atomic number
AtoomgetalElectronegativity
ElektronegatiwiteitSymbol
SimboolApproximate relative atomic mass
Benaderde relatiewe atoommassa

TABLE 4A: STANDARD REDUCTION POTENTIALS
TABEL 4A: STANDAARD- REDUKSIEPOTENSIALE

Half-reactions/Halfreaksies	E^{θ} (V)
$F_2(g) + 2e^- \rightleftharpoons 2F^-$	+ 2,87
$Co^{3+} + e^- \rightleftharpoons Co^{2+}$	+ 1,81
$H_2O_2 + 2H^+ + 2e^- \rightleftharpoons 2H_2O$	+ 1,77
$MnO_4^- + 8H^+ + 5e^- \rightleftharpoons Mn^{2+} + 4H_2O$	+ 1,51
$Cl_2(g) + 2e^- \rightleftharpoons 2Cl^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^- \rightleftharpoons 2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^- \rightleftharpoons 2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^- \rightleftharpoons Mn^{2+} + 2H_2O$	+ 1,23
$Pt^{2+} + 2e^- \rightleftharpoons Pt$	+ 1,20
$Br_2(l) + 2e^- \rightleftharpoons 2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^- \rightleftharpoons NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^- \rightleftharpoons Hg(l)$	+ 0,85
$Ag^+ + e^- \rightleftharpoons Ag$	+ 0,80
$NO_3^- + 2H^+ + e^- \rightleftharpoons NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^- \rightleftharpoons Fe^{2+}$	+ 0,77
$O_2(g) + 2H^+ + 2e^- \rightleftharpoons H_2O_2$	+ 0,68
$I_2 + 2e^- \rightleftharpoons 2I^-$	+ 0,54
$Cu^+ + e^- \rightleftharpoons Cu$	+ 0,52
$SO_2 + 4H^+ + 4e^- \rightleftharpoons S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^- \rightleftharpoons 4OH^-$	+ 0,40
$Cu^{2+} + 2e^- \rightleftharpoons Cu$	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^- \rightleftharpoons SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^- \rightleftharpoons Cu^+$	+ 0,16
$Sn^{4+} + 2e^- \rightleftharpoons Sn^{2+}$	+ 0,15
$S + 2H^+ + 2e^- \rightleftharpoons H_2S(g)$	+ 0,14
$2H^+ + 2e^- \rightleftharpoons H_2(g)$	0,00
$Fe^{3+} + 3e^- \rightleftharpoons Fe$	- 0,06
$Pb^{2+} + 2e^- \rightleftharpoons Pb$	- 0,13
$Sn^{2+} + 2e^- \rightleftharpoons Sn$	- 0,14
$Ni^{2+} + 2e^- \rightleftharpoons Ni$	- 0,27
$Co^{2+} + 2e^- \rightleftharpoons Co$	- 0,28
$Cd^{2+} + 2e^- \rightleftharpoons Cd$	- 0,40
$Cr^{3+} + e^- \rightleftharpoons Cr^{2+}$	- 0,41
$Fe^{2+} + 2e^- \rightleftharpoons Fe$	- 0,44
$Cr^{3+} + 3e^- \rightleftharpoons Cr$	- 0,74
$Zn^{2+} + 2e^- \rightleftharpoons Zn$	- 0,76
$2H_2O + 2e^- \rightleftharpoons H_2(g) + 2OH^-$	- 0,83
$Cr^{2+} + 2e^- \rightleftharpoons Cr$	- 0,91
$Mn^{2+} + 2e^- \rightleftharpoons Mn$	- 1,18
$Al^{3+} + 3e^- \rightleftharpoons Al$	- 1,66
$Mg^{2+} + 2e^- \rightleftharpoons Mg$	- 2,36
$Na^+ + e^- \rightleftharpoons Na$	- 2,71
$Ca^{2+} + 2e^- \rightleftharpoons Ca$	- 2,87
$Sr^{2+} + 2e^- \rightleftharpoons Sr$	- 2,89
$Ba^{2+} + 2e^- \rightleftharpoons Ba$	- 2,90
$Cs^+ + e^- \rightleftharpoons Cs$	- 2,92
$K^+ + e^- \rightleftharpoons K$	- 2,93
$Li^+ + e^- \rightleftharpoons Li$	- 3,05

Increasing oxidising ability/Toenemende oksiderende vermoë

Increasing reducing ability/Toenemende reduserende vermoë



TABLE 4B: STANDARD REDUCTION POTENTIALS
TABEL 4B: STANDAARD- REDUKSIEPOTENSIALE

Half-reactions/Halfreaksies	E^θ (V)
$\text{Li}^+ + \text{e}^- \rightleftharpoons \text{Li}$	- 3,05
$\text{K}^+ + \text{e}^- \rightleftharpoons \text{K}$	- 2,93
$\text{Cs}^+ + \text{e}^- \rightleftharpoons \text{Cs}$	- 2,92
$\text{Ba}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ba}$	- 2,90
$\text{Sr}^{2+} + 2\text{e}^- \rightleftharpoons \text{Sr}$	- 2,89
$\text{Ca}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ca}$	- 2,87
$\text{Na}^+ + \text{e}^- \rightleftharpoons \text{Na}$	- 2,71
$\text{Mg}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mg}$	- 2,36
$\text{Al}^{3+} + 3\text{e}^- \rightleftharpoons \text{Al}$	- 1,66
$\text{Mn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mn}$	- 1,18
$\text{Cr}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cr}$	- 0,91
$2\text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-$	- 0,83
$\text{Zn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Zn}$	- 0,76
$\text{Cr}^{3+} + 3\text{e}^- \rightleftharpoons \text{Cr}$	- 0,74
$\text{Fe}^{2+} + 2\text{e}^- \rightleftharpoons \text{Fe}$	- 0,44
$\text{Cr}^{3+} + \text{e}^- \rightleftharpoons \text{Cr}^{2+}$	- 0,41
$\text{Cd}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cd}$	- 0,40
$\text{Co}^{2+} + 2\text{e}^- \rightleftharpoons \text{Co}$	- 0,28
$\text{Ni}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ni}$	- 0,27
$\text{Sn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Sn}$	- 0,14
$\text{Pb}^{2+} + 2\text{e}^- \rightleftharpoons \text{Pb}$	- 0,13
$\text{Fe}^{3+} + 3\text{e}^- \rightleftharpoons \text{Fe}$	- 0,06
$2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g})$	0,00
$\text{S} + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2\text{S}(\text{g})$	+ 0,14
$\text{Sn}^{4+} + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}$	+ 0,15
$\text{Cu}^{2+} + \text{e}^- \rightleftharpoons \text{Cu}^+$	+ 0,16
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{SO}_2(\text{g}) + 2\text{H}_2\text{O}$	+ 0,17
$\text{Cu}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cu}$	+ 0,34
$2\text{H}_2\text{O} + \text{O}_2 + 4\text{e}^- \rightleftharpoons 4\text{OH}^-$	+ 0,40
$\text{SO}_2 + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons \text{S} + 2\text{H}_2\text{O}$	+ 0,45
$\text{Cu}^+ + \text{e}^- \rightleftharpoons \text{Cu}$	+ 0,52
$\text{I}_2 + 2\text{e}^- \rightleftharpoons 2\text{I}^-$	+ 0,54
$\text{O}_2(\text{g}) + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2\text{O}_2$	+ 0,68
$\text{Fe}^{3+} + \text{e}^- \rightleftharpoons \text{Fe}^{2+}$	+ 0,77
$\text{NO}_3^- + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{NO}_2(\text{g}) + \text{H}_2\text{O}$	+ 0,80
$\text{Ag}^+ + \text{e}^- \rightleftharpoons \text{Ag}$	+ 0,80
$\text{Hg}^{2+} + 2\text{e}^- \rightleftharpoons \text{Hg}(\ell)$	+ 0,85
$\text{NO}_3^- + 4\text{H}^+ + 3\text{e}^- \rightleftharpoons \text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+ 0,96
$\text{Br}_2(\ell) + 2\text{e}^- \rightleftharpoons 2\text{Br}^-$	+ 1,07
$\text{Pt}^{2+} + 2\text{e}^- \rightleftharpoons \text{Pt}$	+ 1,20
$\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 2\text{H}_2\text{O}$	+ 1,23
$\text{O}_2(\text{g}) + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$	+ 1,23
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightleftharpoons 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+ 1,33
$\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-$	+ 1,36
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 4\text{H}_2\text{O}$	+ 1,51
$\text{H}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$	+ 1,77
$\text{Co}^{3+} + \text{e}^- \rightleftharpoons \text{Co}^{2+}$	+ 1,81
$\text{F}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{F}^-$	+ 2,87

Increasing oxidising ability/Toenemende oksiderende vermoë

Increasing reducing ability/Toenemende reduserende vermoë

