

SA's Leading Past Year

Exam Paper Portal

STUDY

You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers


Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



SA EXAM
PAPERS

QUESTION 1

<p>1.1</p>	 <p> $y - 2x + 5 = 4$ $y - 2x = -1 \dots (1)$ $29 - 2y + x = 4$ $-2y + x = -25 \dots (2)$ $(1) \times 2: 2y - 4x = -2 \dots (3)$ $(3) + (2):$ $-3x = -27 \quad \therefore x = 9$ $\quad \quad \quad \therefore y = 17$ </p>	<p>A✓ setting up equation 1st line</p> <p>A✓ setting up equation 1st line</p> <p>CA✓ x – value</p> <p>CA✓ y – value</p>	<p>(4)</p>
<p>1.2</p>	<p> $2a = 4 \quad \therefore a = 2$ $3a + b = 4 \quad \therefore b = -2$ $a + b + c = 5 \quad \therefore c = 5$ $T_n = 2n^2 - 2n + 5$ </p> <p>OR</p> <p> $2a = 4 \quad \therefore a = 2$ $3a + b = 5 \quad \therefore b = -2$ $\therefore c = T_0 = 5$ $T_n = 2n^2 - 2n + 5$ </p> <p>OR</p> $T_n = T_1 + (n - 1)d_1 + \frac{(n - 1)(n - 2)}{2}d_2$ <p>OR</p> $T_n = \frac{(n - 1)}{2} [2a + (n - 2)d] + T_1$	<p>A✓ a – value</p> <p>CA✓ b – value</p> <p>CA✓ c – value</p> <p>CA✓ nth term</p> <p>OR</p> <p>A✓ a – value</p> <p>CA✓ b – value</p> <p>CA✓ c – value</p> <p>CA✓ nth term</p> <p>OR</p> <p>OR</p>	<p>(4)</p> <p>(4)</p>
<p>1.3</p>	<p>$T_{50} = 2(50)^2 - 2(50) + 5 = 4905$</p>	<p>CA✓ substitution (from 1.2)</p> <p>CA✓ answer</p>	<p>(2)</p>
			<p>[10]</p>


QUESTION 2

2	$a + d = 8 \quad \rightarrow (1)$ $a + 6d = 11a \quad \rightarrow (2)$ $d = 8 - a \quad \rightarrow (3)$ $a + 6(8 - a) = 11a$ $a + 48 - 6a = 11a$ $\quad \quad \quad 48 = 16a$ $\quad \quad \quad a = 3$ $\quad \quad \quad d = 5$ 3 ; 8 ; 13 ; ...	A✓equation (1) A✓equation (2) CA✓making d/a the subject CA✓correct substitution of d/a CA✓ a/d value CA✓ d/a value CA✓sequence	[7]
---	--	--	------------

QUESTION 3


3.1.1	$1 ; \frac{1}{p} ; \frac{1}{p^2}$	A✓All three terms	(1)
3.1.2	$-1 < r < 1$ $-1 < \frac{1}{p} < 1$ $p < -1 \text{ or } p > 1$	A✓condition for convergence CA✓ r value in terms of p CACA✓✓answers	(4)
3.2	$S_n = 4 - 4\left(\frac{1}{2}\right)^n$ $T_1 = S_1 = 4 - 4\left(\frac{1}{2}\right)^1 = 2$ $T_1 + T_2 = S_2 = 4 - 4\left(\frac{1}{2}\right)^2 = 3$ $T_1 + T_2 + T_3 = S_3 = 4 - 4\left(\frac{1}{2}\right)^3 = 3\frac{1}{2}$ $\therefore T_2 = 1$ $\therefore T_3 = \frac{1}{2}$	A✓ first term value A✓ Sum of first two terms A✓ Sum of first 3 terms CA✓second term value CA✓third term value	(5)
			[10]

QUESTION 4



<p>4.1</p>	<p>On AB mark off a point P such that $AP = DE$ and on AC mark off a point Q such that $AQ = DF$. Join PQ. In $\triangle ABC$ and $\triangle DEF$ 1. $AP = DE$ (Construction) 2. $AQ = DF$ (Construction) 3. $\hat{A} = \hat{D}$ (Given) $\therefore \triangle APQ \equiv \triangle DEF$ (SAS) Now $\hat{APQ} = \hat{DEF}$ But $\hat{DEF} = \hat{B}$ (Given) $\therefore \hat{APQ} = \hat{B}$ $PQ \parallel BC$ (Corresponding angles =) $\frac{AB}{AP} = \frac{AC}{AQ}$ (Prop. Thm. $PQ \parallel BC$) $\frac{AB}{DE} = \frac{AC}{DF}$ (Construction $AP = DE$ and $AQ = DF$)</p>	<p>✓S Construction ✓S/R ✓S ✓S ✓S/R ✓S/R ✓R</p>	<p>[7]</p>
------------	--	---	------------

4.2



4.2.1	$\frac{AE}{15} = \frac{AD}{3a} = \frac{3}{5} \dots \text{(Prop. Thm. } ED \parallel BC)$ $AE = 9 \text{ units}$	✓S✓R ✓Answer	(3)
4.2.2	$\frac{AH}{15} = \frac{CG}{CB} = \frac{1}{3} \dots \text{(Prop. Thm. } HG \parallel AC)$ $AH = 5 \text{ units}$	✓S✓R ✓Answer	(3)
4.2.3	HE = 4 units and BH = 6 units $\frac{GK}{KH} = \frac{BE}{EH} = \frac{6}{4} = \frac{3}{2} \dots \text{(Prop. Thm. } EK \parallel BG)$	CA✓HE = 4 units CA✓S A✓R	(3)
4.2.4	$\frac{\Delta HEK}{\Delta HEG} = \frac{2}{5} \dots \text{(\Delta s with equal altitudes)}$ $\frac{\Delta HEG}{\Delta HBG} = \frac{4}{10} = \frac{2}{5} \dots \text{(\Delta s with equal altitudes)}$ $\frac{\text{Area of } \Delta HEK}{\text{Area of } \Delta HBG} = \frac{\Delta HEK}{\Delta HEG} \times \frac{\Delta HEG}{\Delta HBG} = \frac{2}{5} \times \frac{2}{5} = \frac{4}{25}$ <p>OR</p> $\frac{\Delta HEK}{\Delta HBG} = \frac{\frac{1}{2} \cdot (4)(2c) \sin \widehat{E\hat{H}K}}{\frac{1}{2} \cdot (10)(5c) \sin \widehat{E\hat{H}K}}$ $\frac{\Delta HEK}{\Delta HBG} = \frac{4}{25}$ <p>OR</p> $\Delta HEK = \frac{2}{5} (\Delta HEG) \dots \text{(\Delta s with equal altitudes)}$ $= \frac{2}{5} \left(\frac{2}{5} \Delta HBG \right) \dots \text{(\Delta s with equal altitudes)}$ $\frac{\text{Area of } \Delta HEK}{\text{Area of } \Delta HBG} = \frac{4}{25}$	✓S/R ✓S/R ✓Answer OR ✓Area of numerator ✓Area of Denominator ✓Answer OR ✓S/R ✓S/R ✓Answer	(3)
			(3)


QUESTION 6

<p>6.1.1</p>	<p style="text-align: center;">$\sin 20^\circ = \sqrt{1 - p^2}$</p>	<p>M✓ A✓ Answer</p>	<p>(2)</p>
<p>6.1.2</p>	$\begin{aligned} \cos 6^\circ &= \cos(20^\circ - 14^\circ) \\ &= \cos 20 \cos 14 + \sin 20 \sin 14 \\ &= p \cdot \sqrt{1 - q^2} + \sqrt{1 - p^2} \cdot q \end{aligned}$	<p>A✓ Writing as difference A✓ Expansion A✓ ($\cos 20^\circ$) value CA ✓ $\sqrt{1 - q^2}$ CA ✓ $\sqrt{1 - p^2}$ A ✓ ($\sin 14^\circ$) value</p>	<p>(6)</p>
<p>6.2</p>	$\begin{aligned} &\sqrt{\frac{\frac{1}{2} \sin 2x}{\tan(540^\circ + x) \left(\frac{1}{\cos^2 x} - \tan^2 x \right)}} \\ &= \sqrt{\frac{\frac{1}{2} (2 \sin x \cos x)}{\tan x \cdot \left(\frac{1 - \sin^2 x}{\cos^2 x} \right)}} \\ &= \sqrt{\frac{\frac{1}{2} (2 \sin x \cos x)}{\frac{\sin x \cdot \cos^2 x}{\cos x \cdot \cos^2 x}}} \\ &= \sqrt{\sin x \cos x \times \frac{\cos x}{\sin x}} \\ &= \sqrt{\cos^2 x} \\ &= \cos x \end{aligned}$	<p>A✓ expansion of $\sin 2x$ A✓ reduction to $\tan x$</p> <p>A✓ $1 - \sin^2 x = \cos^2 x$ A✓ $\tan x = \frac{\sin x}{\cos x}$</p> <p>CA✓ simplifying $\sqrt{\cos^2 x}$</p> <p>CA✓ answer</p>	<p>(6)</p>
			<p>[14]</p>

QUESTION 7

7.1.1	$\cos 4x$ $= \cos(2 \cdot 2x)$ $= 2 \cos^2 2x - 1$ $= 2(2 \cos^2 x - 1)^2 - 1$ $= 2(4 \cos^4 x - 4 \cos^2 x + 1) - 1$ $= 8 \cos^4 x - 8 \cos^2 x + 1$	A✓ $\cos 4x = \cos(2 \cdot 2x)$ A✓ expansion of $\cos 4x$ A✓ expansion of $\cos 2x$ A✓ squaring bracket	(4)
7.1.2	$2 \cos 4x = 1$ $\cos 4x = \frac{1}{2}$ $4x = 60^\circ + 360k$ or $4x = 300^\circ / -60^\circ + 360k$ $x = 15^\circ + 90k$ or $x = 75^\circ / -15^\circ + 90k; k \in \mathbb{Z}$	A✓ setting up equation CA✓ both solutions A✓ $90k$ A✓ $k \in \mathbb{Z}$ CA✓ $x = 15^\circ, 75^\circ / -15^\circ$	(5)
7.1.3	$16 \cos^4 x - 16 \cos^2 x + 2$ $= 2 \cos 4x$ $\therefore \text{Minimum Value} = -2$	A✓ A✓	(2)
7.2	$\frac{2 \sin^2 22.5^\circ - 1}{4 \sin 22.5^\circ \cos 22.5^\circ}$ $= \frac{-(1 - 2 \sin^2 22.5^\circ)}{2 \cdot 2 \sin 22.5^\circ \cos 22.5^\circ}$ $= \frac{-\cos 45^\circ}{2 \cdot \sin 45^\circ}$ $= \frac{-1}{2} \times \frac{\frac{1}{\sqrt{2}}}{\frac{1}{\sqrt{2}}}$ $= -\frac{1}{2} \times 1 = -\frac{1}{2}$	A✓ re-writing numerator and denominator A✓ Numerator A✓ Denominator A✓ Special angle values/ co - ratios A✓ Answer	(5)
			[16]

QUESTION 8



8.1	f : x -int., Maximum and Minimum and Shape g : y -int., Maximum and Minimum and Shape	AAA✓✓✓ AAA✓✓✓	(6)
8.2	$\frac{g(x)}{f(x)} \geq 1$ $g(x) \geq f(x); f(x) > 0$ $x \in (0^\circ; 180^\circ); x \neq 90^\circ$	AA✓✓ Answer	(2)
			[8]

Total Marks: 100