

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great
Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



**SA EXAM
PAPERS**



**NATIONAL
SENIOR CERTIFICATE/
NASIONALE SENIOR
SERTIFIKAAT**

GRADE/GRAAD 12

SEPTEMBER 2022

**PHYSICAL SCIENCES P2
MARKING GUIDELINE/
FISIESE WETENSKAPPE V2
NASIENRIGLYN**

MARKS/ PUNTE: 150

This marking guideline consists of 19 pages./
Hierdie nasienriglyn bestaan uit 19 bladsye.

QUESTION/VRAAG 1

- 1.1 B ✓✓ (2)
- 1.2 A ✓✓ (2)
- 1.3 B ✓✓ (2)
- 1.4 C ✓✓ (2)
- 1.5 C ✓✓ (2)
- 1.6 A ✓✓ (2)
- 1.7 B ✓✓ (2)
- 1.8 D ✓✓ (2)
- 1.9 B ✓✓ (2)
- 1.10 A ✓✓ (2)
- [20]**

QUESTION/VRAAG 2

- 2.1 It is a series of organic compounds that can be described by the same general formula. ✓✓ (2 or 0)

’n Reeks organiese verbindings wat deur dieselfde algemene formule beskryf kan word. (2 of 0)

OR/OF


A series/group of organic compounds in which one member differs from the next with $-CH_2-$ group. ✓✓ (2 or 0)

’n Reeks organiese verbindings waarin die een lid van die volgende verskil met ’n CH_2 -groep. (2)

- 2.2.1 D ✓ (1)

- 2.2.2 C_nH_{2n-2} ✓ (1)

- 2.2.3



Marking criteria/ Nasienkriteria

- Only functional group correct / Slegs funksionele groep korrek: Max/ Maks ½
- Whole structure correct/ Hele struktuur korrek: 2/2

(2)

- 2.3 3-ethyl-2-methylhexanoic acid / 3-etiel-2-metielheksanoësuur

Marking criteria


- Correct stem i.e hexanoic acid ✓
- All substituents (ethyl and methyl) correctly identified ✓
- IUPAC name completely correct including numbering, sequence and hyphens ✓

Nasienkriteria

- Korrekte stam d.i heksanoësuur
- Alle substituentte (etiel en metiel) korrek geïdentifiseer
- IUPAC-naam heeltemal korrek insluitende nommering, volgorde en koppeltekens

(3)


- 2.4.1



Marking criteria/Nasienkriteria

- Longest chain contains 3 carbons / Langste koolstofketting bevat 3 koolstowwe ✓
- Two methyl substituents on C2 / Twee metielsubstituentte op C2 ✓
- Whole structure is correct / Hele struktuur korrek ✓

(3)



Marking criteria/Nasienkriteria

- Reactants / Reaktanse
- Products / Produkte
- Balancing / Balansering

(3)
[15]

QUESTION/VRAAG 3

3.1.1 **Marking criteria/Nasienkriteria**

If any of the underlined key words/phrases in the **correct context** are omitted: - 1 mark per word/phrase.

Indien enige van die sleutelwoorde/frases in die korrekte konteks weggelaat word: - 1 punt per woord/frase.

The temperature at which the vapour pressure of a liquid equals the atmospheric pressure. $\checkmark\checkmark$

Die temperatuur waarby die dampdruk van die vloeistof gelyk is aan die atmosferiese druk.

(2)

3.1.2 As the number of C atoms increases:

- The surface area/chain length/molecular mass of the alcohols increases $\checkmark$
- The strength of London forces/induced dipole forces/dispersion forces increase. $\checkmark$

Soos die aantal C-atome toeneem:

- *Die oppervlak-area/kettinglengte/molekulêre massa van die alkohole verhoog.*
- *Die sterkte van die Londonkragte/geïnduseerde dipoolkragte/verspreidingskragte verhoog*

OR/OF

As the number of C atoms decreases:

- The surface area/chain length/molecular mass of the alcohols decreases $\checkmark$
- The strength of London forces/induced dipole forces/dispersion forces decrease. $\checkmark$

Soos die aantal C-atome afneem:

- *Die oppervlak-area/kettinglengte/molekulêre massa van die alkohole verlaag.*
- *Die sterkte van die Londonkragte/geïnduseerde dipoolkragte/verspreidingskragte verswak*

(2)

3.1.3

Marking criteria

- Identify the intermolecular forces in both compounds. ✓✓
- Compare the strength of the intermolecular forces. ✓

Nasienkriteria

- Die intermolekulêre kragte korrek geïdentifiseer in beide verbindings
- Vergelyk die sterkte van die intermolekulêre kragte

- Alcohols have both (London forces) and hydrogen bonds ✓
- Ketones have both (London forces) and dipole-dipole forces ✓
- Hydrogen bonds in the alcohols are stronger than the dipole-dipole forces in ketones ✓
- *Alkohole het beide (Londonkragte) en waterstofbindings*
- *Ketone het beide (Londonkragte) en dipool-dipool kragte*
- *Waterstofbindings in die alkohole is sterker as die dipool-dipoolkragte in ketone*

OR/OF

- Alcohols have both (London forces) and hydrogen bonds ✓
- Ketones have both (London forces) and dipole-dipole forces ✓
- the dipole-dipole forces in Ketones are weaker than the hydrogen bonds in the alcohols ✓
- *Alkohole het beide (Londonkragte) en waterstofbindings*
- *Ketone het beide (Londonkragte) en dipool-dipool kragte*
- *Die dipool-dipoolkragte in ketone is swakker as die waterstofbindings in die alkohole*

(3)

3.1.4 To have one independent variable ✓ **OR** To have a fair test

Om slegs een onafhanklike veranderlike te het **OF** *Om 'n regverdige toets te hê*

(1)

3.1.5 Ketone ✓

Lower boiling point / Laer kookpunt ✓

(2)

3.2.1 Propanoic acid / Propanoësuur ✓

(1)

3.2.2

Marking criteria

- Identify the intermolecular forces correctly in both compounds. ✓
- Compare the strength of the intermolecular forces. ✓
- Compare the energy required to overcome the intermolecular forces. ✓

Nasienkriteria

- *Die intermolekulêre kragte is korrek in beide verbindings geïdentifiseer*
- *Vergelyk die sterkte van die intermolekulêre kragte.*
- *Vergelyk die energie wat benodig word om die intermolekulêre kragte te oorkom.*

- Both have hydrogen bonds
- Propan-1-ol has ONE site for hydrogen bonds
- Propanoic acid has TWO sites for hydrogen bonds } ✓
- The intermolecular forces of propanoic acid are stronger than that of propan-1-ol ✓
- More energy is needed to overcome the intermolecular forces of propanoic acid. ✓
- *Beide het waterstofbindings*
- *Propan-1-ol het EEN plek vir waterstofbindings*
- *Propanoësuur het TWEE plekke vir waterstofbindings*
- *Die intermolekulêrekragte in propanoësuur is sterker as dié in propan-1-ol*
- *Meer energie word benodig om die intermolekulêre kragte te oorkom in propanoësuur*

OR/OF

- Both have hydrogen bonds.
- Propan-1-ol has ONE site for hydrogen bonds
- Propanoic acid has two sites for hydrogen bonds } ✓
- The intermolecular forces of propan-1-ol are weaker than that of propanoic acid ✓
- Less energy is needed to overcome the intermolecular forces of propan-1-ol. ✓
- *Beide het waterstofbindings*
- *Propan-1-ol het EEN plek vir waterstofbindings*
- *Propanoësuur het TWEE plekke vir waterstofbindings*
- *Die intermolekulêrekragte in propan-1-ol is swakker as dié in propanoësuur.*
- *Minder energie word benodig om die intermolekulêre kragte te oorkom in propan-1-ol.*


(3)
[14]

QUESTION/VRAAG 4

4.1.1 Esterification / condensation / Esterifikasie / konsensasie ✓ (1)

4.1.2 (Mild) heat / (Matige) hitte ✓ (1)

4.1.3

**Marking criteria/ Nasienkriteria**

- Only functional group correct / Slegs funksionele groep korrek: Max/Maks ½
- Whole structure correct / Hele struktuur korrek: 2/2

(2)

4.1.4 Propyl ✓ ethanoate ✓ / Propiel etanoaat (2)

4.1.5 Pentanoic acid / Propanoësuur ✓✓ (2)

4.1.6 Substitution reaction / Substitusie-reaksie ✓ (1)

4.1.7 H₂O ✓ (1)

4.1.8 CH₃CH₂CH₂Cl ✓✓

Marking criteria/ Nasienkriteria


- Only functional group correct / Slegs funksionele groep korrek: Max/Maks ½
- Whole structure correct / Hele struktuur korrek: 2/2

(2)

4.2.1 (Concentrated / Gekonsentreerde) H₂SO₄ ✓ (2)

4.2.2 H₂O in excess ✓ / catalyst/ (Add small amount of HCl/H₃PO₄) (1)

4.2.3

**Marking criteria/ Nasienkriteria****(Organic molecules / Organiese molekules)**

- Only functional group correct / Slegs funksionele groep korrek: Max/Maks ½
- Whole structure correct / Hele struktuur korrek: 2/2

(5)

[20]

QUESTION/VRAAG 5

5.1

Marking criteria/ Nasienkriteria

If any of the underlined key words/phrases in the **correct context** are omitted: - 1 mark per word/phrase.

Indien enige van die sleutelwoorde/frases in die korrekte konteks

weggelaat word: - 1 punt per woord/frase

ANY ONE

- Change in concentration ✓ of reactant / product per (unit) time. ✓
- Change in amount/number of moles/volume/mass of products or reactants per (unit) time.
- Amount/number of moles/volume/mass of products formed/reactants used reactants per (unit) time.

ENIGE EEN

- Verandering in konsentrasie van reaktanse/produkte per (eenheid) tyd
- Verandering in hoeveelheid/getal mol/volume/massa van reaktanse of produkte per (eenheid) tyd.
- Hoeveelheid/getal mol/volume/massa van produkte gevorm/ reaktanse gebruik per (eenheid) tyd

OR/OF

The rate of change in concentration / amount of moles / number of moles / volume / mass. **(2 or 0).**

*Die tempo van verandering in konsentrasie / hoeveelheid mol / getal mol/volume/massa **(2 of 0)*** (2)

5.2 Concentration / Konsentrasie (of/van HCl) ✓ (1)

5.3 Equal to / Gelyk aan ✓

The same amount of (the limiting reagent), $\text{Na}_2\text{S}_2\text{O}_3$, is used. ✓
Dieselfde hoeveelheid (van beperkte reagens) $\text{Na}_2\text{S}_2\text{O}_3$ was gebruik. (2)

5.4.1 Experiment 3 / Eksperiment 3 ✓ (1)

5.4.2 For T_2

- Higher temperature increases kinetic energy of particles ✓
- Greater number of particles have sufficient energy. ✓
- More effective collision per unit time ✓

Vir T_2

- Hoër temperatuur verhoog die kinetiese energie van die deeltjies
- Groter aantal deeltjies het genoeg energie
- Meer effektiewe botsings per eenheidstyd

OR/OF

For T_1


- Lower temperature decreases kinetic energy of particles
- Fewer particles have sufficient energy.
- Less effective collision per unit time

Vir T_1

- Laer temperatuur verlaag die kinetiese energie van die deeltjies
- Minder aantal deeltjies het genoeg energie
- Minder effektiewe botsings per eenheidstyd

(3)

5.4.3

**Marking criteria/Nasienkriteria**

- $X < E_a$

(2)

5.5

Marking criteria

- Formula $n = m/M$
- Substitution into $n = m/M$
- **Using** ratio HCl: $\text{Na}_2\text{S}_2\text{O}_3$ 2 : 1
- Substitution into rate equation
- Final answer

Nasienkriteria

- *Formule* $n = m/M$
- *Vervanging in* $n = m/M$
- **Gebruik** van verhouding HCl: $\text{Na}_2\text{S}_2\text{O}_3$ 2 : 1
- *Vervanging in tempo vergelyking*
- *Finale antwoord*

$$n = \frac{m}{M} \checkmark$$

$$n = \frac{0,7118}{158} \checkmark$$

$$n(\text{Na}_2\text{S}_2\text{O}_3) = 4,505 \times 10^{-3} \text{ mol}$$

$$n(\text{HCl}) = 2 (4,505 \times 10^{-3}) \checkmark$$

$$n(\text{HCl}) = 9,01 \times 10^{-3} \text{ mol}$$

$$\text{rate/ tempo} = - \frac{\Delta n}{\Delta t}$$

$$\text{rate/ tempo} = - \frac{0 - 9,01 \times 10^{-3}}{34} \checkmark$$

$$\text{rate/ tempo} = 2,65 \times 10^{-4} (\text{mol} \cdot \text{s}^{-1}) \checkmark$$

Accept / Aanvaar

$$\text{rate/ tempo} = \frac{\Delta n}{\Delta t}$$

$$\text{rate/tempo} = \frac{- 9,01 \times 10^{-3}}{34} \checkmark$$

$$\text{rate / tempo} = - 2,65 \times 10^{-4} (\text{mol} \cdot \text{s}^{-1}) \checkmark$$

(5)

5.6

REMAINS THE SAME / BLY DIESELFDE ✓

(1)

[17]

QUESTION/VRAAG 6

- 6.1.1 (A reaction in which) products can be converted back to its reactants ✓✓
(and vice versa)

*(Is 'n reaksie waar) produkte terug na reaktanse, en omgekeerd,
omgeskakel kan word.*

(2 or/ of 0) (2)

- 6.1.2 Turns more pink / Raak meer pienk ✓ (1)

- 6.1.3 Turns more blue / Raak meer blou ✓ (1)

- 6.1.4 Exothermic / Eksotermies ✓ (1)

- 6.1.5 • Increase in temperature shifted the equilibrium position left ✓/Reverse
reaction is favoured

- Increase in temperature favours the endothermic reaction ✓

- *Toename in temperatuur verskuif die ewewigsposisie na links/
Terugwaartse reaksie word bevoordeel.*

- *Toename in temperatuur bevoordeel 'n endotermiese reaksie.* (2)

6.2

OPTION 1: MOLE CALCULATIONS**OPSIE 1: MOL BEREKENINGE****Marking criteria:**

- Substitution into formula $n = \frac{N}{N_A}$ ✓
- Using ratio $N_2O_4 : NO_2 = 1 : 2$ ✓
- $n(NO_2)$ equilibrium = $n_{initial} + \Delta n$ ✓
- $n(N_2O_4)$ equilibrium = $n_{initial} - \Delta n$ ✓
- Divide **equilibrium** amounts of N_2O_4 and NO_2 by 4 dm^3 ✓
- Correct K_c expression (formulae in square brackets) ✓
- Substitution into equilibrium concentration into K_c expression ✓
- Final answer ✓

Nasienkriteria:

- Vervanging in formule $n = \frac{N}{N_A}$
- Gebruik** verhouding $N_2O_4 : NO_2 = 1 : 2$ ✓
- $n(NO_2)$ ewewig = $n_{initial} + \Delta n$ ✓
- Deel **ewewig**hoeveelhede van N_2O_4 en NO_2 deur 4 dm^3
- Korrekte K_c -uitdrukking (formules met vierkanthakies)
- Vervanging in ewewigkonsentrasies in K_c -uitdrukking
- Finale antwoord

$$n = \frac{N}{N_A}$$

$$n = \frac{3,01 \times 10^{23}}{6,02 \times 10^{23}} \checkmark \quad (a)$$

$$n = 0,5 \text{ mol}$$

	N_2O_4 (g)	$2 NO_2$ (g)
Initial quantity (mol) <i>Aanvangshoeveelheid (mol)</i>	0,5	-
Change (mol) <i>Verandering (mol)</i>	0,4	0,8
Equilibrium (mol) <i>Ewewig (mol)</i>	0,1 ✓ (d)	0,8
Concentration ($\text{mol} \cdot \text{dm}^{-3}$) <i>Konsentrasie ($\text{mol} \cdot \text{dm}^{-3}$)</i>	0,025	0,2

✓ (b)
ratio

✓ (c)

✓ (e)

$K_c = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]}$ ✓ (f)	No K_c expression, correct substitution / <i>Geen K_c-uitdrukking, korrekte, korrekte substitusie.</i> Max / Maks 7/8
$K_c = \frac{(0,2)^2}{(0,025)}$ ✓ (g)	Wrong K_c expression/Verkeerde K_c – <i>uitdrukking.</i> Max. Maks. 5/8
$K_c = 1,6$ ✓ (h)	

OPTION 2: CONCENTRATION CALCULATIONS/**OPSIE 2: KONSENTRASIEBEREKENINGE****Marking criteria**

- Substitution into formula $n = \frac{N}{N_A}$ ✓
- Using ratio $\text{N}_2\text{O}_4 : \text{NO}_2 = 1 : 2$ ✓
- $c(\text{NO}_2)$ equilibrium = $c_{\text{initial}} + \Delta c$ ✓
- $c(\text{N}_2\text{O}_4)$ equilibrium = $c_{\text{initial}} - \Delta c$ ✓
- Divide n_{initial} and Δn of N_2O_4 by 4 dm^3 ✓
- Correct K_c expression (formulae in square brackets) ✓
- Substitution into equilibrium concentration into K_c expression ✓
- Final answer ✓

Nasienkriteria:

- Vervanging in formule $n = \frac{N}{N_A}$
- Gebruik** verhouding $\text{N}_2\text{O}_4 : \text{NO}_2 = 1 : 2$
- Ewewig $c(\text{NO}_2) = \text{begin } c + \Delta c$
- Ewewig $c(\text{N}_2\text{O}_4) = \text{begin } c - \Delta c$
- Deel **aanvangs en verandering** hoeveelhede van N_2O_4 en NO_2 deur 4 dm^3
- Korrekte K_c -uitdrukking (formules met vierkanthakies)
- Vervanging in ewewigskonsentrasies in K_c -uitdrukking
- Finale antwoord

✓(b)

✓(e)

$$n = \frac{N}{N_A}$$

$$n = \frac{3,01 \times 10^{23}}{6,02 \times 10^{23}} \checkmark \text{ (a)}$$

$$n = 0,5 \text{ mol}$$

	N ₂ O ₄ (g)	2 NO ₂ (g)
Initial concentration (mol·dm ⁻³) <i>Aanvangs konsentrasie (mol·dm⁻³)</i>	0,125	-
Change in concentration (mol·dm ⁻³) <i>Verandering in konsentrasie (mol·dm⁻³)</i>	0,1	0,2
Equilibrium concentration (mol·dm ⁻³) <i>Ewewig konsentrasie (mol·dm⁻³)</i>	0,025 ✓ (c)	0,2

$$K_c = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]} \checkmark \text{ (f)}$$

$$K_c = \frac{(0,2)^2}{(0,025)} \checkmark \text{ (g)}$$

$$K_c = 1,6 \checkmark \text{ (h)}$$

No K_c expression, correct substitution / *Geen K_c-uitdrukking, korrekte substitusie.*
Max / Maks 7/8

Wrong K_c expression / *Verkeerde K_c-uitdrukking.*
Max. Maks. 5/8

(8)
[15]

QUESTION/VRAAG 7

7.1.1 An acid is a proton (H^+ ion) donor / 'n Suur is 'n proton (H^+ -ioon) skenker ✓✓ (2)

7.1.2 HCl and/en Cl^- ✓✓ **OR/OF** H_3O^+ and/en H_2O ✓✓ (2)

7.1.3 Solution I. ✓

- HCl is a stronger acid than CH_3COOH / HCl has a higher K_a ✓ (than CH_3COOH)
- HCl will produce a higher concentration of H_3O^+ ✓ (than CH_3COOH)
- OR
- CH_3COOH is a weaker acid than HCl / CH_3COOH has a lower K_a (than HCl)
- CH_3COOH will produce a lower concentration of H_3O^+ (than HCl)

Oplossing I.

- HCl is 'n sterker suur as CH_3COOH / HCl het 'n hoër K_a -waarde as CH_3COOH
- HCl sal 'n hoër konsentrasie van H_3O^+ produseer as CH_3COOH
- OF
- CH_3COOH is 'n swakker suur as HCl / CH_3COOH het 'n laer K_a -waarde as HCl
- CH_3COOH sal 'n laer konsentrasie H_3O^+ produseer as HCl (3)

7.2.1

$ \begin{aligned} n &= cV \checkmark \\ &= 1 \times 10 / 1\,000 \checkmark \\ &= 0,01 \text{ mol } \checkmark \end{aligned} $

(3)

7.2.2

Marking criteria

- Formula $\text{pH} = -\log [\text{H}_3\text{O}^+]$ ✓
- pH value substituted into formula ✓
- Substitution in K_w formula ✓
- Substitution into $n = cV$ ✓
- Final answer ✓

Nasienkriteria

- Formule $\text{pH} = -\log [\text{H}_3\text{O}^+]$
- pH-waarde vervang in formule
- Vervanging in K_w formule
- Vervanging in $n = cV$
- Finale antwoord

Marking criteria

- Formula $\text{pOH} + \text{pH} = 14$ ✓
- pH value substituted into formula ✓
- Substitution in pOH formula ✓
- Substitution into $n = cV$ ✓
- Final answer ✓

Nasienkriteria

- Formule $\text{pOH} + \text{pH} = 14$
- pH-waarde vervang in formule
- Vervanging in pOH formule
- Vervanging in $n = cV$
- Finale antwoord

OPTION 1 / OPSIE 1

$$\text{pH} = -\log [\text{H}_3\text{O}^+] \checkmark$$

$$13 \checkmark = -\log [\text{H}_3\text{O}^+]$$

$$[\text{H}_3\text{O}^+] = 1 \times 10^{-13} \text{ mol} \cdot \text{dm}^{-3}$$

$$K_w = [\text{OH}^-][\text{H}_3\text{O}^+] = 1 \times 10^{-14}$$

$$[\text{OH}^-][\text{H}_3\text{O}^+] = 1 \times 10^{-14}$$

$$[\text{OH}^-](1 \times 10^{-13}) = 1 \times 10^{-14} \checkmark$$

$$[\text{OH}^-] = 0,1 \text{ mol} \cdot \text{dm}^{-3}$$

$$[\text{NaOH}] = 0,1 \text{ mol} \cdot \text{dm}^{-3}$$

$$c = \frac{n}{V}$$

$$0,1 = \frac{0,01}{V} \checkmark$$

$$V = 0,1 \text{ (dm}^3\text{)} \checkmark$$

OR/OF

From 7.2.1
Vanaf 7.2.1

OPTION 2 / OPSIE 2

$$\text{pOH} + \text{pH} = 14 \checkmark$$

$$\text{pOH} + 13 \checkmark = 14$$

$$\text{pOH} = 1$$

$$\text{pOH} = -\log [\text{OH}^-]$$

$$1 = -\log [\text{OH}^-] \checkmark$$

$$[\text{OH}^-] = 0,1 \text{ mol} \cdot \text{dm}^{-3}$$

$$[\text{NaOH}] = 0,1 \text{ mol} \cdot \text{dm}^{-3}$$

$$c_1V_1 = c_2V_2$$

$$(1)(10) = (0,1)V_2 \checkmark$$

$$V_2 = 100 \text{ cm}^3$$

$$V = 0,1 \text{ (dm}^3\text{)} \checkmark$$

(5)

7.2.3

Marking criteria - Formula $n = cV$ ✓ - Substitution of acid values into $n = cV$ ✓ AND Using ratio Acid : Base = 1 : 2 ✓ - Substitution of V and c into $n = cV$ for V base reacting ✓ - Subtracting $V_{\text{remaining}} = V_{\text{initial}} - V_{\text{reacting}} \quad \checkmark$ - Final answer ✓ Nasienkriteria - Formule $n = cV$ - Vervanging van suur waardes in formule $n = cV$ EN Gebruik verhouding Suur : Basis = 1 : 2 - Vervanging van V en c in $n = cV$ vir V basis wat reageer - Aftrekking $V_{\text{oorbly}} = V_{\text{aanvangs}} - V_{\text{reageer}}$ - Finale antwoord	Marking criteria / - Formula $\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$ ✓ - Substitution LHS $\frac{c_a V_a}{c_b V_b}$ ✓ - Substitution RHS $\frac{n_a}{n_b}$ ✓ - Subtracting $V_{\text{remaining}} = V_{\text{initial}} - V_{\text{reacting}} \quad \checkmark$ - Final answer ✓ Nasienkriteria - Formule $\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$ - Vervang LK $\frac{c_a V_a}{c_b V_b}$ - Vervang RK $\frac{n_a}{n_b}$ - Aftrekking $V_{\text{oorbly}} = V_{\text{aanvangs}} - V_{\text{reageer}}$ - Finale antwoord
OPTION 1/OPSIE 1 $\begin{aligned} n \text{ acid reacting} &= cV \checkmark \\ &= 0,09 \times 15/1\,000 \\ &= 1,35 \times 10^{-3} \text{ mol} \end{aligned} \quad \checkmark$ $\begin{aligned} n \text{ base reacting} &= 2 \times 1,35 \times 10^{-3} \text{ mol} \\ &= 2,7 \times 10^{-3} \text{ mol} \end{aligned}$ $n = cV$ $2,7 \times 10^{-3} = 0,1 V_{\text{base reacting / basis reageer}} \quad \checkmark$ $0,027 \text{ dm}^3 = V_{\text{base reacting / basis reageer}}$	OPTION 2/OPSIE 2 $\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b} \quad \checkmark$ $\frac{(0,09)(15)}{(0,1)V_b} \quad \checkmark = \frac{1}{2} \quad \checkmark$ $V_b = 27 \text{ cm}^3$ $V_b = 0,027 \text{ dm}^3$

$$\begin{aligned} V_{\text{remaining/oorbly}} &= 0,1 - 0,027 \quad \checkmark \\ &= 0,073 \text{ dm}^3 \quad \checkmark \end{aligned}$$

(5)
[20]

QUESTION/VRAAG 8

8.1 Loss of electrons / Verlies aan elektrone ✓✓ (2 orlof 0) (2)

8.2.1 $1 \text{ mol} \cdot \text{dm}^{-3}$ ✓ (1)

8.2.2 Platinum ✓ (1)

8.2.3 Cu ✓ (1)

8.2.4 $\text{O}_2 + 4\text{H}^+ + 4\text{e}^- \rightarrow 2 \text{H}_2\text{O}$ ✓✓

Marking criteria / Nasienkriteria

- $\text{O}_2 + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons 2 \text{H}_2\text{O}$ 1/2
- $2 \text{H}_2\text{O} \leftarrow \text{O}_2 + 4\text{H}^+ + 4\text{e}^-$ 2/2
- $2 \text{H}_2\text{O} \rightleftharpoons \text{O}_2 + 4\text{H}^+ + 4\text{e}^-$ 0/2
- $2 \text{H}_2\text{O} \rightarrow \text{O}_2 + 4\text{H}^+ + 4\text{e}^-$ 0/2

- Ignore if the charge omitted on electron / Ignoreer indien lading op elektron weggelaat is.

(2)

8.2.5 $2 \text{Cu} + \text{O}_2 + 4 \text{H}^+ \rightarrow 2 \text{Cu}^{2+} + 2 \text{H}_2\text{O}$ ✓ (✓ bal)

Marking criteria/Nasienkriteria

- Reactants/ Reaktanse
- Products / Produkte
- Balancing / Balansering

(3)

8.3.1 $E^\theta_{\text{cell}} = E^\theta_{\text{cathode/reduction/oxidising agent}} - E^\theta_{\text{anode/oxidation/reducing agent}}$ ✓

$$E^\theta_{\text{cell}} = (1,23) \checkmark - (0,34) \checkmark$$

$$E^\theta_{\text{cell}} = 0,89 \text{ V} \checkmark$$

Notes/Aantekeninge

- Any other formula using unconventional abbreviation, e.g.
 $E^\theta_{\text{cell}} = E^\theta_{\text{OA}} - E^\theta_{\text{RA}}$ followed by the correct substitution : ¾
- Enige ander formule wat onkonvensionele afkortings gebruik bv.
- $E^\theta_{\text{sel}} = E^\theta_{\text{OM}} - E^\theta_{\text{RM}}$ gevolg met korrekte vervangings: ¾

(4)

8.3.2 Concentration of the reactants decreases ✓
 Rate of the forward reaction decreases ✓

*Konsentrasie van reaktanse verlaag
 Tempo van voortwaartse reaksie verlaag*

(2)

8.3.3 Equilibrium / Ewewig ✓

(1)
[17]

QUESTION/VRAAG 9

9.1.1 **Marking criteria/ Nasienriglyne**


If any of the underlined key words/phrases in the **correct context** are omitted: - 1 mark per word/phrase.

Indien enige van die sleutelwoorde/frases in die korrekte konteks weggelaat word: - 1 punt per woord/frase

(It is a cell in which) electrical energy ✓ is converted into chemical energy ✓

(Dit is 'n sel waarin) elektriese energie omgeskakel word na chemiese energie.

(2)



Ignore phases / Ignoreer fases

(2)


Marking criteria / Nasienkriteria

- $2 \text{Cl}^- \rightleftharpoons \text{Cl}_2(\text{g}) + 2\text{e}^-$ 1/2
- $\text{Cl}_2(\text{g}) + 2\text{e}^- \leftarrow 2 \text{Cl}^-$ 2/2
- $\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2 \text{Cl}^-$ 0/2

Ignore if the charge omitted on electron / Ignoreer indien lading weggelaat is op elektron



(1)



H_2O is reduced to H_2 ✓

H_2O is 'n sterker oksideermiddel as Na^+

H_2O word gereduseer na H_2

(2)



(1)

9.2.2 $n_{\text{Cu}} = \frac{1}{2} \times 6$ ✓
 $= 3 \text{ mol}$

$m_{\text{Cu}} = nM = 3 \times 63,5$ ✓

$= 190,5 \text{ g}$

$0,95 \text{ } \checkmark m_{\text{IMPURE sample}} = 190,5$

$m_{\text{IMPURE sample}} = 200,53 \text{ g}$ ✓

Marking criteria

- Use of ratio of electrons to Cu
- Subst. into $n = m/M$
- Division by 0,95
- Final answer

Nasienkriteria

- *Gebruik van verhouding van elektrone tot Cu*
- *Vervanging in $n = m/M$*
- *Deel deur 0,95*
- *Finale antwoord*

(4)
[12]

TOTAL/TOTAAL: 150