

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great
Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



**SA EXAM
PAPERS**



KWAZULU-NATAL PROVINCE

**EDUCATION
REPUBLIC OF SOUTH AFRICA**

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

PHYSICAL SCIENCES P2 (CHEMISTRY)

COMMON TEST

JUNE 2023

MARKS : 150

TIME : 3 Hours

This question paper consists of 14 pages and 2 data sheets.


INSTRUCTIONS AND INFORMATION

1. Write your centre number and examination number in the appropriate spaces on the ANSWER BOOK.
2. This question paper consists of NINE questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two subquestions, for example between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. You may use appropriate mathematical instruments.
8. You are advised to use the attached DATA SHEETS.
9. Show ALL formulae and substitutions in ALL calculations.
10. Round off your final numerical answers to a minimum of TWO decimal places.
11. Give brief motivations, discussions et cetera where required.
12. Write neatly and legibly.

QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Choose the answer and write only the letter (A-D) next to the question number (1.1-1.10) in the ANSWER BOOK, for example 1.11 D.

- 1.1 What is the homologous series to which the following compound belongs?



- A Ketones
 B Alcohols
 C Haloalkanes
 D Carboxylic acids (2)

- 1.2 Which ONE of the following is a tertiary alcohol?

- A Pentan-1-ol
 B Pentan-2-ol
 C 2 – methylbutan – 2 - ol
 D 3 – methylbutan – 2 - ol (2)

- 1.3 Which of the following statements would apply to organic compounds that belong to the same homologous series? They have the same . . .

- I boiling points.
 II functional group.
 III molecular formula.


- A II only
 B I and II only
 C II and III only
 D I, II and III (2)

1.4 Which ONE of the following statements is TRUE for the reaction of an UNSATURATED HYDROCARBON with bromine water?




- A The reaction is slow.
- B The bromine water decolourises quickly.
- C The reaction is a substitution reaction.
- D Ultraviolet light is needed for this reaction.

(2)

1.5 The Maxwell–Boltzmann distribution curve for a reaction mixture is shown below:



The TEMPERATURE of the reaction mixture is now DECREASED. Which one of the following graphs shows the new distribution curve as a **dotted** line?




(2)


- 1.6 The mass of a catalyst was determined at intervals during a reaction.

Which ONE of the following graphs best represents the mass of the catalyst with time?


A




B



C






D



(2)

- 1.7 Each of the reactions represented below is at equilibrium in a closed container.

In which ONE of these reactions will an INCREASE IN PRESSURE (by decreasing the volume) at constant temperature favour the formation of products?



(2)

- 1.8 The expression for the equilibrium constant (K_c) for a hypothetical reaction is given as follows:

$$K_c = \frac{[R] [S]^2}{[P]^3}$$

Which ONE of the following balanced equations represents the hypothetical reaction?

- A $3P(s) \rightleftharpoons R(g) + 2S(g)$
- B $3P(l) \rightleftharpoons R(aq) + 2S(aq)$ ✓
- C $3P(aq) + Q(s) \rightleftharpoons R(g) + S_2(g)$
- D $3P(aq) + Q(s) \rightleftharpoons R(aq) + 2S(g)$ (2)

- 1.9 Which ONE of the following pairs represents the conjugate acid and the conjugate base of HPO_4^{2-} ?

	CONJUGATE ACID	CONJUGATE BASE
A	PO_4^{3-}	$H_2PO_4^-$
B	$H_2PO_4^-$	PO_4^{3-}
C	$H_2PO_4^-$	H_3PO_4
D	$H_2PO_4^{2-}$	PO_4^{2-}

(2)

- 1.10 CH_3COOH is a weak acid. Which ONE of the following statements is correct for a $0,01 \text{ mol} \cdot \text{dm}^{-3}$ aqueous solution of CH_3COOH ?

- A There are more hydronium ions than acetate ions.
- B There are fewer acetate ions than unionized CH_3COOH .
- C There are more hydronium ions than unionized CH_3COOH
- D There are equal quantities of unionized CH_3COOH , hydronium ions and acetate ions.

(2)
[20]

QUESTION 2 (Start on a new page.)

The letters **A** to **F** in the table below represent six organic compounds.

A	$\begin{array}{ccccccc} & \text{H} & & \text{CH}_3 & & \text{H} & & \text{H} \\ & & & & & & & \\ \text{H} & - \text{C} & - & \text{C} & - & \text{C} = & \text{C} & - & \text{C} & = & \text{C} & - & \text{H} \\ & & & & & & & \\ & \text{H} & & \text{CH}_3 & & \text{CH}_3 & & \text{H} \end{array}$		
B	$\begin{array}{ccccccc} & & & & & \text{H} & \\ & & & & & & \\ & & & & & \text{C} & - & \text{H} \\ & & & & & & \\ \text{H} & - & \text{C} & - & \text{C} & - & \text{C} & - & \text{O} & - & \text{H} \\ & & & & & & & & & & \\ & & \text{H} & & \text{H} & & \text{C} & - & \text{H} \\ & & & & & & & & \\ & & & & & & \text{H} & & \end{array}$		
C	$\text{CH}_3\text{CO}_2\text{H}$	D	$\text{CH}_3\text{CO}_2\text{C}_2\text{H}_5$
E	$\text{C}_3\text{H}_8\text{O}$	F	2 – bromobutane

Use the information in the table to answer the questions that follow.

2.1 Write down the IUPAC name of:

2.1.1 Compound A. (3)

2.1.2 Compound B. (3)

2.2 Write down the letter that represents:

2.2.1 A haloalkane. (1)

2.2.2 A compound that belongs to the same homologous series as compound B. (1)


- 2.3 Compound D is produced when compound C is reacted with a primary alcohol.
- 2.3.1 Define the term *functional group*. (2)
- 2.3.2 Write down the NAME of the FUNCTIONAL GROUP of the homologous series to which compound C belongs. (1)
- 2.3.3 Write down the structural formula of the functional group of the homologous series to which compound D belongs. (2)
- 2.3.4 Write a balanced equation using MOLECULAR FORMULAE to represent the reaction that takes place. (5)
- 2.3.5 Write down the NAME or FORMULA of the substance that can be used to accelerate the above reaction. (1)
- 2.3.6 Write down the IUPAC name of the primary alcohol used in this reaction. (2)
- 2.4 Define the term *functional isomer*. (2)
- 2.5 Write down the STRUCTURAL FORMULA of the straight chain FUNCTIONAL ISOMER of compound D. (2)
- [25]**

QUESTION 3 (Start on a new page.)

- 3.1 X and Y are two saturated hydrocarbons with the same molecular formula. Both X and Y have 4 carbons each. X has a lower boiling point than Y
- 3.1.1 Define *boiling point*. (2)
- 3.1.2 Fully explain the difference in the boiling points of these two compounds. (3)
- 3.1.3 Write down the STRUCTURAL FORMULA for compound X. (2)
- 3.2 In an experiment the boiling points of alcohols are compared to that of the carboxylic acids. The boiling point of pentan-1-ol is found to be 138 °C.
- 3.2.1 Write down the IUPAC name of the carboxylic acid that must be used in this experiment to ensure that the comparison is fair. Give a reason for the answer. (3)
- 3.2.2 Will the boiling point of the carboxylic acid be LESS THAN, EQUAL or GREATER THAN 138 °C? (1)
- 3.2.3 Fully explain the answer to QUESTION 3.2.2. (3)
- [14]**

QUESTION 4 (Start on a new page.)

In the flow diagram below, 1, 2, 3, and 4 represent organic reactions. P and Q represent organic compounds. Q is a HYDROCARBON.

**4.1 Consider REACTION 1:**

4.1.1 Name the type of reaction that takes place. Choose from SUBSTITUTION, ELIMINATION or ADDITION. (1)

4.1.2 Write down the structural formula of Q. (2)

4.1.3 Apart from heat, state one other reaction condition (2)

4.2 REACTION 2 is an example of an addition reaction.

4.2.1 Write down the TYPE of addition reaction represented here. (1)

4.2.2 Write down the NAME or FORMULA of the inorganic reactant required for this reaction. (1)

4.2.3 Name ONE reaction condition for reaction 2. (1)

4.3 Bromine is used as an inorganic reactant in REACTION 3.

4.3.1 Name the type of reaction that takes place. (1)

4.3.2 State the reaction condition for this reaction. (1)

4.3.3 Write down the structural formula of compound P. (2)

4.4 Consider REACTION 4:

4.4.1 Apart from heat, state one other reaction condition. (1)


4.4.2 Using condensed structural formula, write a balanced equation to represent the reaction that takes place. (3)

4.5 Compound P can be converted directly to compound Q in an elimination reaction. Write down the TYPE of elimination reaction taking place. (1)

[17]

QUESTION 5 (Start on a new page)

Calcium carbonate chips are added to excess dilute hydrochloric acid solution in a flask placed on a balance as illustrated below.



The cotton wool plug placed in the mouth of the flask prevents spillage of reactants and products, while simultaneously allowing the formed gas to escape. The balanced equation for the reaction that takes place is:



The mass of the flask and its contents is recorded in intervals of 30 seconds. The results obtained are shown in the table below.

Time in seconds (s)	Mass of the flask and its contents in grams (g)
0	217,50
30	215,00
60	212,95
90	211,05
120	209,20
150	207,40
180	X
210	203,95
240	203,95
270	Y

- 5.1 Define *reaction rate*. (2)
- 5.2 Write down the evidence from the table that suggests that the rate of the above reaction DECREASES with time. (2)
- 5.3 Use the collision theory to explain why the rate of the reaction decreases with time. Assume that the temperature remains constant. (3)

- 5.4 How will an increase in the volume of the hydrochloric acid used affect the rate of this reaction? Choose from INCREASES, DECREASES or REMAINS THE SAME (1)
- 5.5 The ~~average~~ rate of the reaction for the first 180s is $6,58 \times 10^{-2} \text{ g.s}^{-1}$. Calculate the value of X in the table. (4)
- 5.6 Write down the mass, in grams, represented by Y. Give a reason for the answer. (2)
- 5.7 Calculate ~~the~~ total volume of carbon dioxide gas produced during the above reaction at S.T.P. (6)
- 5.8 It is observed that when a catalyst is added to the above reaction, the reaction reaches completion in a shorter time.
Use the collision theory to explain the observation. (3)
- 5.9 The CaCO_3 chips are replaced with an equal mass of CaCO_3 powder. How will each of the following be affected by this change? (Choose from INCREASES, DECREASES or REMAINS THE SAME)
- 5.9.1 The mass in grams represented by Y. (1)
- 5.9.2 Time taken to reach the mass represented by X. (1)
- [25]**


QUESTION 6 (Start on a new page)

- 6.1 An investigation is performed to determine the effect of TEMPERATURE, PRESSURE and CONCENTRATION on equilibrium in the production of ammonia in a sealed container. The volume of the container is kept constant.

The balanced equation below represents the reaction that takes place in the sealed container.




The graph below represents the results obtained:



- 6.1.1 What information about the reaction does the graph provide between 0 and t₁? (2)
- 6.1.2 Will the K_c value for this reaction between 0 and t₁ be GREATER THAN 1, EQUAL TO 1 or LESS THAN 1? (1)
- 6.1.3 At which time, t₁, t₂ or t₃ was the concentration of a reactant increased? Give a reason for the answer. (2)
- 6.1.4 State Le Chatelier's principle. (2)
- 6.1.5 Which ONE of the factors, TEMPERATURE, Pressure or CONCENTRATION was changed at t₃? (1)
- 6.1.6 Was the factor identified in QUESTION 6.1.5 INCREASED or DECREASED? Explain the answer by referring to Le Chatelier's principle. (4)


- 6.1.7 A partial graph of reaction rate versus time is drawn below. The dotted line represents the reverse reaction



Copy the above graph in your answer book. Complete the graph for each of the time periods period till t_4 . Use the solid line for the forward reaction rate and the dotted line for the reverse reaction rate.

(5)

- 6.2 The equation below represents the reversible reaction that takes place when sulphur dioxide gas, SO_2 , and oxygen gas, O_2 , are allowed to react in a sealed container.



- 6.2.1 Write down the meaning of the term *reversible reaction*. (2)

- 6.2.2 The activation energy for the FORWARD reaction is compared to the activation energy of the REVERSE reaction. Which reaction, FORWARD or REVERSE, will have a higher activation energy? (1)

- 6.3 A hypothetical reaction is represented by the balanced equation below:



2,5 moles of X (g) were mixed with 1,75 moles of Y (g) in a sealed 2 dm^3 container. The reaction reaches equilibrium at $T^\circ\text{C}$. At equilibrium the concentration of C (g) was $0,5 \text{ mol.dm}^{-3}$.

Calculate the value of the equilibrium constant, K_c .

(7)

[27]

QUESTION 7 (Start on a new page.)

7.1 Sulphuric acid, H_2SO_4 , ionizes in water in TWO steps as follows:




7.1.1 Define an acid in terms of the Arrhenius theory. (2)

7.1.2 Write down the NAME or FORMULA of the substance that acts as an ampholyte in the above equations. Give a reason for the answer. (2)

7.2 The K_a value for TWO acids are as follows:

NAME OF ACID	FORMULA OF ACID	K_a VALUE
Oxalic Acid	$(\text{COOH})_2$	$5,6 \times 10^{-2}$
Carbonic Acid	H_2CO_3	$4,3 \times 10^{-2}$

7.2.1 Oxalic acid ionizes in water according to the following balanced equation:



Write down the formula of the TWO BASES in this reaction. (2)

7.2.2 Which ONE, oxalic acid or carbonic acid, will have a higher conductivity? Explain the answer by referring to the information in the table. (3)

7.3 Water is added to a $0,01 \text{ mol} \cdot \text{dm}^{-3}$ solution of hydrochloric acid. How does the addition of the water affect each of the following? (Choose from: INCREASES, DECREASES or REMAINS THE SAME).

7.3.1 The strength of the solution of the hydrochloric acid. (1)

7.3.2 The concentration of the solution of the hydrochloric acid. Give a reason for the answer. (2)

7.4 A solution of sodium hydrogen carbonate, (NaHCO_3) of volume 250 cm^3 , is prepared by dissolving x g of sodium hydrogen carbonate in sufficient distilled water in a volumetric flask.

20 cm^3 of the prepared sodium hydrogen carbonate solution neutralises $12,58 \text{ cm}^3$ of a sulphuric acid (H_2SO_4) solution of $0,29 \text{ mol} \cdot \text{dm}^{-3}$.

The balanced equation for the reaction is:



7.4.1 Define the equivalence point of a titration. (2)

7.4.2 Calculate the concentration of the NaHCO_3 solution that neutralised the H_2SO_4 solution. (4)

7.4.3 Calculate x, the mass of sodium hydrogen carbonate used to prepare the initial solution of sodium hydrogen carbonate. (4)

[22]
[150]

**DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)**

TABLE 1: PHYSICAL CONSTANTS

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure <i>Standaarddruk</i>	p^θ	$1,013 \times 10^5 \text{ Pa}$
Molar gas volume at STP <i>Molêre gasvolume by STD</i>	V_m	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard temperature <i>Standaardtemperatuur</i>	T^θ	273 K
Charge on electron <i>Lading op electron</i>	E	$-1,6 \times 10^{-19} \text{ C}$
Avogadro's constant <i>Avogadro-konstante</i>	N_A	$6,02 \times 10^{23} \text{ mol}^{-1}$

TABLE 2: FORMULAE

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ or/of $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$
$\frac{c_a v_a}{c_b v_b} = \frac{n_a}{n_b}$	$\text{pH} = -\log[\text{H}_3\text{O}^+]$
$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14} \text{ at/by } 298 \text{ K}$	

KEYISLEUTEL

Copyright reserved