

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



SA EXAM PAPERS

SA EXAM
PAPERS



LIMPOPO
PROVINCIAL GOVERNMENT
REPUBLIC OF SOUTH AFRICA

DEPARTMENT OF
EDUCATION

NATIONAL
SENIOR CERTIFICATE

GRADE/*GRAAD* 12



MATHEMATICS P1/*WISKUNDE VI*
SEPTEMBER 2023
MARKING GUIDELINES/*NASIENRIGLYNE*

MARKS/*PUNTE*: 150

These marking guidelines consist of 21 pages/*Hierdie nasienriglyne bestaan uit 21 bladsye.*



QUESTION/VRAAG 1

1.1	1.1.1	$x(7-x)=0$ $x=0$ or/of $x=7$	$\checkmark x=0$ $\checkmark x=7$	(2)
	1.1.2	$3x^2 - 2x - 6 = 0$ $x = \frac{-(-2) \pm \sqrt{(-2)^2 - 4(3)(-6)}}{2(3)}$ $x = -1,12$ or/of $x = 1,79$	$\checkmark$ subst in correct formula/vervang in korrekte formule $\checkmark x = -1,12$, $\checkmark x = 1,79$ N.B. (-1 mark for incorrect rounding/-1 vir verkeerde afronding)	(3)
	1.1.3	$(3-x)(x+7) < 0$ CV: $x=3$ or/of $x=-7$  $\therefore x < -7$ or/of $x > 3$ OR/OF $(-\infty; -7)$ or/of $(3; \infty)$ OR/OF $(3-x)(x+7) < 0$ $-(x-3)(x+7) < 0$ $(x-3)(x+7) > 0$ CV: $x = -7$ or/of $x = 3$  $\therefore x < -7$ or/of $x > 3$ OR/OF $(-\infty; -7)$ or/of $(3; \infty)$	$\checkmark$ critical values/ kritieke waardes $\checkmark \checkmark$ answer/antwoord OR/OF $\checkmark$ critical values/ kritieke waardes $\checkmark \checkmark$ answer/antwoord	(3)



1.1.4	$\sqrt[3]{32} = 8^{3x} \cdot 2^{6x}$ $32^{\frac{1}{3}} = 8^{3x} \cdot 4^{3x}$ $32^{\frac{1}{3}} = (8 \times 4)^{3x}$ $32^{\frac{1}{3}} = 32^{3x}$ $3x = \frac{1}{3}$ $x = \frac{1}{9}$ OR / OF $\sqrt[3]{32} = 8^{3x} \cdot 2^{6x}$ $2^{\frac{5}{3}} = (2^3)^{3x} \cdot 2^{6x}$ $2^{\frac{5}{3}} = 2^{9x} \cdot 2^{6x}$ $2^{\frac{5}{3}} = 2^{15x}$ $15x = \frac{5}{3}$ $x = \frac{1}{9}$	$\checkmark 32^{\frac{1}{3}} = 8^{3x} \cdot 4^{3x}$ $\checkmark 32^{\frac{1}{3}} = 32^{3x}$ $\checkmark 3x = \frac{1}{3}$ $\checkmark x = \frac{1}{9}$ OR/OF $\checkmark 2^{\frac{5}{3}} = 2^{9x} \cdot 2^{6x}$ $\checkmark 2^{\frac{5}{3}} = 2^{15x}$ $\checkmark 15x = \frac{5}{3}$ $\checkmark x = \frac{1}{9}$	(4)
1.1.5	$x - 4 - 2\sqrt{x-1} = 0$ $x - 4 = 2\sqrt{x-1}$ $(x-4)^2 = (2\sqrt{x-1})^2$ $x^2 - 8x + 16 = 4(x-1)$ $x^2 - 12x + 20 = 0$ $(x-10)(x-2) = 0$ $x = 10$ or/of $x \neq 2$	$\checkmark$ isolating the square root/soleer vierkants wortel $\checkmark$ squaring both sides/kwadreer beide kante $\checkmark$ standard form/vorm $\checkmark x = 10$ or/of $x \neq 2$	(4)



1.2	$x - 2y = 1$ $x = 1 + 2y \dots\dots\dots(1)$ $x^2 - xy + 2y^2 = 2 \dots\dots\dots(2)$ subst (1) into (2) $(1 + 2y)^2 - y(1 + 2y) + 2y^2 = 2$ $4y^2 + 4y + 1 - y - 2y^2 + 2y^2 - 2 = 0$ $4y^2 + 3y - 1 = 0$ $(4y - 1)(y + 1) = 0$ $y = \frac{1}{4}$ or/of $y = -1$ $x = \frac{3}{2}$ or/of $x = -1$ OR/OFF $y = \frac{x-1}{2} \dots\dots\dots(1)$ $x^2 - xy + 2y^2 = 2 \dots\dots\dots(2)$ Subst. (1) into (2) $x^2 - x\left(\frac{x-1}{2}\right) + 2\left(\frac{x-1}{2}\right)^2 = 2$ $x^2 - \frac{x^2-x}{2} + \frac{2x^2-4x+2}{4} = 2$ $4x^2 - 2x^2 + 2x + 2x^2 - 4x + 2 - 8 = 0$ $4x^2 - 2x - 6 = 0$ $2x^2 - x - 3 = 0$ $(2x-3)(x+1) = 0$ $x = \frac{3}{2}$ or/of $x = -1$ $y = \frac{1}{4}$ or/of $y = -1$	✓ equation/vgl 1 ✓ substitution/ <i>vervang</i> ✓ simpl/ <i>vereenv</i> ✓ standard form/ <i>vorm</i> ✓ y-values/ <i>waardes</i> ✓ x-values/ <i>waardes</i> OR/OFF ✓ equation/vgl 1 ✓ substitution/ <i>vervang</i> ✓ simpl/ <i>vereenv</i> ✓ standard form/ <i>vorm</i> ✓ x-values/ <i>waardes</i> ✓ y-values/ <i>waardes</i>	(6)
-----	--	--	-----

1.3	1.3.1	Then the amount that should be paid by each friend $= \frac{3500}{x}$ $\therefore$ The amount paid by each friend less four friends $= \frac{3500}{x-4}$ Extra amount paid = 43,75 $\rightarrow \frac{3500}{x-4} - \frac{3500}{x} = 43,75$ $43,75x^2 - 175x - 14000 = 0$ $x^2 - 4x - 320 = 0$ $(x-20)(x+16) = 0$ $x = 20$ or $x \neq -16$ $\rightarrow 20$ friends are going on the trip. OR/OF Let the cost per person be y $y = \frac{3\ 500}{x}$ $xy = 3\ 500$ $(x-4)(y+43,75) = 3\ 500$ $xy - 4y + 43,75x - 175 = 3\ 500$ $3\ 500 - 4\left(\frac{3\ 500}{x}\right) + 43,75x - 175 = 3\ 500$ $43,75x^2 - 175x - 14\ 000 = 0$ $x^2 - 4x - 320 = 0$ $(x-20)(x+16) = 0$ $\therefore x = 20$ or $x \neq -16$	$\checkmark \frac{3500}{x}$ $\checkmark \frac{3500}{x-4}$ $\checkmark$ equation/vgl $\checkmark$ standard form/vorm $\checkmark$ values with selection/ waardes net seleksie OR/OF $\checkmark y = \frac{3\ 500}{x}$ $\checkmark$ equation/vgl $\checkmark$ simplification/ vereenvoudig $\checkmark$ standard form/vorm $\checkmark$ values with selection/ waardes met seleksie	(5)
	1.3.2	Each friend pays: $\frac{3500}{16} = R218,75$ OR/OF Cost = $175 + 43,75$ $= R218,75$	$\checkmark$ answer/antwoord OR/OF $\checkmark R218,75$	(1)
				(1)
				[28]



		OR/OF $21 + (n-2)(14) + 21 + (n-1)(14) = 308$ $21 + 14n - 28 + 21 + 14n - 14 = 308$ $28n = 308$ $n = 11$ $T_{11} = 21 + 10(14) = 161$ $T_{10} = 21 + 9(14) = 147$	OR/OF ✓ $d = 14$ ✓ method/metode ✓ $n = 11$ ✓ $T_{10} = 147$	(4)
2.2		$\begin{array}{ccccccc} a+b+c & & 4a+2b+c & & 9a+3b+c & & 68 \\ & \swarrow & & \swarrow & & \swarrow & \\ & & 3a+b & & 21 & & 29 \\ & & & \swarrow & & \swarrow & \\ & & & & 8 & & 8 \end{array}$ $2a = 8$ $a = 4$ $5a + b = 21$ $5(4) + b = 21$ $b = 1$ $16a + 4b + c = 68$ $16(4) + 4(1) + c = 68$ $c = 0$ $\therefore T_n = 4n^2 + n$	✓ 2nd difference/ 2^e verskil ✓ $2a = 8$ ✓ $5a + b = 21$ ✓ $16a + 4b + c = 68$	(4)
				[13]



QUESTION/VRAAG 3

3.1	3.1.1	$T_1 = a^2$ $T_2 = (ar)^2$ OR / OF $T_2 = a^2 r^2$	✓ answer/antwoord	(1)
	3.1.2	$\frac{a}{1-r} = 6 \dots\dots\dots(1)$ $\frac{a^2}{1-r^2} = 6 \dots\dots\dots(2)$ $\frac{a^2}{1-r^2} = \left(\frac{a}{1-r}\right)\left(\frac{a}{1+r}\right)$ $6 = 6\left(\frac{a}{1+r}\right)$ $a = 1+r \dots\dots\dots(3)$ subst. into (1) $\frac{1+r}{1-r} = 6$ $1+r = 6-6r$ $r = \frac{5}{7}$ $a = \frac{12}{7}$	✓ both equations/ beide vergelykings ✓ $6 = 6\left(\frac{a}{1+r}\right)$ ✓ equation/vgl (3) ✓ substitution/ vervanging ✓ value/waarde $r = \frac{5}{7}$ ✓ value/waarde $a = \frac{12}{7}$	(6)



		OR/OF $\frac{a}{1-r} = 6$ $a = 6(1-r) \dots\dots\dots(1)$ $\frac{a^2}{1-r^2} = 6$ $a^2 = 6(1-r^2) \dots\dots\dots(2)$ From equation (1) $a^2 = 36(1-r)^2$ $\therefore 36(1-r)^2 = 6(1-r^2)$ $36 - 72r + 36r^2 = 6 - 6r^2$ $42r^2 - 72r + 30 = 0$ $7r^2 - 12r + 5 = 0$ $(7r-5)(r-1) = 0$ $r = \frac{5}{7} \text{ or } r \neq 1$ $a = 6\left(1 - \frac{5}{7}\right)$ $= \frac{12}{7}$	OR/OF $\checkmark a = 6(1-r)$ $\checkmark a^2 = 6(1-r^2)$ $\checkmark 36(1-r)^2 = 6(1-r^2)$ $\checkmark 42r^2 - 72r + 30 = 0$ $\checkmark r = \frac{5}{7} \text{ or / of } r \neq 1$ $\checkmark a = \frac{12}{7}$	(6)
	3.1.3	$T_6 = a^2 r^{10}$ $= \left(\frac{12}{7}\right)^2 \left(\frac{5}{7}\right)^{10}$ $= 0,10$ OR/OF $T_6 = (ar^5)^2$ $= \left[\frac{12}{7} \left(\frac{5}{7}\right)^5\right]^2$ $= \left(\frac{12}{7}\right)^2 \left(\frac{5}{7}\right)^{10}$ $= 0,10$	$\checkmark \left(\frac{12}{7}\right)^2 \left(\frac{5}{7}\right)^{10}$ $\checkmark \text{ answer/antwoord}$ OR/OF $\checkmark \left(\frac{12}{7}\right)^2 \left(\frac{5}{7}\right)^{10}$ OR/OF $\left[\frac{12}{7} \left(\frac{5}{7}\right)^5\right]^2$ $\checkmark \text{ answer/antwoord}$	(2)



3.2	water left after 1 hour = $\frac{99}{100} \times 50$ $\therefore a = 50(0,99), \quad r = 0,99$ $T_n = ar^{n-1}$ $T_8 = 50(0,99)(0,99)^7$ $= 50(0,99)^8$ $= 46,14$ 46,14 litres of water will be left after eight hours OR/OF water evaporated after an hour = 1% of 50 = 0,5 water evaporated after 2 hours = 1% of 49,5 = 0,495 water evaporated after 3 hours = 1% of 49,005 = 0,49005 $a = 0,5 \quad r = 0,99$ $S_n = \frac{a(r^n - 1)}{r - 1}$ $S_8 = \frac{0,5(0,99^8 - 1)}{0,99 - 1}$ $= 3,86$ $\therefore$ water that will be left after eight hours = $50 - 3,86$ $= 46,14$ liters OR/OF $T_0 = 50$ $T_1 = 50(0,99)$ $T_2 = 50(0,99)(0,99)$ $T_8 = 50(0,99)(0,99)^7$ $= 50(0,99)^8$ $= 46,14$ litres	✓ $a = 50(0,99)$ ✓ subst in correct formula/vervang in korrekte formule ✓ 46,14 litres OR/OF ✓ $a = 0,5$ ✓ subst in correct formula/vervang in korrekte formule ✓ 46,14 litres OR/OF ✓ $T_1 = 50(0,99)$ ✓ substitution/vervanging ✓ answer/antwoord	(3) (3) (3)
			[12]



QUESTION/VRAAG 4

4.1	4.1.1	$p = -1$ $q = -3$	✓ value/waarde p ✓ value/waarde q	(2)
	4.1.2	$f(x) = \frac{a}{x-1} - 3$ subs (0 ; -5) $-5 = \frac{a}{0-1} - 3$ $-5 = -a - 3$ $a = 2$	✓ subst. p and q / verv. p en q ✓ subst/verv (0 ; -5) ✓ value/waarde a	(3)
	4.1.3	f is translated 6 units up and 2 units to the right to form g .	✓ 6 units up/ eenhede op ✓ 2 units to the right/ eenhede na regs	(2)
	4.1.4	$y = 3$	✓ answer/antwoord	(1)
4.2	4.2.1	$x > 0$	✓ answer/antwoord	(1)
	4.2.2	$y = \log_b x$ $b^y = x$ $b^{-2} = 25$ $b = \frac{1}{5}$ $f(x) = \log_{\frac{1}{5}} x$	✓ subst/verv (25 ; -2) ✓ answer/antwoord	(2)
	4.2.3	$y = \log_{\frac{1}{5}} x$ $x = \log_{\frac{1}{5}} y$ $y = \left(\frac{1}{5}\right)^x$	✓ swop x and y /ruil x en y ✓ answer/antwoord	(2)
	4.2.4	$x > -5$	✓✓ answer/antwoord	(2)
				[15]



QUESTION/VRAAG 5

5.1		$y = a(x+1)(x+5)$ $5 = a(1)(5)$ $a = 1$ $y = (x+1)(x+5)$ $y = x^2 + 6x + 5$	✓ subst/verv $(-5 ; 0)$ & $(-1 ; 0)$ ✓ subst/verv $(0 ; 5)$ ✓ value/waarde a	(3)
5.2		$x^2 + 6x + 5 = 2x + 10$ $x^2 + 4x - 5 = 0$ $(x+5)(x-1) = 0$ $x \neq -5$ or $x = 1$ $\therefore y = 2(1) + 10 = 12$ OR $y = (1)^2 + 6 + 5 = 12$ $Q(1 ; 12)$	✓ equating/gelykstel ✓ standard form/vorm ✓ x-values with selection/x-waardes met seleksie ✓ y-value/waarde	(4)
5.3		For turning point: $f'(x) = 2x + 6 = 0$ $x = -3$ Then $f(-3) = (-3)^2 + 6(-3) + 5 = -4$ -4 is minimum value. Therefore, $f(x) \neq -5$ for all value of x OR/OF Axis of symmetry: $x = -\frac{b}{2a} = -\frac{6}{2} = -3$ $y = (-3)^2 + 6(-3) + 5 = -4$ $f(x) = -4$ is minimum at $x = -3$ $-5 < -4$, therefore, $(x) \neq -5$ for all x	✓ $x = -3$ ✓ value/waarde y ✓ conclusion/gevolgtrekking OR/OF ✓ x-value//waardes ✓ value/waarde y ✓ conclusion/gevolgtrekking	(3) (3)
5.4	5.4.1	$SR = 2x + 10 - (x^2 + 6x + 5)$ $= -x^2 - 4x + 5$ $\frac{dSR}{dx} = -2x - 4 = 0$ $x = -2$ $f'(x) = 2x + 6$ $\therefore f'(-2) = 2$	✓ SR ✓ value/waarde x ✓ $f'(x)$ ✓ $f'(-2)$	(4)



		OR / OF $SR = 2x + 10 - (x^2 + 6x + 5)$ $= -x^2 - 4x + 5$ $\max SR \text{ at } x = \frac{-(-4)}{2(-1)}$ $= -2$ $f'(x) = 2x + 6$ $\therefore f'(-2) = 2$	OR/OF ✓ $-x^2 - 4x + 5$ ✓ $x = -2$ ✓ $f'(x) = 2x + 6$ ✓ $f'(-2) = 2$	(4)
	5.4.2	When $x = -2$ $y = x^2 + 6x + 5$ $y = (-2)^2 + 6(-2) + 5 = -3$ $R(-2; -3)$ and $m = 2$ $y = mx + c$ $-3 = 2(-2) + c$ $c = 1$ $y = 2x + 1$ OR / OF When $x = -2$ $y = x^2 + 6x + 5$ $y = (-2)^2 + 6(-2) + 5 = -3$ $R(-2; -3)$ and $m = 2$ $y - (-3) = 2[x - (-2)]$ $y + 3 = 2x + 4$ $y = 2x + 1$	✓ y-value/waarde ✓ substitution/ vervanging ✓ answer/antwoord OR/OF ✓ y-value//waarde ✓ substitution/ vervanging ✓ answer/antwoord	(3)
5.5		$g(x) = 2x + 10$ $g^{-1}(x) = \frac{x}{2} - \frac{10}{2}$ $g(x) - g^{-1}(x) = 15$ for $x = 15$ $\therefore g(x) - g^{-1}(x) > 15$ for $x > 0$	✓ $g^{-1}(x)$ ✓✓ $x > 0$	(3)
				[20]



QUESTION/VRAAG 6

6.1		$1 + i_{eff} = 1 + i_{nom}$ $1 + i_{eff} = \left(1 + \frac{0,05}{12}\right)^{12}$ $i_{eff} = 0,05116...$ $= 5,12\%$	✓ substitution/ vervanging ✓ answer/antwoord	(2)
6.2	6.2.1	$P = \frac{x \left[1 - (1+i)^{-n} \right]}{i}$ $800\,000 = \frac{x \left[1 - \left(1 + \frac{0,1075}{12}\right)^{-300} \right]}{\frac{0,1075}{12}}$ $x = 7\,696,74$ Matome's monthly payments = R7 696,74	✓ value/waarde n ✓ value/waardes i ✓ subst in correct formula/vervang in korrekte formule ✓ answer/antwoord	(4)
	6.2.2	Balance after 240th payment: $\text{Balance} = x(1+i)^n - \frac{x \left[(1+i)^n - 1 \right]}{i}$ $= 800\,000 \left(1 + \frac{0,1075}{12}\right)^{240} - \frac{7\,696,74 \left[\left(1 + \frac{0,1075}{12}\right)^{240} - 1 \right]}{\frac{0,1075}{12}}$ $= 356\,036,20$ OR/OF $P = \frac{x \left[1 - (1+i)^{-n} \right]}{i}$ $= \frac{7\,696,74 \left[1 - \left(1 + \frac{0,1075}{12}\right)^{-60} \right]}{\frac{0,1075}{12}}$ $= 356\,034,62$ Balance was R356 034, 62	✓✓ Tick for each term/ merk elke term ✓ answer/antwoord OR/OF ✓ $n = -60$ ✓ subst in correct formula/vervang in korrekte formule ✓ answer/antwoord	(3)



6.2.3	The balance after missing 6 instalments: $A = P(1+i)^n$ $= 356\,034,62 \left(1 + \frac{0,1075}{12}\right)^6$ $= 375\,605,22$ The number of payments: $375\,605,22 = \frac{9\,000 \left[1 - \left(1 + \frac{0,1075}{12}\right)^{-n}\right]}{\frac{0,1075}{12}}$ $-n = \log_{\left(1 + \frac{0,1075}{12}\right)} \left[1 - \frac{375\,605,22 \left(\frac{0,1075}{12}\right)}{9\,000}\right]$ $n = 52,5$ Yes, Matome was able to settle the loan in time. He settled the loan in 53 months instead of 54 months. OR/OR $A = P(1+i)^n$ $= 356\,034,62 \left(1 + \frac{0,1075}{12}\right)^6$ $= 375\,605,2205$ $P = \frac{x \left[1 - (1+i)^{-n}\right]}{i}$ $375\,605,2205 = \frac{x \left[1 - \left(1 + \frac{0,1075}{12}\right)^{-54}\right]}{\frac{0,1075}{12}}$ $x = 8\,803,68$ Yes, Matome paid more than the required amount.	✓ subst in correct formula/vervang in korrekte formule ✓ answer/antwoord ✓ subst in correct formula/vervang in korrekte formule ✓ value/waarde n ✓ conclusion/ OR/OF ✓ subst in correct formula/vervang in korrekte formule ✓ answer/antwoord ✓ subst in correct formula/vervang in korrekte formule ✓ answer/antwoord ✓ conclusion/ gevolgtrekking gevolgtrekking	(5) (5)
			[14]




QUESTION/VRAAG 7

7.1		$f(x) = 4x^2 - x$ $f(x+h) = 4(x+h)^2 - (x+h)$ $= 4x^2 + 8xh + 4h^2 - x - h$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{4x^2 + 8xh + 4h^2 - x - h - (4x^2 - x)}{h}$ $= \lim_{h \rightarrow 0} \frac{8xh + 4h^2 - h}{h}$ $= \lim_{h \rightarrow 0} \frac{h(8x + 4h - 1)}{h}$ $= \lim_{h \rightarrow 0} (8x + 4h - 1)$ $= 8x - 1$ OR/OF $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{4(x+h)^2 - (x+h) - (4x^2 - x)}{h}$ $= \lim_{h \rightarrow 0} \frac{4x^2 + 8xh + 4h^2 - x - h - 4x^2 + x}{h}$ $= \lim_{h \rightarrow 0} \frac{8xh + 4h^2 - h}{h}$ $= \lim_{h \rightarrow 0} \frac{h(8x + 4h - 1)}{h}$ $= \lim_{h \rightarrow 0} (8x + 4h - 1)$ $= 8x - 1$	✓ expansion/uitbrei ✓ correct substitution/ korrekte vervanging ✓ simplification/ vereenvoudig ✓ common factor/ gemeensk faktor ✓ answer/antwoord OR/OF ✓ correct substitution ✓ expansion ✓ simplification/ vereenvoudig ✓ common factor ✓ answer/antwoord	(5)
7.2	7.2.1	$y = -x^3 - 16x^2 + 6x$ $\frac{dy}{dx} = -3x^2 - 32x + 6$	✓ $-3x^2$ ✓ $-32x$ ✓ 6	(3)



7.2.2	$f(x) = \frac{4}{5x^{-3}} - \frac{3}{\sqrt[3]{x^2}}$ $= \frac{4}{5}x^3 - 3x^{\frac{2}{3}}$ $f'(x) = \frac{12}{5}x^2 + 2x^{\frac{5}{3}}$	$\checkmark \frac{4}{5}x^3$ $\checkmark -3x^{\frac{2}{3}}$ $\checkmark \frac{12}{5}x^2$ $\checkmark 2x^{\frac{5}{3}}$	(4)
			[12]

QUESTION/VRAAG 8

8.1		$\checkmark$ x-intercepts/afsnitte $\checkmark$ y- intercept/afsnit $\checkmark$ turning points/DP $\checkmark$ shape/vorm	(4)
8.2	$y = a(x-1)^2(x+2)$ $2 = a(-1)^2(2)$ $2 = 2a$ $a = 1$ $f(x) = (x^2 + x - 2)(x - 1)$ $= x^3 - 3x + 2$	$\checkmark$ subst in correct formula/vervang in korrekte formule $\checkmark$ subst/verv (0 ; 2) $\checkmark a = 1$ $\checkmark b = 0$ $\checkmark c = -3 \text{ \& } d = 2$	(5)



8.3		(1 ; 0) and/en (4 ; 0)	✓ (1 ; 0) ✓ (4 ; 0)	(2)
8.4		$p > 4$ or/of $p < 0$	✓✓ answer/antwoord	(2)
				[13]

QUESTION/VRAAG 9

9.1		The area of the square = x^2 and the area of the circle = πr^2 Total area (A) = $x^2 + \pi r^2$(1) Perimeter of the square + circumference of the circle = 12 $4x + 2\pi r = 12$ $r = \frac{6-2x}{\pi}$(2) subst. (2) into (1) $A = x^2 + \pi \left(\frac{6-2x}{\pi} \right)^2$ $A = x^2 + \frac{4x^2 - 24x + 36}{\pi}$ $A = \frac{4x^2 + \pi x^2 - 24x + 36}{\pi}$ $A = \frac{(4 + \pi)x^2 - 24x + 36}{\pi}$	✓ $A = x^2 + \pi r^2$ ✓ $4x + 2\pi r$ ✓ equating to 12 ✓ $r = \frac{6-2x}{\pi}$ ✓ substitution/ vervangings ✓ simplification/ vereenvoudig	(6)
-----	--	---	---	-----



		OR/OF The area of the square = x^2 remaining wire = $12 - 4x$ $2\pi r = 12 - 4x$ $r = \frac{12 - 4x}{2\pi}$ area of the circle = $\pi \left(\frac{12 - 4x}{2\pi} \right)^2$ $= \pi \left(\frac{6 - 2x}{\pi} \right)^2$ $= \frac{36 - 24x + 4x^2}{\pi}$ Total area = area of a square + area of a circle Total area (A) = $x^2 + \frac{36 - 24x + 4x^2}{\pi}$ $= \frac{\pi x^2 + 36 - 24x + 4x^2}{\pi}$ $= \frac{(\pi + 4)x^2 - 24x + 36}{\pi}$	OR/OF ✓ x^2 ✓ $12 - 4x$ ✓ $r = \frac{12 - 4x}{2\pi}$ ✓ $\frac{36 - 24x + 4x^2}{\pi}$ ✓ $x^2 + \frac{36 - 24x + 4x^2}{\pi}$ ✓ $\frac{\pi x^2 + 36 - 24x + 4x^2}{\pi}$	(6)
--	--	--	---	-----



9.2		Area is minimum when: $x = \frac{\frac{24}{\pi}}{2\left(\frac{\pi+4}{\pi}\right)}$ $x = 1,68$ OR/<i>OF</i> For minimum area: $A(x) = \frac{(4+\pi)}{\pi}x^2 - \frac{24}{\pi}x + \frac{36}{\pi}$ $A'(x) = 2\left(\frac{4+\pi}{\pi}\right)x - \frac{24}{\pi} = 0$ $2\left(\frac{4+\pi}{\pi}\right)x = \frac{24}{\pi}$ $x = 1,68 \text{ m}$ For minimum area, the side of the square should be 1,68 m	✓ method/<i>metode</i> ✓ answer/<i>antwoord</i> OR/<i>OF</i> ✓ method/<i>metode</i> ✓ answer/<i>antwoord</i>	(2) (2)
				[8]



QUESTION/VRAAG 10

10.1	10.1.1	$P(G) = \frac{50}{100} = \frac{1}{2} = 0,5$	✓ answer/antwoord	(1)
	10.1.2	$P(S) = \frac{35}{100} = \frac{7}{20} = 0,35$	✓ answer/antwoord	(1)
	10.1.3	$P(G \text{ and } S) = \frac{15}{100} = 0,15$ $P(S) \times P(G) = \frac{1}{2} \times \frac{7}{20} = \frac{7}{40} = 0,175$ $P(G \text{ and } S) \neq P(G) \times P(S)$ Events G and S are not independent	✓ 0, 15 ✓ 0, 175 ✓ Conclusion/ gevolgtrekking	(3)
10.2		$4 \times 7 \times 7 \times 3$ $= 588$	✓ $4 \times 7 \times 7 \times 3$ ✓ answer/antwoord	(2)
10.3	10.3.1	$7! = 5\,040$	✓ 7! or/of 5 040	(1)
	10.3.2	$3 \times 5! \times 4 = 1\,440$	✓ 4 ✓ $3 \times 5!$ ✓ answer/antwoord	(3)
	10.3.3	The number of ways of arranging all the letters with all the three vowels following each other : $3! \times 5! = 720$ $\therefore P(\text{all vowels will not follow each other})$ $= 1 - \frac{720}{5\,040} = \frac{6}{7}$	✓ 3! ✓ $3! \times 5!$ ✓ $1 - \frac{720}{5\,040}$ ✓ $\frac{6}{7}$	(4)
				[15]

TOTAL/TOTAAL : 150