

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



SA EXAM PAPERS

SA EXAM
PAPERS



KWAZULU-NATAL PROVINCE

EDUCATION
REPUBLIC OF SOUTH AFRICA

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

MATHEMATICS P2
MARKING GUIDELINE

SEPTEMBER 2023

MARKS: 150


This marking guideline consists of 14 pages.

NB: CA APPLIES TO ALL SUB-QUESTIONS IN THIS MARKING GUIDELINE.**QUESTION 1**

1.1	$a = 34,90$ $b = 0,08$ $y = 34,90 + 0,08x$ Answer only: Full Marks	✓A Value of a ✓A Value of b ✓CA Equation	(3)
1.2	$r = 0,88$	✓A✓A Answer	(2)
1.3	$y = 34,90 + 0,08x$ $y = 34,90 + 0,08(560)$ $y = 79,7$ minutes = 1.33 hours Total cost = R150 + R150 = R300	✓CA Substitution ✓CA 79,7 minutes ✓CA Answer	(3)
1.4.1	Area = $100 \times 70 = 7000$ square meters $y = 34,90 + 0,08(7000)$ $y = 594,9$ minutes = 9.92 hours	✓CA Calculation	(1)
1.4.2	No. The time taken will exceed his daily 8 hour working hours.	✓CA Justification	(1)
			[10]

QUESTION 2


Daily travelling time x (in minutes)	Number of employees (f)	Midpoint of Interval (x)	$f \cdot x$
$0 \leq x < 10$	20	5	100
$10 \leq x < 20$	35	15	525
$20 \leq x < 30$	30	25	750
$30 \leq x < 40$	10	35	350
$40 \leq x < 50$	5	45	225
Total	100		1950



2.1	Estimated Mean = $\frac{1950}{100} = 19,5$ Answer only: Full Marks	✓A 1950 ✓A 100 ✓CA Answer	(3)
2.2	$10 \leq x < 20$	✓A ✓A Answer	(2)
2.3	See Diagram	✓A Minimum and Maximum value ✓A 1 st and 3 rd Quartiles ✓A 2 nd Quartile	(3)
	Five number summary: 0 ; 12 ; 18 ; 26 ; 50 Accept: ± 1 deviation on quartiles		


2.4.1	True	✓A	Answer	(1)
2.4.2	False	✓A	Answer	(1)
2.4.3	True	✓A	Answer	(1)
				[11]

QUESTION 3

				
3.1	$m_{AD} = \frac{7+3}{1+4} = \frac{10}{5} = 2$	✓A Substitution of points A and D ✓CA Answer	(2)	
3.2	$m_{AD} = m_{BC} = 2 \dots (AD \parallel BC)$ $y = mx + c$ $-8 = 2(-2) + c$ $-4 = c$ $y = 2x - 4$	✓CA Gradient of BC ✓CA Substitution of point C and gradient ✓CA Answer	(3)	


3.3	Let $y = 0$: $0 = 2x - 4$ $x = 2$ $F(2 ; 0)$	✓CA $0 = 2x - 4$ ✓CA x - value	(2)
3.4	$M(3 ; 2)$	✓A x - value ✓A y - value	(2)
3.5	$m_{CD} = \frac{-3 + 8}{-4 + 2} = \frac{5}{-2}$ Inclination of CD: $\tan \theta_1 = 180^\circ - 68,2^\circ = 111,8^\circ$ Inclination of CF: $\tan \theta_2 = 2$ $\theta_2 = 63,43^\circ$ Therefore $\alpha = 111,8^\circ - 63,43^\circ = 48,37^\circ$ OR $DF = \sqrt{(2 + 4)^2 + (0 + 3)^2} = \sqrt{45}$ $DC = \sqrt{29}$ $CF = \sqrt{80}$ $(\sqrt{45})^2 = (\sqrt{29})^2 + (\sqrt{80})^2 - 2\sqrt{29}\sqrt{80}\cos \alpha$ $\alpha = 48,37^\circ$	✓A Gradient of CD ✓A Inclination of CD ✓A Inclination of CF ✓A subtraction of the angles ✓A Distance of DF ✓A Distance of DC ✓A Distance of CF ✓A subst. into cosine rule	(4)
3.6	$DC = \sqrt{(-3 + 8)^2 + (-4 + 2)^2} = \sqrt{29}$ $FC = \sqrt{(0 + 8)^2 + (2 + 2)^2} = \sqrt{80}$ $\text{Area of } \triangle DCF = \frac{1}{2}(\sqrt{29})(\sqrt{80}) \sin 48,37^\circ$ $= 18 \text{ square units.}$	✓A Length of CD ✓CA Length of FC ✓CA Substitution into area formula ✓CA Answer	(4)
			[17]

QUESTION 4

4.1.1	$x^2 + y^2 - 8x + 6y = 15$ $\text{LHS} = (2)^2 + (-9)^2 - 8(2) + 6(-9)$ $= 4 + 81 - 16 - 54$ $= 15$ $= \text{RHS}$	✓A Subst. of point ✓A Simplification	(2)
4.1.2	 $(x - 4)^2 + (y + 3)^2 = 40$ Centre: C(4; -3) P(2; -9) $m_{\text{Radius}} = \frac{-3 + 9}{4 - 2} = \frac{6}{2} = 3$ $m_{\text{Tangent}} = -\frac{1}{3}$ Equation of Tangent: $y = mx + c$ $-9 = -\frac{1}{3}(2) + c$ $-\frac{25}{3} = c$ $y = -\frac{1}{3}x - \frac{25}{3}$	✓A writing the equation as $(x - 4)^2 + (y + 3)^2 = 40$ ✓CA Centre of circle ✓CA Gradient of radius ✓CA Gradient of tangent ✓CA Substitution ✓CA Answer	(6)
4.1.3	$r^2: (x - 4)^2 + (y + 3)^2 = 40$ $r^2 = 40$ $(\text{distance } Q \text{ to the centre})^2 = (-10 - 4)^2 + (12 + 3)^2$ $= 421$ $(\text{Length of tangent})^2 = 421 - 40 = 381$ $\text{Length of tangent} = \sqrt{381}$	✓CA Calculation of r^2 ✓CA distance calculation ✓CA Tangent calculation ✓CA Answer	(4)

4.2.1	$(x - 3)^2 + (y + 2)^2 = 25$ Let $x = 0$: $(0 - 3)^2 + (y + 2)^2 = 25$ $(y + 2)^2 = 16$ $y + 2 = \pm 4$ $y = -6$ or $y = 2$ $B(0; 2)$	✓A Letting $x = 0$ ✓A Simplification ✓CA y - values ✓CA Answer	(4)
4.2.2	$C(6; 2)$	✓CA x - value ✓CA y - value	(2)
4.2.3 (a)	$T(3; -2)$ and $M(12; 10)$ $TM^2 = (12 - 3)^2 + (10 + 2)^2 = 225$ $TM = 15$ units	✓A Coordinates of M ✓CA Answer	(2)
(b)	Radius, center T = 5 units and Radius, center M = 10 units Sum of radii = 15 units Circles touch . $TM =$ Sum of radii	✓CA Sum of radii ✓CA Justification	(2)
			[22]


QUESTION 5

5.1			
5.1.1	$\cos 218^\circ$ $= -\cos 38^\circ$ $= -\frac{\sqrt{1 - p^2}}{1}$	✓A Calculation of $\sqrt{1 - p^2}$ ✓A Reduction ✓CA Answer	(3)

5.1.2	$\cos 14^\circ$ $= \cos(52^\circ - 38^\circ)$ $= \cos 52^\circ \cos 38^\circ + \sin 52^\circ \sin 38^\circ$ $= \left(\frac{p}{1}\right)\left(\frac{\sqrt{1-p^2}}{1}\right) + \left(\frac{\sqrt{1-p^2}}{1}\right)\left(\frac{p}{1}\right)$ $= 2p\sqrt{1-p^2}$ OR $\cos 24^\circ = \sin 76^\circ$ $= \sin 2(38^\circ)$ $= 2 \sin 38^\circ \cos 38^\circ$ $= 2p\sqrt{1-p^2}$	✓ A Writing as difference ✓ A Expansion ✓ CA Answer ✓ $\sin 76^\circ$ ✓ double angle ✓ answer	(3)
5.1.3	$\sin 26^\circ \cos 26^\circ$ $= \frac{1}{2} \sin 52^\circ$ $= \frac{1}{2} \sqrt{1-p^2}$	✓ A Double angle ✓ CA Answer	(2)
5.2	$\frac{2 \sin 165^\circ \cos 195^\circ}{\cos 45^\circ \sin 15^\circ - \cos 15^\circ \sin 45^\circ}$ $= \frac{2 \sin 15^\circ \cdot (-\cos 15^\circ)}{\cos 45^\circ \sin 15^\circ - \cos 15^\circ \sin 45^\circ}$ $= \frac{-2 \sin 30^\circ}{\sin(15^\circ - 45^\circ)}$ $= \frac{-2 \sin 30^\circ}{\sin(-30^\circ)}$ $= \frac{-2 \sin 30^\circ}{-\sin 30^\circ}$ $= 2$	✓ A $-\cos 15^\circ$ ✓ A $-2 \sin 30^\circ$ ✓ A $\sin(15^\circ - 45^\circ)$ ✓ A $-\sin 30^\circ$ ✓ CA Answer	(5)
5.3.1	$K = \sqrt{3} \cos x + \sin x$ $K = 2 \left(\frac{\sqrt{3}}{2} \cos x + \frac{1}{2} \sin x \right)$ $K = 2(\sin 60^\circ \cos x + \cos 60^\circ \sin x)$ $K = 2 \sin(60^\circ + x)$	✓ A $2 \left(\frac{\sqrt{3}}{2} \cos x + \frac{1}{2} \sin x \right)$ ✓ A $\sin 60^\circ$ and $\cos 60^\circ$ ✓ A $2 \sin(60^\circ + x)$	(3)

5.3.2	$t = 2$ and $\theta = 60^\circ$	✓CA t – value and $\theta = 60^\circ$	(1)
5.3.3	2	✓CA Answer	(1)
5.4.1	$\text{LHS} = \frac{2 \tan \theta - \sin 2\theta}{2 \sin^2 \theta}$ $= \frac{2 \left(\frac{\sin \theta}{\cos \theta} \right) - 2 \sin \theta \cos \theta}{2 \sin^2 \theta} \times \frac{\cos \theta}{\cos \theta}$ $= \frac{2 \sin \theta - 2 \sin \theta \cos^2 \theta}{2 \sin^2 \theta \cos \theta}$ $= \frac{2 \sin \theta (1 - \cos^2 \theta)}{2 \sin^2 \theta \cos \theta}$ $= \frac{2 \sin \theta \cdot \sin^2 \theta}{2 \sin^2 \theta \cdot \cos \theta}$ $= \frac{\sin \theta}{\cos \theta}$ $= \tan \theta$ $= \text{LHS}$	✓A $\frac{\sin \theta}{\cos \theta}$ ✓A $2 \sin \theta \cos \theta$ ✓A simplification ✓A factorizing ✓A $1 - \cos^2 \theta = \sin^2 \theta$ ✓A simplified to $\frac{\sin \theta}{\cos \theta}$	(6)
5.4.2	$2 \sin^2 \theta = 0$ $\sin \theta = 0$ $\therefore \theta = 180^\circ \text{ and } 360^\circ$ $\theta = 270^\circ$	✓ 180° and 360° ✓ 270°	(2)
			[26]

QUESTION 6


6.1			
	Graph of f: 1A mark for x – intercepts 1A marks for minimum and maximum points 1A mark for shape Graph of g: 1A mark for end points 1A mark for x – intercepts 1A mark for y – intercept (6)		
6.2.1	360°	✓A Answer	(1)
6.2.2	$x \in [-180^\circ; -150^\circ) \cup (30^\circ; 180^\circ]$	$[-180^\circ; -150^\circ)$ ✓A : Notation ✓CA : values $(30^\circ; 180^\circ]$ ✓A : Notation ✓CA : values	(4)
6.2.3	$f(x) = 1.5 + g(x)$ $f(x) - g(x) = 1.5$ $x = 90^\circ$ or $x = 150^\circ$	✓A $f(x) - g(x) = 1.5$ ✓A Answer	(2)
			[13]

QUESTION 7

7.1	$\frac{PQ}{QR} = \tan \theta$ $PQ = QR \tan \theta$	✓ A Answer	(1)
7.2	$\frac{QR}{\sin \hat{S}} = \frac{QS}{\sin \angle QRS}$ $\frac{QR}{\sin \beta} = \frac{x}{\sin \alpha}$ $QR = \frac{x \sin \beta}{\sin \alpha}$ $PQ = \frac{x \sin \beta \tan \theta}{\sin \alpha}$	✓ A Sine rule formula ✓ A Subs. Sine rule ✓ A Making QR a subj. of the formula ✓ A Subst. of QR	(4)
7.3	$\text{Area of } \triangle QSR = \frac{1}{2}(x)(8 - 2x) \sin 60^\circ$ $= \frac{1}{2}(x)(8 - 2x) \left(\frac{\sqrt{3}}{2}\right)$ $= \frac{\sqrt{3}}{4}(x)(8 - 2x)$ $= \sqrt{3}(x) \left(2 - \frac{1}{2}x\right)$ $= 2\sqrt{3}x - \frac{\sqrt{3}}{2}x^2$	✓ A Subst. into Area rule ✓ A $\frac{\sqrt{3}}{2}$ ✓ A Simplifying	(3)

7.4	For Max Area: $x = -\frac{b}{2a}$ $x = -\frac{(2\sqrt{3})}{2\left(-\frac{\sqrt{3}}{2}\right)}$ $x = 2$	✓ A Formula ✓ A Substitution into formula ✓ CA Answer	(3)
			[11]

QUESTION 8


8.1	 <u>Constr:</u> Join MO and produce to D. $\widehat{AOD} = \widehat{OAM} + \widehat{AMO} \dots$ (Ext. $\angle$ of Δ) $\widehat{BOD} = \widehat{OBM} + \widehat{BMO} \dots$ (Ext. $\angle$ of Δ) But $\widehat{OAM} = \widehat{AMO}$ and $\widehat{OBM} = \widehat{BMO} \dots$ (Radii =) $\therefore \widehat{AOD} + \widehat{BOD} = 2\widehat{AMO} + 2\widehat{BMO}$ $\widehat{AOB} = 2(\widehat{AMO} + \widehat{BMO})$ $\widehat{AOB} = 2\widehat{M}$ NOTE No construction : No marks	✓ A Construction ✓ A S/R ✓ A S ✓ A S ✓ A S	(5)
8.2.1	$\widehat{O\hat{V}A} = \widehat{O\hat{D}A} = 90^\circ \dots$ (Radius $\perp$ Tangent) VODA is a cyclic quad. (Converse of opposite angles of quad. Supplementary)	✓ A S/R ✓ A R	(2)
8.2.2	$\widehat{O_1} = 40^\circ \dots$ (Exterior angle of cyclic quad = int. opp. Angle)	✓ S ✓ R	(2)

[14]


QUESTION 9

9.1	Divides the other two sides, proportionally.	✓S divides the other two sides proportionally	(2)
9.2			
9.2.1	$\frac{KQ}{3} = \frac{6}{4} \dots\dots\dots(\text{Prop. Thm; } KL \parallel QR)$ $KQ = 4,5 \text{ units}$	✓ S/R ✓ Answer	(2)
9.2.2	KL = QM(Opposite sides of $\parallel^m$ QKLM are equal) KL = NR (Opposite sides of $\parallel^m$ NKLR are equal) Therefore QM = NR (both = KL) OR $\frac{QN}{QR} = \frac{QK}{KP}$ $\frac{4,5}{3} = \frac{3}{2}$ $\frac{MR}{QM} = \frac{RL}{PL}$ $= \frac{6}{4} = \frac{3}{2}$ $\therefore \frac{QN}{NR} = \frac{MR}{QM}$ $\frac{QM+1,8}{NR} = \frac{NR+1,8}{QM}$ $\therefore QM(QM+1,8) = MR(NR+1,8)$ $\therefore QM = NR$	✓ S/R ✓ S/R ✓ $\frac{QN}{QR} = \frac{QK}{KP}$ Prop Thm, $KN \parallel PR$ OR $\frac{MR}{QM} = \frac{RL}{PL}$ Prop Thm, $ML \parallel QP$ ✓ $\frac{QM+1,8}{NR} = \frac{NR+1,8}{QM}$	(2)

QUESTION 10

			
10.1	In Δ's MQA and MBQ $\hat{M}_1 = \hat{M}_1$(Common) $\hat{A}_1 = \hat{Q}_1$(Tan-Chord Theorem) $\hat{AQM} = \hat{B}_1$(Remaining $\angle$s of Δ's) $\Delta MQA \parallel \Delta MBQ$(AAA)	✓ S/R ✓ S/R ✓ R	(3)
10.2	In Δ's MAR and MRB $\hat{M}_2 = \hat{M}_2$(Common) $\hat{A}_2 = \hat{R}_1$(Tan-Chord Thm) $\hat{ARM} = \hat{RBM}$(Remaining $\angle$s of Δ's) $\Delta MAR \parallel \Delta MRB$(AAA) $\frac{MA}{MR} = \frac{MR}{MB}$(Δ's similar) $MR^2 = AM \cdot MB$	✓ S ✓ S/R ✓ R ✓ S A✓R	(5)
10.3	$\frac{MQ}{MB} = \frac{MA}{MQ}$(from 10.1) $MQ^2 = MB \cdot MA = \dots \dots (1)$ Also $MR^2 = AM \cdot MB$ Now $\frac{MQ^2}{MR^2} = \frac{MB \cdot MA}{AM \cdot MB}$ $\therefore \frac{MQ^2}{MR^2} = 1$	✓ $\frac{MQ}{MB} = \frac{MA}{MQ}$ ✓ $MQ^2 = MB \cdot MA$ ✓✓ $\frac{MQ^2}{MR^2} = \frac{MB \cdot MA}{AM \cdot MB}$	(4)
			[12]

QUESTION 11

			
11.1	$BC^2 = 8^2 - 4,8^2 \dots\dots\dots(\text{TOP})$ $BC = 6,4 \text{ cm}$	✓ S ✓ S	(2)
11.2	$\triangle BED$	✓ S	(1)
11.3	$\frac{BA}{BE} = \frac{AC}{ED} = \frac{BC}{BD}$ $\frac{8}{BE} = \frac{4,8}{ED} = \frac{6,4}{4}$ $ED = \frac{4,8 \times 4}{6,4} = 3 \text{ cm}$ Area of $\triangle EDB$ $= \frac{1}{2}(4)(3) = 6 \text{ cm}^2$ Area of $\triangle ABC$ $= \frac{1}{2}(4,8)(6,4) = 15,6 \text{ cm}^2$ Therefore, Area of ADEC $= 15,6 - 6 = 9,6 \text{ cm}^2$	$\triangle BAC \parallel \triangle BED$ ✓A S ✓CA Value of ED ✓CA Area of $\triangle EDB$ ✓CA Area of $\triangle ABC$ ✓CA Area of ADEC	(5)
			[8]

GRAND TOTAL: 150