

# SA's Leading Past Year

## Exam Paper Portal




You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ [www.saexampapers.co.za](http://www.saexampapers.co.za)





**LIMPOPO**

**PROVINCIAL GOVERNMENT**  
REPUBLIC OF SOUTH AFRICA

DEPARTMENT OF  
**EDUCATION**

**NATIONAL**  
**SENIOR CERTIFICATE**

**GRADE 12**

**PHYSICAL SCIENCES**  
**PAPER 2 (CHEMISTRY)**  
**MARKING GUIDELINES**  
**SEPTEMBER 2023**

**MARKS: 150**

This marking guideline consists of 9 pages including THE COVER PAGE


**QUESTION 1**

- 1.1 D ✓✓ (2)  
 1.2 C ✓✓ (2)  
 1.3 B ✓✓ (2)  
 1.4 C ✓✓ (2)  
 1.5 A ✓✓ (2)  
 1.6 C ✓✓ (2)  
 1.7 B ✓✓ (2)  
 1.8 D ✓✓ (2)  
 1.9 C ✓✓ (2)  
 1.10 A ✓✓ (2)

**[20]****QUESTION 2**

- 2.1 A series of organic compounds that can be described by the same general formula OR in which one member differs from the next with a CH<sub>2</sub> group. ✓✓ (2)

2.2.1




|                          |   |
|--------------------------|---|
| Functional group         | ✓ |
| Whole structure correct. | ✓ |

(2)

- 2.2.2 1,1-dimethyl✓propan✓-1-ol✓ / 1,1-dimethyl-1-propanol (3)

- 2.2.3 ketone ✓ (1)

2.2.4



(1)

- 2.3.1 C<sub>n</sub>H<sub>2n+2</sub> ✓ (1)

- 2.3.2  $\underline{2\text{C}_4\text{H}_{10} + 13\text{O}_2} \quad \checkmark \rightarrow \underline{8\text{CO}_2 + 10\text{H}_2\text{O}} \quad \checkmark$  bal. ✓ (3)

- 2.4.1 C ✓ (1)

- 2.4.2 ethyl✓ethanoate ✓ (2)

**QUESTION 3**

- 3.1 The pressure exerted by a vapor at equilibrium with its liquid in a closed system. ✓✓ (2)
- 3.2 A ✓ (1)
- 3.3.1 SATURATED ✓ (1)
- 3.3.2 Only single bonds between C- atoms. ✓ (1)
- 3.3.3 London forces ✓ / dispersion forces / induced dipole forces (1)
- 3.4.1 Compounds with the same molecular formula, ✓ but different structural formulae. ✓ (2)
- 3.4.2
- 2-methylbutane is a spherical molecule that offers a smaller surface area to other molecules. ✓ // Pentane is a linear molecule which offers a larger surface area to other molecules.
  - Less/smaller surface area where intermolecular forces (London forces) can interact with other molecules. ✓ // Greater/larger surface area where intermolecular forces (London forces) can interact with other molecules.
  - Less energy required to overcome the intermolecular forces. ✓ // More energy required to overcome the intermolecular forces. (3)
- 3.5
- pentan-1-ol (D): H-bonds are stronger ✓ than the weaker London forces ✓ in (B), alkanes.
  - Therefore, more energy is required to ✓ overcome the stronger intermolecular forces in (D).
  - Consequently (D) has a higher boiling point. ✓ (4)

**OR**


- Pentane (B): London forces are weaker in alkanes than the stronger H-bonds in alcohols in (D).
- Less energy is required to overcome the forces of attraction in (B).

Consequently (B) has a lower boiling point.



**[15]****QUESTION 4**

4.1



4.2


addition / hydrohalogenation ✓

(1)

4.3

butan-2-ol ✓ / 2-butanol

|   |   |
|---|---|
| Hydroxyl group on the 2 <sup>nd</sup> carbon. | ✓ |
| Whole structure correct.                      | ✓ |



(3)

4.4

hydrolysis ✓

(1)

4.5.1

Water ✓

(1)

4.5.2

 $\text{H}_2\text{SO}_4$  ✓ /  $\text{HCl}$  /  $\text{H}_3\text{PO}_4$ 

(1)

4.5.3

hydration ✓ (accept addition)

(1)

4.6.1

but-2-ene / 2-buten ✓

(1)

4.6.2

dehydrohalogenation ✓ / elimination

(1)

**[14]****QUESTION 5**

5.1.1

Sufficient kinetic energy (molecules move fast enough) during the collisions. ✓Molecules must be correctly orientated. ✓

(2)

5.1.2

Increased temperature:

More molecules move fast enough or have sufficient  $E_k$ . ✓  
 There are more effective collisions per unit time ✓ /  $E_k \geq$   
 activation energy.

5.2.1 Activation energy ✓ (2)  
 (1)

5.2.2 (a) Increase in the concentration of one or both reactants. ✓ (1)  
 (b) Increase in temperature. ✓ (1)

5.3.1 How will a change in concentration affect the reaction rate?  
**OF**  
 What is the relationship between the concentration and reaction rate?

- Identify dependent and independent variable. ✓ ✓
- Ask a question (?) about the relationship between dependent and independent variable. ✓

5.3.2  $\text{HNO}_3$  ✓ / Nitric acid (3)  
 The magnesium is used up. / Magnesium is the limited reagent. ✓ (2)


5.3.3

| Opsie 1  | Opsie 2   |
|--|---|
| $\Delta n = 1,0 - 0,8 \checkmark = 0,2 \text{ mol}$<br><br>$n = \frac{m}{M}$<br>$0,2 = \frac{m}{24} \checkmark$<br>$\therefore m = 4,8 \text{ g}$<br><br>$\text{Gem. reaksietempo} = \frac{\Delta m}{\Delta t}$<br>$= \frac{4,8 \checkmark}{30 - 0 \checkmark}$<br>$= 0,16 \text{ g} \cdot \text{s}^{-1} \checkmark$ | $n = \frac{m}{M}$<br>$n = \frac{m}{M}$<br>$1 = \frac{m}{24} \checkmark$<br>$0,8 = \frac{m}{24} \checkmark$<br>$\therefore m = 24 \text{ g}$<br>$\therefore m = 19,2 \text{ g}$<br><br>$\text{Gem. reaksietempo} = \frac{\Delta m}{\Delta t}$<br>$= \frac{19,2 - 24 \checkmark}{30 - 0 \checkmark}$<br>$= 0,16 \text{ g} \cdot \text{s}^{-1} \checkmark$ |

(5)

5.3.4

- Steeper slope below original graph. ✓
- Intercept x-axis earlier. ✓

(2)  
[19]**QUESTION 6**

6.1.1 Increases ✓ (1)

6.1.2 Forward ✓ (1)

6.1.3 When the equilibrium in a closed system is disturbed, the system will re-instate a new equilibrium by favouring the reaction that will oppose the disturbance. ✓✓ (2)

6.1.4 Increase in temperature increases  $K_c$ . ✓  
 Increase in  $K_c$  indicates that the forward reaction has been favoured. ✓  
 Increase in temperature favours the endothermic reaction. ✓  
 Therefore, the forward reaction is endothermic. ✓ (4)

6.1.5 Add a catalyst. ✓ Decrease pressure OR  
 Increase the volume of the container. ✓ (2)

|             | $2\text{SO}_2$ | $\text{O}_2$    | $2\text{SO}_3$ |
|-------------|----------------|-----------------|----------------|
| Initial mol | 8              | y               | 0              |
| Mol reacted | -2x            | -x              | +2x ✓          |
| Mol at eq   | 2              | y-3 ✓           | 6              |
| [ ] at eq   | 1              | $\frac{y-3}{2}$ | 3 ✓ (÷2)       |


$$K_c = \frac{[\text{SO}_3]^2}{[\text{SO}_2]^2 [\text{O}_2]} \quad \checkmark \quad \text{correct } K_c \text{ expression}$$

$$9 = \frac{(3)^2}{(1)^2 (y-3)} \quad \checkmark \quad \text{correct substitution}$$

$$Y = 5\text{mol} \quad \checkmark$$

(6)

[16]

**QUESTION 7**7.1 A proton donor ( $H^+$  ion donor).✓✓ (2)7.2 WEAK ACIDS: ionizes incompletely in water to form only a few  $H_3O^+$  ions.✓  
DILUTED ACIDS: contains a large amount of water added to it.✓7.3.1 EXOTHERMIC ✓ (2)  
(1)7.3.2  $pH = -\log[H_3O^+]$  ✓  
 $2,3 \checkmark = -\log[H_3O^+]$   
 $\therefore [H_3O^+] = 10^{-2,3}$   
 $= 0,005 \text{ mol.dm}^{-3}$  ✓ (3)7.3.3  $n(\text{HCl}) \text{ initial} = cV$  ✓  
 $= 0,25(0,5)$  ✓  
 $= 0,125 \text{ mol}$  ✓  
 $n(\text{HCl}) \text{ after addition} = cV$   
 $= 0,005(1)$  ✓  
 $= 0,005 \text{ mol}$  ✓ (3)Number of moles of HCl reacted with NaOH:  
 $= 0,125 - 0,005 = 0,12 \text{ mol}$  ✓

So 0.12 mol of NaOH reacted with 0.12 mol of HCl:

$$[NaOH]_{\text{initial}} = \frac{n}{V}$$
$$= \frac{0,12}{0,5} \checkmark$$
$$= 0,24 \text{ mol.dm}^{-3} \checkmark$$
 (8)

[16]

**QUESTION 8**

8.1.1 Hydrogen✓ (1)



8.1.2

In terms of the reducing agent:

- Cu is a weaker reducing agent ✓ than H<sub>2</sub> ✓ and will not reduce H<sup>+</sup> (to H<sub>2</sub>). ✓

(3)

In terms of the oxidizing agent:

- H<sup>+</sup> is a weaker oxidising agent than Cu<sup>2+</sup> and will not oxidise Cu (to Cu<sup>2+</sup>).


(NOTE: Compare the two reducing agents in the two half reactions OR the two oxidizing agents in the two half reactions.)

**OR**

H<sup>+</sup> (H<sub>2</sub>SO<sub>4</sub>) is a weaker oxidizing agent than Cu (to Cu<sup>2+</sup>).


(NOTE: No marks if referring to the relative positions on the table.)

8.1.3



(3)

8.2.1



(3)

8.2.2

25 °C ✓ / 298K and 1mol dm<sup>-3</sup> ✓

(2)

8.2.3

$$\begin{aligned} E^\ominus_{\text{cell}} &= E^\ominus_{\text{cathode}} - E^\ominus_{\text{anode}} \\ &= 0,8 - (-2,36) \\ &= 3,16\text{V} \end{aligned}$$

(4)

8.2.4

$$I = \frac{P}{V} = \frac{6\text{W}}{3\text{V}} = 2\text{ A,}$$

The light bulb is manufactured to work effectively when connected to a 3 V source that can deliver a current of 2 A. This cell produces a large enough potential difference, but the current is probably too small ✓ due to a very large internal resistance.

(1)

8.2.5

Anode: Mg is oxidized and therefore forms the anode.

The amount of AgNO<sub>3</sub> available determines how much of the anode (Mg) will go into solution.

1 mol Mg reacts with 2 mol Ag<sup>+</sup>

$$\begin{aligned} n(\text{AgNO}_3) &= cV \\ &= 1(0,4) \\ &= 0,4\text{ mol} \end{aligned}$$

$$\begin{aligned}\therefore n(\text{Mg}) &= \frac{1}{2} n(\text{Ag}^+) \\ &= \frac{1}{2}(0,4) \checkmark \\ &= 0,2 \text{ mol}\end{aligned}$$

$$\begin{aligned}\therefore \text{maximum loss in mass} &= nM \checkmark \\ &= 0,2(24) \checkmark \\ &= 4,8 \text{ g Mg} \checkmark\end{aligned}$$

(6)  
[23]**QUESTION 9**9.1 NEGATIVE  $\checkmark$  (1)9.2 To improve electrical conductivity.  $\checkmark \checkmark$  (2)9.3 Decrease  $\checkmark$   
 $\text{Cu} \rightarrow \text{Cu}^{2+} + 2\text{e}^- \checkmark \checkmark$  (3)9.4  $n(\text{Cu}) = \frac{1}{2}(2) \checkmark$   $\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu}$   
 $= 1 \text{ mol}$ 

$$\begin{aligned}m(\text{Cu}) &= nM \\ &= 1(63,5) \checkmark \\ &= 63,5 \text{ g}\end{aligned}$$

$$\begin{aligned}\% \text{ purity} &= \frac{m_{\text{pure}}}{m_{\text{impure}}} \times 100 \\ &= \frac{63,5 \checkmark}{95,7 \checkmark} \times 100 \\ &= 66,35 \text{ g} \checkmark\end{aligned}$$

(5)  
[11]**TOTAL: 150**



**PAPER 2: THE PAPER SHOULD BE MARKED OUT OF 140.**

**QUESTION 1**

1.6 Removed.

**- 2 marks**

**QUESTION 2**

2.2.2 2-methylbutan-2-ol

**Marking criteria:**

Correct stem : butan-2-ol ✓

Correct substituent : 2-methyl✓

IUPAC name correct including hyphens✓ (3)

**QUESTION 3**

3.3.2 No multiple bonds between C- atoms. ✓ (1)

**QUESTION 4**

4.6.1 **Correct spelling** is but-2-ene

**QUESTION 5**

5.3.1 Accept any of the following factors for the **independent variable**:

- Change the surface area of Mg.
- Add a catalyst.
- Change in the temperature for the reactants.

E.g How will a Change the surface area of Mg affect the reaction rate? ✓✓ (2)

5.3.2 Removed.

- 2 marks

5.3.3 *Average rate*  $= -\frac{\Delta m}{\Delta t}$  ✓

$$= -\frac{0,8 \text{ ✓} - 1,0 \text{ ✓}}{30 - 0 \text{ ✓}}$$

$$= 6,67 \times 10^{-3} \text{ g} \cdot \text{s}^{-1} \text{ ✓} \quad (5)$$

**QUESTION 6**

6.1.4 Endothermic. ✓

- Increase in temperature increases  $K_c$  ✓
- Increase in  $K_c$  indicates that the forward reaction is favoured ✓
- Increase in temperature favours the endothermic reaction ✓ (4)

- 6.1.5
- Decrease pressure ✓
  - Decrease temperature ✓ (2)

6.2 Removed

- 6 marks

**QUESTION 7**

7.2 Weak acids ionize incompletely in water to form low concentration of  $\text{H}_3\text{O}^+$  ✓

Dilute acids contain large amount of water. ✓ (2)

7.3.3 **POSITIVE MARKING FROM 7.3.2****Marking criteria:**

- Formula  $n(\text{HCl})_{\text{initial}} = cV$  ✓
- Substitution of  $c(\text{HCl})_{\text{initial}} = 0,25$  and  $V = 0,5$  ✓
- Substitution of  $c(\text{HCl})_{\text{after addition}} = 0,005$  and  $V = 1$  ✓
- Calculation of  $n(\text{HCl})_{\text{reacted}}$  ✓
- Ratio  $\text{NaOH} : \text{HCl}$  ✓
- Substitution in the formula for  $[\text{NaOH}]$  ✓
- Final answer ✓



$$\begin{aligned}
 n(\text{HCl})_{\text{initial}} &= c V \checkmark \\
 &= 0,25 (0,5) \checkmark \\
 &= 0,125 \text{ mol}
 \end{aligned}$$

$$\begin{aligned}
 n(\text{HCl})_{\text{after addition}} &= c V \\
 &= 0,005 (1) \checkmark \\
 &= 0,005 \text{ mol} \checkmark
 \end{aligned}$$

$$\begin{aligned}
 \text{Number of moles of HCl reacted with NaOH :} \\
 &= 0,125 - 0,005 = 0,12 \text{ mol} \checkmark
 \end{aligned}$$

$$\text{Ratio: } 0,12 \text{ mol HCl} : 0,12 \text{ mol NaOH} \checkmark$$

$$\begin{aligned}
 [\text{NaOH}]_{\text{initial}} &= \frac{n}{V} \\
 &= \frac{0,12}{0,5} \checkmark \\
 &= 0,24 \text{ mol} \cdot \text{dm}^{-3} \checkmark
 \end{aligned}
 \tag{8}$$

### QUESTION 8

8.1.2 H<sub>2</sub> is a stronger reducing agent ✓ than Cu ✓ and Cu will not  
reduce H<sup>+</sup> (to H<sub>2</sub>)

#### ACCEPT

Zn is a stronger reducing agent than Cu, ✓ therefore Zn will be oxidized.

Cu is a weak reducing agent, ✓ than Zn therefore will not undergo oxidation and will not produce gas. ✓

(3)

8.2.2 Temperature of 25°C / 298K ✓

Concentration of 1mol · dm<sup>-3</sup> ✓ (2)

8.2.4 **OR**

Voltage drop due to the internal resistance. ✓ (1)

8.2.5 **MARK ALLOCATION**

**Remove 1 mark from**

$n(\text{AgNO}_3) = cV$

allocate it in the ratio

**QUESTION 9**

$$9.4 \quad n(\text{Cu}) = \frac{1}{2} (2) \checkmark$$

$$= 1 \text{ mol}$$

$$m(\text{Cu}) = n M$$

$$= 1(63,5) \checkmark$$

$$= 63,5 \text{ g}$$

**POSITIVE MARKING**

$$\% \text{ purity} = \frac{m_{\text{pure}}}{m_{\text{impure}}} \times 100$$

$$= \frac{63,5 \checkmark}{95,7 \checkmark} \times 100$$

$$= 66,35 \text{ g} \checkmark$$

(5)

**TOTAL MARKS : 140**