

# SA's Leading Past Year

## Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ [www.saexampapers.co.za](http://www.saexampapers.co.za)



# SA EXAM PAPERS

SA EXAM PAPERS



**LIMPOPO**

PROVINCIAL GOVERNMENT  
REPUBLIC OF SOUTH AFRICA

DEPARTMENT OF  
**EDUCATION**

**NATIONAL  
SENIOR CERTIFICATE**

**GRADE 12**

**PHYSICAL SCIENCES  
PAPER (CHEMISTRY)**

**SEPTEMBER 2023**

**MARKS: 150**

**TIME: 3 hours**



EPHSCP2

**This question paper consists of 20 pages including DATA SHEETS**

Copyright reserved

Please turn over



SA EXAM  
PAPERS

**INSTRUCTIONS AND INFORMATION**

1. The question paper consists of nine questions. Answer **ALL** the questions.
2. Start **EACH** question on a **NEW** page.
3. Number your answers correctly according to the numbering system used in this question paper.
4. Leave **ONE** line between two sub-questions, e.g., between **QUESTION 2.1** and **QUESTION 2.2**.
5. A non-programmable calculator may be used.
6. Appropriate mathematical instruments may be used.
7. Show **ALL** formulae and substitutions in **ALL** calculations.
8. Round off your **FINAL** numerical answers to a minimum of **TWO** decimal places.
9. Give brief motivations, discussions, etc. where required.
10. You are advised to use the attached data sheets.
11. Write neatly and legibly.

**QUESTION 1: MULTIPLE-CHOICE QUESTIONS**

Four options are provided as possible answers to the following questions. Choose the answer and write only the letter (A–D) next to the question number (1.1–1.10), for example 1.10 E

1.1 Which ONE of the following pairs represents compounds that are isomers?

- A.  $\text{CHCl}_3$  and  $\text{CHBr}_3$
- B.  $\text{CH}_3\text{OH}$  and  $\text{CH}_3\text{CH}_2\text{OH}$
- C.  $\text{CH}_3 - \text{CH} = \text{CH}_2$  and  $\text{CH}_2 = \text{CH} - \text{CH}_3$
- D.  $\text{CH}_3\text{COOCH}_3$  and  $\text{CH}_3\text{CH}_2\text{COOH}$  (2)

1.2 Which ONE of the following organic reactions will take place only when exposed to light?

- A.  $\text{CH}_2\text{CH}_2 + \text{H}_2 \rightarrow \text{CH}_3\text{CH}_3$
- B.  $\text{CH}_3\text{CH}_3 \rightarrow \text{CH}_2\text{CH}_2 + \text{H}_2$
- C.  $\text{CH}_3\text{CH}_3 + \text{Cl}_2 \rightarrow \text{CH}_3\text{CH}_2\text{Cl} + \text{HCl}$
- D.  $\text{CH}_2\text{CH}_2 + \text{Cl}_2 \rightarrow \text{CH}_2\text{ClCH}_2\text{Cl}$  (2)

1.3 What is the name of the reaction between an unsaturated hydrocarbon and hydrogen gas?

- A. halogenation
- B. hydrogenation
- C. dehydrohalogenation
- D. hydration (2)

1.4 When an alkane, containing  $n$  carbons, undergoes combustion in excess oxygen, the mole ratio of the combusted products,  $\text{CO}_2 : \text{H}_2\text{O}$ , will be equal to:

- A.  $n : n$
- B.  $n : n + 2$
- C.  $n : n + 1$
- D.  $n : 2n$  (2)



- 1.5 Ammonia gas  $\text{NH}_3(\text{g})$  and oxygen gas,  $\text{O}_2$ , react in a closed container at a constant temperature according to the following equation below.




Which ONE of the combinations of pressure and temperature will result in an increase in the yield of nitrogen monoxide  $\text{NO}(\text{g})$ ?

	Pressure	Temperature
A.	decrease	decrease
B.	decrease	increase
C.	increase	increase
D.	increase	decrease

(2)

- 1.6 Dinitrogen tetra oxide,  $\text{N}_2\text{O}_4(\text{g})$  decomposes to produce nitrogen dioxide,  $\text{NO}_2(\text{g})$ , in a closed container according to the following balanced equation:



Consider the following statements regarding this equation:


- (i): The forward reaction is endothermic.
- (ii): The temperature of the container decreases.
- (iii): For each mole of  $\text{NO}_2(\text{g})$  which is formed,  $x\text{kJ}$  of energy is released

Which ONE of the following statements are/is true about this reaction?

- A. (i) only
- B. (i) and (ii) only
- C. (ii) and (iii) only
- D. (i), (ii) and (iii)

(2)


- 1.7 Four gases, **P**, **Q**, **R**, and **S** are in equilibrium in a closed container. The diagram below shows their relative concentrations over a period of time. The equilibrium position is altered at  $t_1$  by the addition of more of gas **Q**.



The equilibrium reaction is shown by:

- A.  $P + Q \rightleftharpoons R + S$
- B.  $Q + S \rightleftharpoons P + R$
- C.  $P + S \rightleftharpoons Q + R$
- D.  $Q \rightleftharpoons P + R + S$  (2)
- 1.8 The ionisation constant,  $K_w$ , for pure water at  $25^\circ\text{C}$  is:
- A. 14
- B. 7
- C.  $1 \times 10^{-7}$
- D.  $1 \times 10^{-14}$  (2)
- 1.9 The electrode potential of the  $\text{Ag}^+/\text{Ag}$ -electrode is determined with the aid of a standard  $\text{H}^+/\text{H}_2$ , Pt-electrode as reference electrode. The equation for the half-reaction which occurs at the ANODE of this cell, is:
- A.  $\text{Ag} \rightarrow \text{Ag}^+ + \text{e}^-$
- B.  $\text{Ag}^+ + \text{e}^- \rightarrow \text{Ag}$
- C.  $\text{H}_2 \rightarrow 2\text{H}^+ + 2\text{e}^-$
- D.  $2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$  (2)

- 1.10 An electrochemical cell, consisting of two standard half cells, is allowed to deliver current for a period. Which one of the following combinations best represents the change in CONCENTRATION of the two electrolytes over the period.



(2)

[20]



**QUESTION 2 (Start on a new page.)**

The table below shows six organic molecules A to F

<b>A.</b> butane	<b>B.</b> $\text{CH}_3\text{-CH}_2\text{-OH}$
<b>C.</b> $\text{CH}_3\text{COOH}$	<b>D.</b> <pre>       H             H-C-H           H       H   H                 H-C - C - C - C-H                   H   O   H   H               H           </pre>
<b>E.</b> methanal	<b>F.</b> <pre>       H   H   O   H                         H-C - C - C - C-H                           H   H   H   H           </pre>

- 2.1 Define the term *homologous series*. (2)
- 2.2 Write down the:
- 2.2.1 Structural formula of compound C (2)
- 2.2.2 IUPAC name of compound D (3)
- 2.2.3 Homologous series to which F belongs (1)
- 2.2.4 Structural formula of the functional group of E (1)
- 2.3 For compound A, write down:
- 2.3.1 The general formula. (1)
- 2.3.2 The balanced equation for the oxidation reaction. (3)



NSC

- 2.4 Compound **B** reacts with another organic compound from the table to form an ester.

2.4.1 Write down the LETTER of the other organic compound that reacts with **B**. (1)

2.4.2 Write down the NAME of the ester that forms. (2)

[16]

**QUESTION 3 (Start on a new page.)**

The table below shows data collected for four organic compounds, represented by the letters **A**, **B**, **C** and **D**, during a practical investigation:

	Organic compound	Boiling point (°C)
<b>A</b>	propane	-42
<b>B</b>	pentane	36
<b>C</b>	2-methylbutane	27,8
<b>D</b>	pentan-1-ol	137

- 3.1 Define the term *vapour pressure*. (2)

- 3.2 Which ONE of the compounds **A** or **B** has the higher vapour pressure? (1)

- 3.3 Consider compound **C**.

3.3.1 Is compound **C** a SATURATED or UNSATURATED hydrocarbon? (1)

3.3.2 Give a reason for your answer in QUESTION 3.3.1. (1)

3.3.3 Which type of intermolecular forces are mainly present between these molecules? (1)

- 3.4 Compounds **B** and **C** are structural isomers of each other.

3.4.1 Define the term *structural isomer*. (2)


3.4.2 Refer to structure, intermolecular forces, and energy in your explanation to explain why **B** has a higher boiling point than **C** (3)

- 3.5 Explain the difference in the boiling points of **B** and **D**. Refer to intermolecular forces and energy in your explanation. (4)

[15]

**QUESTION 4 (Start on a new page.)**

The flow chart below shows the preparation of various organic compounds using but-1-ene as starting material.



- 4.1 Using STRUCTURAL formulas, write down a balanced chemical equation for Reaction 1 (4)
- Write down the:
- 4.2. Name of the type of the reaction that occurs when but-1-ene is converted to compound X (1)
- 4.3 Write the structural formula and IUPAC name of the secondary alcohol, Y, that forms. (3)
- 4.4 Name the type of substitution reaction that occurs when compound X is converted to the secondary alcohol Y (1)
- 4.5 But-1-ene can be converted directly to the secondary alcohol, without the formation of the intermediate product X, with the help of a catalyst.
- 4.5.1 Except but-1-ene, write the NAME of the reactant needed for this reaction. (1)
- 4.5.2 Write down the FORMULA of the catalyst that can be used. (1)
- 4.5.3 Name the type of reaction that will occur during this direct

conversion.

(1)

4.6 Instead of using water as a reactant in reaction 2, concentrated sodium hydroxide is used, and the mixture is heated.

4.6.1 Write down the IUPAC name of the compound **W** that forms. (1)

4.6.2 Name the TYPE of reaction that occurs. (1)


**[14]****QUESTION 5 (Start on a new page.)**

5.1 The collision theory can be used to explain how different factors affect the rate of a chemical reaction.

5.1.1 Name TWO conditions that determine whether a collision between two molecules will lead to a chemical reaction. (2)

5.1.2 In terms of the collision theory, explain why the rate of a chemical reaction increases with increasing temperature. (2)

5.2 Curve **R** represents the Maxwell-Boltzmann distribution curve for gas in a closed container at a certain temperature. Curve **Q** and **S** represents the curves of the same molecules at different conditions.




5.2.1 Write down the NAME of the energy represented by **P**. (1)

5.2.2 Write down the change in the condition(s) which results in:

(a) Curve **Q** (1)


(b) Curve **S** (1)

- 5.3 A teacher uses the reaction between magnesium ribbon,  $\text{Mg(s)}$  and nitric acid,  $\text{HNO}_3(\text{aq})$ , to investigate ONE of the factors that influence reaction rate. The balanced equation of the reaction between  $\text{Mg(s)}$  and  $\text{HNO}_3(\text{aq})$  is given below.



- 5.3.1 Write down an investigative question for this investigation. (3)

The results obtained when using DILUTED nitric acid are shown on the graph below.




- 5.3.2 Which compound,  $\text{Mg}$  or  $\text{HNO}_3$ , is in excess, and give a reason for your answer using the information on the graph. (2)
- 5.3.3 Calculate the average reaction rate (in grams per second) during the first 30 s. (5)
- 5.3.4 Copy the above graph in your answer book. On the same set of axes, use a **DOTTED LINE** to show the curve that will be obtained when CONCENTRATED nitric acid was used. No numerical values are required. (2)

[19]




**QUESTION 6 (Start on a new page.)**

- 6.1 The graph below shows the effect of the temperature on the value of  $K_c$  for the following reaction taking place in a closed container.



- 6.1.1 Write down the effect of increasing temperature has on the amount of  $\text{NO}_2(\text{g})$  formed. (1)
- 6.1.2 Which reaction was favoured due to an increase in temperature? Write only FORWARD or REVERSE. (1)
- 6.1.3 State Le Chatelier's Principle. (2)
- 6.1.4 Is the forward reaction EXOTHERMIC or ENDOTHERMIC? Explain your answer. (4)
- 6.1.5 Write down TWO changes, that can be introduced to decrease the rate of the forward reaction (2)
- 6.2 1 mol  $\text{SO}_2(\text{g})$  and x mole  $\text{O}_2(\text{g})$  is initially placed in an empty  $2\text{dm}^3$  container and sealed at a specific temperature. At equilibrium 6 mole  $\text{SO}_3(\text{g})$  was present in the container. If the value for the following equilibrium at this temperature is 9, calculate x, the initial quantity  $\text{O}_2(\text{g})$  that was placed in the container. (6)




**QUESTION 7 (Start on a new page.)**

7.1 Define an acid according to Bronsted and Lowry (2)

7.2 Explain the difference between a WEAK ACID and a DILUTE ACID. (2)

7.3 When 500 cm<sup>3</sup> of sodium hydroxide solution, NaOH(aq) is added to 500 cm<sup>3</sup> of 0,25 mol dm<sup>-3</sup> of hydrochloric acid solution HCl(aq), the temperature rises and the pH of the solution changes to 2,3. The reaction between NaOH(aq) and HCl (aq) is as follows:



7.3.1 Is the reaction between NaOH (aq) and HCl (aq) EXOTHERMIC or ENDOTHERMIC? (1)

Calculate the:


7.3.2 Final concentration of the hydronium ions, H<sub>3</sub>O<sup>+</sup>(aq) ions. (3)

7.3.3 Initial concentration of the NaOH (aq) solution in 500cm<sup>3</sup> (8)

[16]

**QUESTION 8 (Start on a new page.)**

An investigation to compare the relative reducing ability of copper, Cu(s) and zinc, Zn (s) is carried out. The two metals are reacted with dilute sulphuric acid and the experiment set up as shown below.



- 8.1 In test tube A, bubbles of gas are observed whilst in test tube B no bubbles are observed.
- 8.1.1 Write down the name of the gas produced in test tube A (1)
- 8.1.2 Refer to the relative reducing abilities of Cu and Zn to explain why no bubbles of gas are produced in test tube B (3)
- 8.1.3 Use the STANDARD REDUCTION POTENTIAL TABLE to write down a balanced equation of the reaction which takes place in test tube A (3)
- 8.2 An electrochemical cell is set up using a redox reaction given below.
- $$Mg(s) + 2AgNO_3(aq) \rightarrow Mg(NO_3)_2(aq) + 2Ag(s)$$
- 8.2.1 Write the cell notation of this cell. (3)



LimpopoDoE/September 2023

8.2.2 Write down the conditions needed for the cell to operate under standard conditions. (2)

8.2.3 Calculate the initial emf of the cell under standard conditions (4)

This cell is now connected to a light bulb which is marked 3 V; 6 W. In theory, the bulb should glow, but in practice it doesn't.


8.2.4 Without any calculation, give an explanation for this. (1)

The volume of the electrolyte in each standard half-cell is 0,4 dm<sup>3</sup>.

8.2.5 Calculate the maximum loss of mass that can occur at the anode. (6)  
[23]

**QUESTION 9 (Start on a new page.)**

The electrolytic cell shown below is used to purify IMPURE copper. Copper sulphate solution,  $\text{CuSO}_4(\text{aq})$ , is used as the electrolyte. Electrode B contains 95,7% Cu by mass.



- 9.1 Is electrode A, the POSITIVE or NEGATIVE electrode? (1)
- 9.2 Pure copper is used to make electrical wires. Give one reason why the copper must be purified first before it is used in electrical wires. (2)
- 9.3 Will the mass of electrode B INCREASE or DECREASE when the cell is operating? Write down the relevant half-reaction to support your answer. (3)
- 9.4 If 2 moles of electrons are transferred from the anode to the cathode during electrolysis, calculate the mass of the IMPURE copper sample. (5)

[11]

**TOTAL: 150**

(I)	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
(II)												(III)	(IV)	(V)	(VI)	(VII)	(VIII)
1 H 1																	
3 Li 7	4 Be 9																
11 Na 23	12 Mg 24																
19 K 39	20 Ca 40	21 Sc 45	22 Ti 48	23 V 51	24 Cr 52	25 Mn 55	26 Fe 56	27 Co 59	28 Ni 59	29 Cu 63.5	30 Zn 65	31 Ga 70	32 Ge 73	33 As 75	34 Se 79	35 Br 80	36 Kr 84
37 Rb 86	38 Sr 88	39 Y 89	40 Zr 91	41 Nb 92	42 Mo 96	43 Tc 98	44 Ru 101	45 Rh 103	46 Pd 106	47 Ag 108	48 Cd 112	49 In 115	50 Sn 119	51 Sb 122	52 Te 128	53 I 127	54 Xe 131
55 Cs 133	56 Ba 137	57 La 139	58 Ce 140	59 Pr 141	60 Nd 144	61 Pm	62 Sm 150	63 Eu 152	64 Gd 157	65 Tb 159	66 Dy 163	67 Ho 165	68 Er 167	69 Tm 169	70 Yb 173	71 Lu 175	72 Hf 179
87 Fr	88 Ra 226	89 Ac															

KEY/SLEUTEL

Atomic number  
Atoomgetal

Electronegativity  
Elektronegativiteit

Symbol  
Simbool

Approximate relative atomic mass  
Benaderde relatieve atoommassa

29 Cu 63.5



**TABLE 4A: STANDARD REDUCTION POTENTIALS**  
**TABEL 4A: STANDAARD-REDUKSIEPOTENSIALE**

Half-reactions/Halfreaksies	$E^{\ominus}$ (V)
$F_2(g) + 2e^- \rightleftharpoons 2F^-$	+ 2,87
$Co^{3+} + e^- \rightleftharpoons Co^{2+}$	+ 1,81
$H_2O_2 + 2H^+ + 2e^- \rightleftharpoons 2H_2O$	+ 1,77
$MnO_4^- + 8H^+ + 5e^- \rightleftharpoons Mn^{2+} + 4H_2O$	+ 1,51
$Cl_2(g) + 2e^- \rightleftharpoons 2Cl^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^- \rightleftharpoons 2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^- \rightleftharpoons 2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^- \rightleftharpoons Mn^{2+} + 2H_2O$	+ 1,23
$Pl^{2+} + 2e^- \rightleftharpoons Pl$	+ 1,20
$Br_2(l) + 2e^- \rightleftharpoons 2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^- \rightleftharpoons NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^- \rightleftharpoons Hg(l)$	+ 0,85
$Ag^+ + e^- \rightleftharpoons Ag$	+ 0,80
$NO_3^- + 2H^+ + e^- \rightleftharpoons NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^- \rightleftharpoons Fe^{2+}$	+ 0,77
$O_2(g) + 2H^+ + 2e^- \rightleftharpoons H_2O_2$	+ 0,68
$I_2 + 2e^- \rightleftharpoons 2I^-$	+ 0,54
$Cu^+ + e^- \rightleftharpoons Cu$	+ 0,52
$SO_2 + 4H^+ + 4e^- \rightleftharpoons S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^- \rightleftharpoons 4OH^-$	+ 0,40
$Cu^{2+} + 2e^- \rightleftharpoons Cu$	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^- \rightleftharpoons SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^- \rightleftharpoons Cu^+$	+ 0,16
$Sn^{4+} + 2e^- \rightleftharpoons Sn^{2+}$	+ 0,15
$S + 2H^+ + 2e^- \rightleftharpoons H_2S(g)$	+ 0,14
$2H^+ + 2e^- \rightleftharpoons H_2(g)$	0,00
$Fe^{3+} + 3e^- \rightleftharpoons Fe$	- 0,06
$Pb^{2+} + 2e^- \rightleftharpoons Pb$	- 0,13
$Sn^{2+} + 2e^- \rightleftharpoons Sn$	- 0,14
$Ni^{2+} + 2e^- \rightleftharpoons Ni$	- 0,27
$Co^{2+} + 2e^- \rightleftharpoons Co$	- 0,28
$Cd^{2+} + 2e^- \rightleftharpoons Cd$	- 0,40
$Cr^{3+} + e^- \rightleftharpoons Cr^{2+}$	- 0,41
$Fe^{2+} + 2e^- \rightleftharpoons Fe$	- 0,44
$Cr^{3+} + 3e^- \rightleftharpoons Cr$	- 0,74
$Zn^{2+} + 2e^- \rightleftharpoons Zn$	- 0,76
$2H_2O + 2e^- \rightleftharpoons H_2(g) + 2OH^-$	- 0,83
$Cr^{2+} + 2e^- \rightleftharpoons Cr$	- 0,91
$Mn^{2+} + 2e^- \rightleftharpoons Mn$	- 1,18
$Al^{3+} + 3e^- \rightleftharpoons Al$	- 1,66
$Mg^{2+} + 2e^- \rightleftharpoons Mg$	- 2,36
$Na^+ + e^- \rightleftharpoons Na$	- 2,71
$Ca^{2+} + 2e^- \rightleftharpoons Ca$	- 2,87
$Sr^{2+} + 2e^- \rightleftharpoons Sr$	- 2,89
$Ba^{2+} + 2e^- \rightleftharpoons Ba$	- 2,90
$Cs^+ + e^- \rightleftharpoons Cs$	- 2,92
$K^+ + e^- \rightleftharpoons K$	- 2,93
$Li^+ + e^- \rightleftharpoons Li$	- 3,05

Increasing oxidising ability/Toenemende oksiderende vermoë

Increasing reducing ability/Toenemende reducerende vermoë

TABLE 4B: STANDARD REDUCTION POTENTIALS  
TABEL 4B: STANDAARD-REDUKSIEPOTENSIALE

Half-reactions/Halfreaksies	$E^0$ (V)
$\text{Li}^+ + \text{e}^- \rightleftharpoons \text{Li}$	-3,05
$\text{K}^+ + \text{e}^- \rightleftharpoons \text{K}$	-2,93
$\text{Cs}^+ + \text{e}^- \rightleftharpoons \text{Cs}$	-2,92
$\text{Ba}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ba}$	-2,90
$\text{Sr}^{2+} + 2\text{e}^- \rightleftharpoons \text{Sr}$	-2,89
$\text{Ca}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ca}$	-2,87
$\text{Na}^+ + \text{e}^- \rightleftharpoons \text{Na}$	-2,71
$\text{Mg}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mg}$	-2,36
$\text{Al}^{3+} + 3\text{e}^- \rightleftharpoons \text{Al}$	-1,66
$\text{Mn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mn}$	-1,18
$\text{Cr}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cr}$	-0,91
$2\text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-$	-0,83
$\text{Zn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Zn}$	-0,76
$\text{Cr}^{3+} + 3\text{e}^- \rightleftharpoons \text{Cr}$	-0,74
$\text{Fe}^{2+} + 2\text{e}^- \rightleftharpoons \text{Fe}$	-0,44
$\text{Cr}^{3+} + \text{e}^- \rightleftharpoons \text{Cr}^{2+}$	-0,41
$\text{Cd}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cd}$	-0,40
$\text{Co}^{2+} + 2\text{e}^- \rightleftharpoons \text{Co}$	-0,28
$\text{Ni}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ni}$	-0,27
$\text{Sn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Sn}$	-0,14
$\text{Pb}^{2+} + 2\text{e}^- \rightleftharpoons \text{Pb}$	-0,13
$\text{Fe}^{3+} + 3\text{e}^- \rightleftharpoons \text{Fe}$	-0,06
$2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g})$	0,00
$\text{S} + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2\text{S}(\text{g})$	+0,14
$\text{Sn}^{4+} + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}$	+0,15
$\text{Cu}^{2+} + \text{e}^- \rightleftharpoons \text{Cu}^+$	+0,16
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{SO}_2(\text{g}) + 2\text{H}_2\text{O}$	+0,17
$\text{Cu}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cu}$	+0,34
$2\text{H}_2\text{O} + \text{O}_2 + 4\text{e}^- \rightleftharpoons 4\text{OH}^-$	+0,40
$\text{SO}_2 + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons \text{S} + 2\text{H}_2\text{O}$	+0,45
$\text{Cu}^+ + \text{e}^- \rightleftharpoons \text{Cu}$	+0,52
$\text{I}_2 + 2\text{e}^- \rightleftharpoons 2\text{I}^-$	+0,54
$\text{O}_2(\text{g}) + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2\text{O}_2$	+0,68
$\text{Fe}^{3+} + \text{e}^- \rightleftharpoons \text{Fe}^{2+}$	+0,77
$\text{NO}_3^- + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{NO}_2(\text{g}) + \text{H}_2\text{O}$	+0,80
$\text{Ag}^+ + \text{e}^- \rightleftharpoons \text{Ag}$	+0,80
$\text{Hg}_2^{2+} + 2\text{e}^- \rightleftharpoons \text{Hg}(\text{l})$	+0,85
$\text{NO}_3^- + 4\text{H}^+ + 3\text{e}^- \rightleftharpoons \text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+0,96
$\text{Br}_2(\text{l}) + 2\text{e}^- \rightleftharpoons 2\text{Br}^-$	+1,07
$\text{Pt}^{2+} + 2\text{e}^- \rightleftharpoons \text{Pt}$	+1,20
$\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 2\text{H}_2\text{O}$	+1,23
$\text{O}_2(\text{g}) + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$	+1,23
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightleftharpoons 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+1,33
$\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-$	+1,38
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 4\text{H}_2\text{O}$	+1,51
$\text{H}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$	+1,77
$\text{Co}^{3+} + \text{e}^- \rightleftharpoons \text{Co}^{2+}$	+1,81
$\text{F}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{F}^-$	+2,87

Increasing oxidising ability/Toenemende oksiderende vermoë

Increasing reducing ability/Toenemende reduserende vermoë

**GEGEWENS VIR FISIESTE WETENSKAPPE GRAAD 12**  
**VRAESTEL 2 (CHEMIE)**  
**DATA FOR PHYSICAL SCIENCES GRADE 12**  
**PAPER 2 (CHEMISTRY)**

**TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESTE KONSTANTES**

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Avogadro's constant <i>Avogadro-konstante</i>	$N_A$	$6,02 \times 10^{23} \text{ mol}^{-1}$
Charge on electron <i>Lading op elektron</i>	$e$	$-1,6 \times 10^{-19} \text{ C}$
Electron mass <i>Elektronmassa</i>	$m_e$	$9,11 \times 10^{-31} \text{ kg}$
Molar gas volume at STP <i>Molêre gasvolume by STD</i>	$V_m$	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard pressure <i>Standaarddruk</i>	$p^\theta$	$1,013 \times 10^5 \text{ Pa}$
Standard temperature <i>Standaardtemperatuur</i>	$T^\theta$	$273 \text{ K}$

**TABLE 2: FORMULAE/TABEL 2: FORMULES**

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ or/of $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$
$K_w = [H_3O^+][OH^-] = 1 \times 10^{-14} \text{ at / by } 298 \text{ K}$	
$\text{pH} = -\log[H_3O^+]$	
$E_{\text{cell}}^\theta = E_{\text{cathode}}^\theta - E_{\text{anode}}^\theta / E_{\text{sel}}^\theta = E_{\text{katode}}^\theta - E_{\text{anode}}^\theta$ or / of $E_{\text{cell}}^\theta = E_{\text{reduction}}^\theta - E_{\text{oxidation}}^\theta / E_{\text{sel}}^\theta = E_{\text{reduksie}}^\theta - E_{\text{oksidasie}}^\theta$ or / of $E_{\text{cell}}^\theta = E_{\text{oxidising agent}}^\theta - E_{\text{reducing agent}}^\theta / E_{\text{sel}}^\theta = E_{\text{oksideermiddel}}^\theta - E_{\text{reduksieermiddel}}^\theta$	