

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



**SA EXAM
PAPERS**
SA EXAM
PAPERS



education

Department of
Education
FREE STATE PROVINCE

**PREPARATORY EXAMINATION
VOORBEREIDENDE EKSAMEN**

GRADE/GRAAD 12

**PHYSICAL SCIENCES: CHEMISTRY (P2)
FISIESE WETENSKAPPE: CHEMIE (V2)**

SEPTEMBER 2024

MARKS/PUNTE: 150

MARKING GUIDELINES/NASIENRIGLYNE

These marking guidelines consist of 14 pages.
Hierdie nasienriglyne bestaan uit 14 bladsye.

QUESTION 1/VRAAG 1

- 1.1 B ✓✓ (2)
- 1.2 B ✓✓ (2)
- 1.3 C ✓✓ (2)
- 1.4 D ✓✓ (2)
- 1.5 B ✓✓ (2)
- 1.6 C ✓✓ (2)
- 1.7 D ✓✓ (2)
- 1.8 C ✓✓ (2)
- 1.9 A ✓✓ (2)
- 1.10 A ✓✓ (2)
- [20]**

QUESTION 2/VRAAG 2

- 2.1.1 A ✓ (1)
- 2.1.2 E ✓ (1)
- 2.1.3 D ✓ (1)
- 2.2.1 Tertiary (alcohol) / *Tersiêre (alkohol)* ✓
The carbon atom bonded to the hydroxyl group is bonded to three other carbon atoms. ✓
Die koolstofatoom wat aan die hidroksielgroep gebind is, is aan drie ander koolstofatome gebind. (2)
- 2.2.2 3-methylheptan-3-ol / *metielheptaan-3-ol*

Marking criteria/Nasienkriteria:

- Correct stem of alcohol, i.e. heptanol. ✓
- Substituent correctly identified, i.e. methyl. ✓
- IUPAC name completely correct including numbering and hyphens. ✓
- *Korrekte stam van alkohol, d.i. heptanol.*
- *Substituent korrek geïdentifiseer, d.i. metiel.*
- *IUPAC-naam heeltemal korrek, insluitend nommering en koppeltekens.*

(3)

- 2.2.3 Hydroxyl (group) / *Hydroksiel (groep)* ✓ (1)



Marking Guidelines/Nasiennriglyne

2.3.1


**Marking criteria/Nasienkriteria:**

- Correct functional group of carboxylic acid. ✓
- Whole structure of carboxylic acid correct. ✓
- Correct functional group of alcohol. ✓
- Whole structure of alcohol correct. ✓

- *Korrekte funksionele groep karboksielsuur.*
- *Hele struktuur van karboksielsuur korrek.*
- *Korrekte funksionele groep alkohol.*
- *Hele struktuur van alkohol korrek.*


(4)

2.3.2



(1)

2.4

**Marking criteria/Nasienkriteria:**

- Methyl substituent on second C atom. ✓
- Two fluorine atoms on third C atom. ✓
- Whole structure correct. ✓

- *Metielsubstituent op tweede C-atoom.*
- *Twee fluooratome op derde C-atoom.*
- *Hele struktuur korrek.*

(3)

$$\begin{aligned} 2.5 \quad n(\text{CO}_2) &= \frac{m}{M} \checkmark (a) \\ &= \frac{2750}{44} \checkmark (b) \\ &= 62,5 \text{ mol} \end{aligned}$$

$$\begin{aligned} n(\text{CH}_4) &= n(\text{CO}_2) \checkmark (c) \\ &= 62,5 \text{ mol} \end{aligned}$$

Marking criteria/Nasienkriteria:

- (a) Formula/Formule $n = \frac{m}{M}$ ✓
- (b) Substitute/Vervanging $M = 44 \text{ g} \cdot \text{mol}^{-1}$ in $n = \frac{m}{M}$ ✓
- (c) Use mole ratio/Gebruik molverhouding:
 $n(\text{CH}_4) = n(\text{CO}_2)$ ✓
- (d) Substitute/Vervanging $M = 16 \text{ g} \cdot \text{mol}^{-1}$ in $n = \frac{m}{M}$ ✓
- (e) Correct final answer/Korrekte finale antwoord:
1 000 g OR 1 kg ✓

$$\begin{aligned} m(\text{CH}_4) &= n(\text{CH}_4) \times M(\text{CH}_4) \\ &= 62,5 \times 16 \checkmark (d) \\ &= 1000 \text{ g} / 1 \text{ kg} \checkmark (e) \end{aligned}$$

(5)

[22]

QUESTION 3/VRAAG 3**3.1 Marking criteria/Nasienkriteria:**

If any of the underlined key phrases in the correct context is omitted, deduct 1 mark.

Indien enige van die onderstreepte sleutel frases in die korrekte konteks weggelaat word, trek 1 punt af.

The temperature at which the vapour pressure of a liquid is equal to the atmospheric pressure. ✓✓

Die temperatuur waarby die dampdruk van 'n vloeistof gelyk is aan die atmosferiese druk. (2)

3.2.1 FROM A TO C

- Increase branching/smaller surface area/more compact ✓
- Weaker intermolecular forces ✓
- Less energy needed to break the intermolecular forces ✓
- Boiling point decreases ✓

VANAF A TOT C

- *Vergroot vertakking/kleiner oppervlakte/meer kompak*
- *Swakker intermolekulêre kragte*
- *Minder energie benodig om die intermolekulêre kragte te breek*
- *Kookpunt neem af* (4)

3.2.2 C/2,2-dimethylpropane / dimetielpropan ✓ (1)

3.2.3 Lowest boiling point / Laagste kookpunt ✓ (1)

3.3 Greater than (103 °C) ✓

- Between compound D/pentanal molecules are dipole-dipole forces ✓ (and London forces) and between compound E/butanoic acid are hydrogen bonds ✓ (dipole-dipole and London forces).
- Dipole-dipole forces are weaker than hydrogen bonds. ✓

OR

Intermolecular forces between compound D/pentanal molecules are weaker than those between compound E/butanoic acid molecules.

OR

Less energy is needed to break the intermolecular forces between pentanal molecules.

Groter as (103 °C)

- *Tussen verbinding D/pentanale molekules is dipool-dipoolkragte (en Londen-kragte) en tussen verbinding E/butaansuur is waterstofbindings (dipool-dipool en Londen-kragte).*
- *Dipool-dipoolkragte is swakker as waterstofbindings.*

OF



Intermolekulêre kragte tussen verbinding D/pentanale molekules is swakker as dié tussen verbinding E/butaansuurmolekules.

OF

Minder energie benodig om die intermolekulêre kragte tussen pentanale molekules te breek.

(4)

3.4 Yes/Ja ✓

(1)

3.5 Comparable molecular masses AND only functional group (homologous series) changed/only 1 independent variable ✓

Vergelykbare molekulêre massas EN enigste funksionele groep (homoloë reeks) verander/slegs 1 onafhanklike veranderlike.

(1)

[14]

QUESTION 4/VRAAG 4

4.1 A reaction of a halogen with a compound. ✓✓
'n Reaksie van 'n halogeen met 'n verbinding.

(2)

4.2.1 Heat/Sunlight/UV light / *Hitte/Sonlig/UV-lig* ✓

(1)

4.2.2 HBr ✓

(1)

4.3 $\text{CH}_3\text{CH}_2\text{Br}$ ✓ + NaOH/ ✓ \longrightarrow CH_2CH_2 ✓ + NaBr/ ✓ + H_2O ✓
KOH/LiOH KBr/LiBr

(5)

4.4 Addition/Hydration / *Addisie/Hidrasie* ✓


(1)

4.5 Methanoic acid / *Metanoësuur* ✓✓

(2)

4.6 Esterification/Condensation / *Verestering/Kondensasie* ✓

(1)



[17]



QUESTION 5/VRAAG 55.1 Mass meter/Scale / *Massameter/Skaal* ✓ (1)5.2 Calcium carbonate / *Kalsiumkarbonaat* ✓ (1)

- 5.3 • Change in concentration/mass/amount/volume ✓ of reactants/products per unit time. ✓
 • Concentration/amount/mass/volume of reactants used/products formed per unit time.
 • Rate of change in concentration/amount/mass/volume. **(2 or 0)**
 • *Verandering in konsentrasie/massa/hoeveelheid/volume van reaktante/produkte per tydseenheid.*
 • *Konsentrasie/hoeveelheid/massa/volume van reaktante gebruik/produkte gevorm per tydseenheid.*
 • *Tempo van verandering in konsentrasie/hoeveelheid/massa/volume.* (2)

5.4 **Marking criteria/Nasiennriglyne:**

- Mass of CO₂ formed/*Massa van CO₂ gevorm*: 270-268,4 or/of 1,6g ✓
- Substituting/*Vervanging* 44 in $n = \frac{m}{M}$ ✓
- Substitute number of moles of CO₂ in the rate formula ✓
Vervang die aantal mol CO₂ in die tempoformule
- Substitute time in the rate formula / *Vervang tyd in die tempoformule* ✓
- Correct final answer/*Korrekte finale antwoord* ✓
- Range/*Gebied*: 0,001 to 0,0013 mol·s⁻¹

$$\begin{aligned} n(\text{CO}_2) &= \frac{m}{M} \\ &= \frac{1,6}{44} \checkmark \text{ (a)} \\ &= \frac{1,6}{44} \checkmark \text{ (b)} \\ &= 0,03636 \text{ mol} \end{aligned}$$

$$\begin{aligned} \text{Rate/Tempo} &= \left(\frac{\Delta n}{\Delta t} \right) \\ &= \left(\frac{0,03636-0}{30-0} \right) \checkmark \text{ (c)} \\ &= \left(\frac{0,03636}{30} \right) \checkmark \text{ (d)} \\ &= 1,21 \times 10^{-3} \text{ (mol} \cdot \text{s}^{-1}) \checkmark \text{ (e)} \end{aligned} \quad (5)$$

5.5 Experiment 1/*Eksperiment 1* ✓ (1)

- 5.6 • High concentration results in more particles per unit volume ✓
 • More effective collisions per unit time/frequency of effective collisions increases ✓
 • Higher rate of reaction ✓/Higher rate of formation of CO₂.
 • *Hoë konsentrasie lei tot meer deeltjies per eenheid volume*
 • *Meer effektiewe botsings per tydseenheid/frekwensie van effektiewe botsings neem toe*
 • *Hoër reaksietempo/Hoër tempo van vorming van CO₂.* (3)

[13]

QUESTION 6/VRAAG 6

- 6.1.1 Rate of forward reaction is equal to rate of reverse reaction. ✓✓
Tempo van voorwaartse reaksie is gelyk aan tempo van terugwaartse reaksie. (2)
- 6.1.2 Addition of a catalyst / Addisie van 'n katalisator ✓✓ (2)
- 6.1.3 (a) Decreased / Toeneem ✓ (1)
- (b) Exothermic / Eksotermies ✓ (1)
- (c)
 - Decrease in temperature favours an exothermic reaction . ✓
 - The forward reaction is favoured ✓/rate of forward reaction decreased less than that of the reverse reaction.
 - *Verlaging in temperatuur bevoordeel 'n eksotermiese reaksie.*
 - *Die voorwaartse reaksie word bevoordeel/tempo van voorwaartse reaksie het minder afgeneem as dié van die terugwaartse reaksie* (2)
- 6.2.1 When the products can be converted back to reactants ✓
Wanneer die produkte terug na reaktante omgeskakel kan word (1)



6.2.2 **CALCULATIONS USING NUMBER OF MOLES/****Mark allocation:**

- (a) USING ratio $\text{NOCl} : \text{NO} : \text{Cl}_2 = 2:2:1$ ✓
 (b) Change in number of moles of NO = equilibrium moles of NO
 Equilibrium number moles of Cl_2 = change in moles of Cl_2
 (c) Equilibrium moles NOCl = initial moles NOCl – change moles of NOCl ✓
 (d) Dividing equilibrium moles NOCl and Cl_2 by 2 AND multiplying equilibrium concentration NO by 2 ✓
 (e) Correct K_c expression (formulae in square brackets) ✓
 (f) Correct substitution of concentrations into K_c correct expression ✓
 (g) Final answer : 0,33 ✓

BEREKENINGE MET GEBRUIK VAN AANTAL MOL**Punttoekenning:**

- (a) *GBRUIK verhouding $\text{NOCl} : \text{NO} : \text{Cl}_2 = 2:2:1$*
 (b) *Verandering in aantal mol NO = ewewigsmol NO*
Ekwilibrumgetal mol Cl_2 = verandering in mol Cl_2
 (c) *Ekwilibrum mol NOCl = aanvanklike mol NOCl – verander mol van NOCl*
 (d) *Deel ewewigsmol NOCl en Cl_2 deur 2 EN vermenigvuldig ewewigkonsentrasie NO met 2*
 (e) *Korrekte K_c -uitdrukking (formules tussen vierkantige hakies)*
 (f) *Korrekte vervanging van konsentrasies in K_c korrekte uitdrukking*
 (g) *Finale antwoord : 0,33*

| | NOCl | NO | Cl₂ | |
|--|-------------|-----------|-----------------------|----------------------------------|
| Initial quantity (mol) <i>Aanvanklike hoeveelheid (mol)</i> | 4 | 0 | 0 | |
| Change <i>Verander (mol)</i> | 1,8 | 1,8 | 0,9 | Ratio ✓ (a) <i>Verhouding</i> |
| Quantity at equilibrium (mol) <i>Hoeveelheid by ewewig (mol)</i> | 2,2 ✓ (c) | 1,8 | 0,9 ✓ (b) | |
| Equilibrium concentration (mol·dm ⁻³) <i>Ekwilibrumkonsentrasie (mol·dm⁻³)</i> | 1,1 | 0,9 | 0,45 | ✓ (d) ÷ & x 2 |

$$K_c = \frac{[\text{NO}]^2[\text{Cl}_2]}{[\text{NOCl}]^2} \quad \checkmark \text{ (e)}$$

$$K_c = \frac{(0,9)^2(0,45)}{1,1} \quad \checkmark \text{ (f)}$$

$$K_c = 0,33 \quad \checkmark \text{ (g)}$$

Wrong K_c expression / *Verkeerde K_c uitdrukking*

Max/Maks: 4/7

No K_c expression followed by correct substitutions / *Geen K_c -uitdrukking gevolg deur korrekte vervangings*

Max/Maks: 6/7

(7)



CALCULATIONS USING CONCENTRATIONS**Mark allocation:**

- (a) Initial concentration of NOCl = initial moles of NOCl \div 2 ✓
 (b) USING ratio $\text{NOCl} : \text{NO} : \text{Cl}_2 = 2:2:1$ ✓
 (c) Change in concentration of NO = equilibrium concentration of NO }
 Equilibrium concentration of Cl_2 = change in concentration of Cl_2 } ✓
 (d) Equilibrium concentration of NOCl = initial concentration of NOCl –
 change in concentration of NOCl ✓
 (e) Correct K_c expression (formulae in square brackets) ✓
 (f) Correct substitution of concentrations into K_c expression ✓
 (g) Final answer : 0,33 ✓

BEREKENINGE DEUR KONSENTRASIES**Puntetoekenning:**

- (a) Aanvanklike konsentrasie van NOCl = aanvanklike mol NOCl \div 2
 (b) GEBRUIK verhouding $\text{NOCl} : \text{NO} : \text{Cl}_2 = 2:2:1$
 (c) Verandering in konsentrasie van NO = ewewigskonsentrasie van NO
 Ekwiilibriumkonsentrasie van Cl_2 = verandering in konsentrasie van Cl_2
 (d) Ekwiilibriumkonsentrasie van NOCl = aanvanklike konsentrasie NOCl –
 verandering in konsentrasie van NOCl
 (e) Korrekte K_c -uitdrukking (formules tussen vierkantige hakies)
 (f) Korrekte vervanging van konsentrasies in K_c uitdrukking
 (g) Finale antwoord: 0,33

| | NOCl | NO | Cl_2 | |
|---|---------------|-------------|---------------|---------------------------|
| Initial concentration (mol·dm ⁻³) Aanvanklike konsentrasie (mol·dm ⁻³) | 2 ✓ (a) | 0 | 0 | |
| Change concentration (mol·dm ⁻³) Verander konsentrasie (mol·dm ⁻³) | 0,9 | 0,9 ✓ (c) | 0,45 | Ratio ✓ (b) Verhouding |
| Equilibrium concentration (mol·dm ⁻³) Ekwiilibriumkonsentrasie (mol·dm ⁻³) | 1,1 ✓ (d) | 0,9 | 0,45 | |

$$K_c = \frac{[\text{NO}]^2[\text{Cl}_2]}{[\text{NOCl}]^2} \quad \checkmark \text{ (e)}$$

$$K_c = \frac{(0,9)^2(0,45)}{1,1} \quad \checkmark \text{ (f)}$$

$$K_c = 0,33 \quad \checkmark \text{ (g)}$$

| | |
|--|---------------|
| Wrong K_c expression / Verkeerde K_c uitdrukking | Max/Maks: 4/7 |
| No K_c expression followed by correct substitutions / Geen K_c -uitdrukking gevolg deur korrekte vervangings | Max/Maks: 6/7 |

(7)
[16]

QUESTION 7/VRAAG 7

7.1.1 The reaction of salt with water ✓✓
Die reaksie van soutwater met water (2)

7.1.2 Acidic/Suur ✓ (1)

7.1.3 $\text{NH}_4^+(\text{aq}) + \text{H}_2\text{O}(\ell) \checkmark \rightarrow \text{NH}_3 + \text{H}_3\text{O}^+(\text{aq}) \checkmark \checkmark$
OR / OF
 $\text{NH}_4^+(\text{aq}) + 2\text{H}_2\text{O}(\ell) \checkmark \rightarrow \text{NH}_4\text{OH} + \text{H}_3\text{O}^+(\text{aq}) \checkmark \checkmark$ (3)

7.2.1 Strong (acid) / Sterk (suur)✓ (1)

7.2.2 Ionises/dissociate completely in water✓ to form a high concentration of H_3O^+ ions. ✓
loneer/dissosieer heeltemal in water om 'n hoë konsentrasie van H_3O^+ ione te vorm. (2)

7.2.3 Methyl orange / Metiel oranje ✓ (1)

7.2.4 Titration of strong acid ✓ and weak base ✓
Titrasie van sterk suur en swak basis (2)

7.2.5 **Marking criteria/Nasienriglyne:**

- Formula / Formule $c = \frac{n}{V}$ ✓
- Substitute / Vervang 0,1 AND / EN 25×10^{-3} in $n = cV$ ✓
- Final answer/Finale antwoord: 0,0025 mol ✓

$$c = \frac{n}{V} \checkmark$$

$$n = 0,1 \times 25 \times 10^{-3} \checkmark$$

$$n = 2,5 \times 10^{-3} \text{ mol} \checkmark / 0,0025 \text{ mol (ACCEPT/AANVAAR: 0,003 mol)} \quad (3)$$



7.2.6 **Marking criteria/Nasiennriglyne:**

- (a) Ratio / *Verhouding* $n(\text{HCl}) = n(\text{NH}_3)$ ✓
 (b) Formula / *Formule* $c = \frac{n}{V}$ / $\text{pH} = -\log[\text{H}_3\text{O}^+]$ ✓
 (c) Substituting $n(\text{NH}_3)$ and volume of $0,04\text{dm}^3$ / *Vervang $n(\text{NH}_3)$ en volume van $0,04\text{dm}^3$*
 (d) Subtraction / *Aftrekking*: $c(\text{HCl})_{\text{initial}} - c(\text{HCl})_{\text{used}}$ ✓✓
 (e) $c(\text{H}_3\text{O}^+) = c(\text{HCl})$ ✓
 (f) Substituting / *Vervanging* $c(\text{H}_3\text{O}^+)$ in $\text{pH} = -\log[\text{H}_3\text{O}^+]$ ✓
 (g) Final answer / *Finale antwoord* = 1,43 ✓

$$n(\text{HCl}) = n(\text{NH}_3) \checkmark (\text{a})$$

$$= 2,5 \times 10^{-3}$$

$$c(\text{HCl}) = \frac{n}{V}$$

$$= \frac{2,5 \times 10^{-3}}{0,04} \checkmark (\text{c})$$

$$= 0,0625 \text{ mol} \cdot \text{dm}^{-3}$$

$$c(\text{HCl})_{\text{excess/oormaat}} = 0,1 - 0,0625 \checkmark \checkmark (\text{d})$$

$$= 0,0375 \text{ mol} \cdot \text{dm}^{-3}$$

ANY/ENIGE 1 ✓(b)

$$c(\text{H}_3\text{O}^+) = c(\text{HCl}) \checkmark (\text{e})$$

$$= 0,0375$$

$$\text{pH} = -\log[\text{H}_3\text{O}^+]$$

$$= -\log(0,0375) \checkmark (\text{f})$$

$$= 1,43 \checkmark (\text{g})$$

(8)
[23]

QUESTION 8/VRAAG 8

8.1 Concentration / *Konsentrasie* = 1 mol·dm⁻³ ✓
Temperature / *Temperatuur* = 25 °C ✓ (2)

8.2.1 Potassium nitrate/KNO₃/Any soluble salt ✓✓
Kaliumnitraat/KNO₃/Enige oplosbare sout (2)

8.2.2 Cl⁻ ✓ (1)

8.2.3 Pt(s) | Cl⁻(aq) | Cl₂(g) ✓ || Au³⁺(aq) | Au(s) ✓ (3)

8.3 OPTION/OPSIE 1

$$E_{\text{cell}}^{\theta} = E_{\text{reduction}}^{\theta} - E_{\text{oxidation}}^{\theta} \checkmark$$

$$0,14 = E_{\text{reduction}}^{\theta} \checkmark - 1,36 \checkmark$$

$$E_{\text{reduction}}^{\theta} = 1,5 \text{ V} \checkmark$$

$$E_{\text{sel}}^{\theta} = E_{\text{reduksie}}^{\theta} - E_{\text{oksidasie}}^{\theta} \checkmark$$

$$0,14 = E_{\text{reduksie}}^{\theta} \checkmark - 1,36 \checkmark$$

$$E_{\text{reduksie}}^{\theta} = 1,5 \text{ V} \checkmark$$


Notes/Aantekeninge:

Accept any other correct formula from the data sheet.

Aanvaar enige ander korrekte formule uit die datablad.

Any other formula using unconventional abbreviations, e.g. $E_{\text{cell}}^{\theta} = E_{\text{O.A.}}^{\theta} - E_{\text{R.}}^{\theta}$, followed by correct substitutions: $\frac{3}{4}$

Enige ander formule wat onkonvensionele afkortings gebruik, bv. $E_{\text{cell}}^{\theta} = E_{\text{O.A.}}^{\theta} - E_{\text{R.}}^{\theta}$, gevolg deur korrekte vervangings: $\frac{3}{4}$

OPTION/OPSIE 2

QUESTION 9/VRAAG 9

9.1 The chemical process in which electrical energy is converted to chemical energy/ use of electrical energy to produce a chemical reaction. ✓✓

Die chemiese proses waarin elektriese energie na chemiese energie omgeskakel word/gebruik van elektriese energie om 'n chemiese reaksie te produseer. (2)

9.2 X ✓ (1)

9.3.1 $2\text{Cl}^- \rightarrow \text{Cl}_2 + 2\text{e}^-$ ✓✓

Marking criteria/Nasienkriteria:

- $2\text{Cl}^- \rightleftharpoons \text{Cl}_2 + 2\text{e}^-$ $\frac{1}{2}$ $\text{Cl}_2 + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-$ $\frac{0}{2}$
 $\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-$ $\frac{0}{2}$ $\text{Cl}_2 + 2\text{e}^- \leftarrow 2\text{Cl}^-$ $\frac{2}{2}$
- Ignore if charge omitted on electron/*Ignoreer as lading op electron weggelaat is*
- If charge (-) omitted on Cl^- /*Indien lading (+) weggelaat is op Cl^- :*
 Max./Maks: $\frac{1}{2}$
 Example/Voorbeeld: $2\text{Cl}^- \rightarrow \text{Cl}_2 + 2\text{e}^-$

(2)

9.3.2 $2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2 + 2\text{OH}^-$ ✓✓

Marking criteria/Nasienkriteria:

- $2\text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{H}_2 + 2\text{OH}^-$ $\frac{1}{2}$ $\text{H}_2 + 2\text{OH}^- \rightleftharpoons 2\text{H}_2\text{O} + 2\text{e}^-$ $\frac{0}{2}$
 $\text{H}_2 + 2\text{OH}^- \leftarrow 2\text{H}_2\text{O} + 2\text{e}^-$ $\frac{2}{2}$ $\text{H}_2 + 2\text{OH}^- \rightarrow 2\text{H}_2\text{O} + 2\text{e}^-$ $\frac{0}{2}$
- Ignore if charge omitted on electron / *Ignoreer as lading op elektron weggelaat is*
- If charge (-) omitted on OH^-
Indien lading (-) op OH^- weggelaat is Max./Maks: $\frac{1}{2}$
 Example / Voorbeeld: $2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2 + 2\text{OH}^-$

(2)



9.4

Marking criteria/Nasienriglyne:

- (a) Substitute / Vervang $0,5 \times 18\,000$ ✓
 (b) Dividing charge by / Deel lading deur $1,6 \times 10^{-19}$ ✓
 (c) Dividing number of electrons by Avogadro's constant. ✓
Deel die aantal elektrone deur Avogadro se konstante.
 (d) Using ratio / Gebruik verhouding $n(\text{Cl}_2) : n(e^-) = 1:2$ ✓
 (e) Substituting / Vervanging $n(\text{Cl}_2)$ and / en 22,4 in $n = \frac{V}{V_m}$ ✓
 (f) Final answer / Finale antwoord $1,04 \text{ dm}^3$ ✓

$$Q = I\Delta t$$

$$= 0,5 \times 18\,000 \text{ ✓(a)}$$

$$= 9\,000 \text{ C}$$

$$n = \frac{Q}{q_e}$$

$$n = \frac{9\,000}{1,6 \times 10^{-19}} \text{ ✓(b)}$$

$$= 5,625 \times 10^{22} \text{ electrons / elektrone}$$

$$n(\text{electrons}) = \frac{5,625 \times 10^{22}}{6,02 \times 10^{23}} \text{ ✓(c)}$$

$$= 0,093 \text{ mol}$$

$$n(\text{Cl}_2) = \frac{1}{2}n(e^-)$$

$$= 0,093 \div 2 \text{ ✓(d)}$$

$$= 0,0465 \text{ mol}$$

$$n = \frac{V}{V_m}$$

$$V = 0,0465 \times 22,4 \text{ ✓(e)}$$

$$= 1,04 \text{ dm}^3 \text{ ✓(f)}$$

(6)

[13]

TOTAL/TOTAAL: 150