

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



**SA EXAM
PAPERS**
SA EXAM
PAPERS



LIMPOPO
PROVINCIAL GOVERNMENT
REPUBLIC OF SOUTH AFRICA

DEPARTMENT OF EDUCATION

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

**PHYSICAL SCIENCES: CHEMISTRY (P2)
SEPTEMBER 2024**

MARKS: 150

TIME: 3 HOURS

This paper consists of 14 pages and 4 data sheets



INSTRUCTIONS AND INFORMATION

1. The question paper consists of nine questions. Answer ALL the questions.
2. Start EACH question on a NEW page.
3. Number your answers correctly according to the numbering system used in this question paper.
4. Leave ONE line between two sub-questions, e.g., between QUESTION 2.1 and QUESTION 2.2.
5. A non-programmable calculator may be used.
6. Appropriate mathematical instruments may be used.
7. Show ALL formulae and substitutions in ALL calculations.
8. Round off your FINAL numerical answers to a minimum of TWO decimal places.
9. Give brief motivations, discussions, etc. where required.
10. You are advised to use the attached data sheets.
11. Write neatly and legibly.



NSC

QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Four options are provided as possible answers to the following questions. Choose the answer and write only the letter (A–D) next to the question number (1.1 – 1.10), for example 1.10 E


1.1 What is the IUPAC name for the compound $\text{CH}_3\text{CH}_2\text{CHFCH}_3$?

- A. 2-fluorobutane
- B. 3-fluorobutane
- C. fluoro-3-butane
- D. fluorobutane

(2)


Base your answers to questions 1.2 and 1.3 on the graph below.

Boiling points of selected alcohols at 101,3 kPa



NSC

- 1.2 What is represented by the number "1" in the IUPAC name for three of these alcohols?
- The number of isomers for each alcohol.
 - The number of -OH groups for each carbon atom in each alcohol molecule.
 - The location of an -OH group on one end of the carbon chain in each alcohol molecule.
 - The location of an -OH group in the middle of the carbon chain in each alcohol molecule. (2)
- 1.3 What can be concluded from this graph?
- At 101,3 kPa, water has a higher boiling point than butan-1-ol.
 - At 101,3 kPa, water has a lower boiling point than ethanol.
 - The greater the number of carbon atoms per alcohol molecule, the lower the boiling point of the alcohol.
 - The greater the number of carbon atoms per alcohol molecule, the higher the boiling point of the alcohol. (2)
- 1.4 Which formula represents the product of the addition reaction between ethene and chlorine, Cl_2 ?



(2)



NSC

- 1.5 A reaction is most likely to occur when the reacting particles collide with proper orientation and proper:
- A. Charge
 - B. Energy
 - C. Mass
 - D. Volume
- (2)
- 1.6 Catalysts can increase the rate of a chemical reaction by providing.....
- A. an alternate reaction pathway with a higher activation energy.
 - B. the same reaction pathway with a higher activation energy.
 - C. an alternate reaction pathway with a lower activation energy.
 - D. the same reaction pathway with a lower activation energy.
- (2)
- 1.7 Given the equation representing a system at equilibrium in a sealed, rigid container:
- $$2\text{HI}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \text{I}_2(\text{g}) + \text{energy}$$
- Increasing the temperature of the system causes the concentration of.....
- A. HI to increase.
 - B. H₂ to increase.
 - C. HI to remain constant.
 - D. H₂ to remain constant.
- (2)
- 1.8 How will the pH change if there is a tenfold increase in the hydronium ion concentration?
- A. A decrease of one unit of pH.
 - B. A decrease of 10 units of pH.
 - C. An increase of one unit of pH.
 - D. An increase of 10 units of pH.
- (2)



NSC

1.9 Which positive ion must be present in an aqueous solution of an Arrhenius acid?

A. Na^+


B. NH_4^+

C. OH^-

D. H_3O^+

(2)

1.10 Consider the reaction below.



In this reaction, electrons are transferred from:

A. Al to H^+

B. H to H^+

C. Al to Al^{3+}

D. H to Al

(2)

[20]

NSC

QUESTION 2 (Start on a new page.)

Carbon is the basic building block of organic compounds that recycles through the earth's air, water, soil, and living organisms including human beings.

2.1 Discuss ONE special property of carbon that makes it possible to form a variety of bonds. (2)

2.2 The IUPAC name of an organic compound is:


1-chloro-2,3-dimethylpentane

2.2.1 Write down the GENERAL FORMULA of the homologous series to which this compound belongs. (1)

2.2.2 Draw the condensed STRUCTURAL formula of this compound. (3)

2.2.3 Is this compound SATURATED or UNSATURATED? (1)

2.3 The organic compound below ($\text{CH}_3\text{COCH}_2\text{CH}_2\text{CH}_3$) has one POSITIONAL isomer and one FUNCTIONAL isomer.



2.3.1 Define the term *positional isomer*. (2)


For this compound, write down the:

2.3.2 Name of the functional group present in this compound. (1)

2.3.3 IUPAC name of its POSITIONAL isomer. (2)

2.3.4 Structural formula of its FUNCTIONAL isomer. (2)

Write down the condensed structural formula of an alcohol.



2.4.1 IS THIS a PRIMARY, SECONDARY or TERTIARY alcohol? Give a reason for the answer. (2)



Write down the condensed structural formula of the compound. (2)

[18]



QUESTION 3 (Start on a new page.)

The table below shows data collected for three organic compounds, represented by the letters **A**, **B** and **C**, during a practical investigation:

	Organic compound	Mr (g.mol ⁻¹)	Boiling Point (°C)
A	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$ 	58	-0,5
B	$\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ 	60	97
C	CH_3COOH 	60	118


- 3.1 Which variable was controlled during this investigation? (1)
- 3.2 Identify the:
- 3.2.1 Dependent variable. (1)
- 3.2.2 Independent variable. (1)
- 3.3 Consider compound **A**:
- 3.3.1 2-methylpropane is an isomer of compound **A**.
- Predict whether the boiling point of 2-methylpropane will be HIGHER THAN, LOWER THAN or THE SAME as the boiling point of compound **A**. (1)
- 3.3.2 Explain your prediction in QUESTION 3.3.1. (3)
- 3.4 Define the term *vapor pressure*. (2)
- 3.5 Refer to intermolecular forces and energy to explain why compound **B** will have a higher vapour pressure than compound **C** at 20 °C. (3)

(3)
[12]



QUESTION 4 (Start on a new page.)

Various organic reactions are shown in the flow diagram.




- 4.1 Name the type of reaction illustrated by:
- 4.1.1 A (1)
- 4.1.2 B (1)
- 4.1.3 C (1)
- 4.1.4 D (1)
- 4.2 Use condensed structural formulas and write a balanced equation for reaction C. (4)
- 4.3 Write down the structural formula for compound X. (2)
- 4.4 Compound X undergoes combustion in the presence of oxygen. Write down the molecular formulas of the two products that form. (2)
- 4.5 Use structural formulas to write a balanced equation for the reaction to produce product Y. (3)
- Instead of heating concentrated potassium hydroxide under reflux during the formation of Y, dilute sodium hydroxide is used.*
- 4.6 Write the IUPAC name of the organic compound that will form. (2)
- 4.7 To which homologous series does product Z belong? (1)

- 4.8 Write down the IUPAC name of product Z if ethanoic acid was used during the reaction.

(2)
[20]

QUESTION 5 (Start on a new page.)

Graph B represents the Maxwell-Boltzmann energy distribution curve for a reaction mixture at 300°C. Area X represents the number of molecules in the mixture that have enough kinetic energy for the reaction to take place.



- 5.1 Give a term for the “*minimum energy needed for a reaction to take place*”, as indicated by Y. (1)
- 5.2 The temperature of the mixture is now increased to 500 °C.
- 5.2.1 Which ONE of graph A or C represents the distribution curve of the mixture at this higher temperature? (1)
- 5.2.2 Give a reason for the answer. (1)
- 5.2.3 Use the collision theory to explain how this increase in temperature will influence the rate of the reaction. (4)
- 5.3 A catalyst is added to the mixture.
- 5.3.1 Write down the definition of a *catalyst*. (2)
- 5.3.2 How will the above-mentioned action affect the size of area X (area to the right of vertical line Y)? Write down only INCREASES, DECREASES or REMAINS THE SAME. (1)
- 5.3.3 Explain your answer to QUESTION 5.3.2. (2)


[12]

NSC


QUESTION 6 (Start on a new page.)

2,0 g of CaCO_3 is sealed in an evacuated $1,0 \text{ dm}^3$ metal flask and a pressure gauge is connected to the flask, in order to determine the equilibrium constant for the decomposition of calcium carbonate (CaCO_3).

The flask is heated to a 800°C at which equilibrium was reached.



The graph obtained for pressure versus time for the decomposition of calcium carbonate is shown below.



- 6.1 Define the term *reaction rate*. (2)
- 6.2 How does the rate of the reverse reaction change from t_0 to t_1 ? (2)
- 6.3 What is the reason for the horizontal line between t_2 and t_3 ? (1)
- 6.4 Draw a sketch graph to show how the mass of CaCO_3 changes for the period t_0 to t_3 . (3)
- 6.5 During a power failure the temperature of the oven drops to 500°C . What effect (only write INCREASES, DECREASES or STAYS THE SAME) does this decrease in temperature have on the following:
- 6.5.1 The rate of the forward reaction. (1)
- 6.5.2 The concentration of CO_2 . (1)
- 6.5.3 The value of K_c . (1)
- 6.6 Give a reason for your answer to QUESTION 6.5.3. (4)
- 6.7 When equilibrium was established at 800°C , the concentration of CO_2 present in the flask was $1,4 \times 10^{-2} \text{ mol} \cdot \text{dm}^{-3}$. Calculate the equilibrium constant (K_c) at 800°C for this reaction. (2)

[17]

QUESTION 7 (Start on a new page)

- 7.1 Sulphuric acid is a diprotic
- 7.1.1 Define an *acid* in terms of the Lowry-Brønsted theory. (2)
- 7.1.2 Give a reason why sulphuric acid is a *diprotic acid*. (1)
- 7.2 The hydrogen carbonate ion can act as both an acid and a base. It reacts with water according to the following balanced equation:
- $$\text{HCO}_3^-(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{H}_2\text{CO}_3(\text{aq}) + \text{OH}^-(\text{aq})$$
- 7.2.1 Write down ONE word for the underlined phrase. (1)
- 7.2.2 $\text{HCO}_3^-(\text{aq})$ acts as a base in the above reaction.
Write down the formula of the conjugate acid of $\text{HCO}_3^-(\text{aq})$. (1)
- 7.3 A chemist spills sulphuric acid of concentration $6 \text{ mol}\cdot\text{dm}^{-3}$ from a flask on a laboratory bench. She neutralises the spilled acid by sprinkling sodium hydrogen carbonate powder onto it. The reaction that takes place is: (Assume that the H_2SO_4 ionises completely.)
- $$\text{H}_2\text{SO}_4(\text{aq}) + 2\text{NaHCO}_3(\text{s}) \rightarrow \text{Na}_2\text{SO}_4(\text{aq}) + 2\text{H}_2\text{O}(\text{l}) + 2\text{CO}_2(\text{g})$$
- The fizzing, due to the formation of carbon dioxide, stops after the chemist has added 27 g sodium hydrogen carbonate to the spilled acid.
- 7.3.1 Calculate the volume of sulphuric acid that spilled. Assume that all the sodium hydrogen carbonate reacts with all the acid. (6)
- The chemist now dilutes some of the $6 \text{ mol}\cdot\text{dm}^{-3}$ sulphuric acid solution in the flask to $0,1 \text{ mol}\cdot\text{dm}^{-3}$.
- 7.3.2 Calculate the volume of the $6 \text{ mol}\cdot\text{dm}^{-3}$ sulphuric acid solution needed to prepare 1 dm^3 of the dilute acid. (2)
- During a titration 25 cm^3 of the $0,1 \text{ mol}\cdot\text{dm}^{-3}$ sulphuric acid solution is added to an Erlenmeyer flask and titrated with a $0,1 \text{ mol}\cdot\text{dm}^{-3}$ sodium hydroxide solution.
- 7.3.3 The chemist uses bromothymol blue as indicator.
What is the purpose of this indicator? (1)
- 7.3.4 Calculate the pH of the solution in the flask after the addition of 30 cm^3 of sodium hydroxide. The endpoint of the titration is not yet reached at this point. (8)
- [22]




QUESTION 8 (Start on a new page.)

NSC

A learner wants to investigate the effect of the area of the metal plates used as electrodes in a galvanic cell on the emf of the cell.

The learner sets up the following cell under standard conditions and measures the emf.




- 8.1 Which electrode will show an increase in mass when this cell is functioning? (1)
- 8.2 Write the equation for the half-reaction occurring at the anode. (2)
- 8.3 Calculate the emf that the learner will read on the voltmeter. (4)
- 8.4 Name TWO variables that should be controlled during this investigation. (2)
- 8.5 The learner now replaces the two metal plates with ones of larger surface area and takes the voltmeter reading again.
- 8.5.1 How would you expect the new emf to compare with the one calculated in QUESTION 8.3? (Only write SMALLER THAN, LARGER THAN or EQUAL TO.) (1)
- 8.5.2 Explain your answer to QUESTION 8.5.1. (2)
- 8.6 The learner now connects a resistor of low resistance across terminals A and B. The learner notes that the reading on the voltmeter immediately drops.
- 8.6.1 Give a reason for this observation. (1)
- 8.6.2 After some time the learner observes a further gradual drop in the reading on the voltmeter. Give a reason for this observation. (1)

[14]

QUESTION 9 (Start on a new page.)

Electroplating is an electrolytic process that can be used to coat metal objects with a less reactive metal. The diagram below shows an electroplating cell that includes a power source connected to a copper rod and a bracelet made from a different metal. The rod and bracelet are in an aqueous copper (II) sulphate solution.



- 9.1 Define the term *electrolyte*. (2)
- 9.2 Is the electrolytic process endothermic or exothermic? (1)
- 9.3 Which electrode, the BRACELET or the Cu rod is the cathode? (1)
- 9.4 Initially the $\text{CuSO}_4(\text{aq})$ has a blue colour.
- 9.4.1 How will the intensity of the blue colour change whilst the cell is functioning? Write down INCREASES, DECREASES or REMAINS THE SAME. (1)
- 9.4.2 Give a reason for the answer in QUESTION 9.4.1. (2)
- 9.4.3 Write down the half-reaction that takes place at the pure copper electrode. (2)
- 9.4.4 If the bracelet is plated with 0,86 g of copper, how many moles of electrons were transferred during the process. (6)

[15]**TOTAL: 150**

GEGEWENS VIR FISIESTE WETENSKAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)
DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)

TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESTE KONSTANTES

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Avogadro's constant <i>Avogadro-konstante</i>	N_A	$6,02 \times 10^{23} \text{ mol}^{-1}$
Charge on electron <i>Lading op elektron</i>	e	$-1,6 \times 10^{-19} \text{ C}$
Electron mass <i>Elektronmassa</i>	m_e	$9,11 \times 10^{-31} \text{ kg}$
Molar gas volume at STP <i>Molêre gasvolume by STD</i>	V_m	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard pressure <i>Standaarddruk</i>	p^θ	$1,013 \times 10^5 \text{ Pa}$
Standard temperature <i>Standaardtemperatuur</i>	T^θ	273 K

TABLE 2: FORMULAE/TABEL 2: FORMULES

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ or/of $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$ $\frac{CaVa}{CbVb} = \frac{na}{nb}$
$K_w = [H_3O^+][OH^-] = 1 \times 10^{-14}$ at / by 298 K pH = $-\log[H_3O^+]$	
$E_{cell}^\theta = E_{cathode}^\theta - E_{anode}^\theta$ / $E_{sel}^\theta = E_{katode}^\theta - E_{anode}^\theta$	
or / of	
$E_{cell}^\theta = E_{reduction}^\theta - E_{oxidation}^\theta$ / $E_{sel}^\theta = E_{reduksie}^\theta - E_{oksidasie}^\theta$	
or / of	
$E_{cell}^\theta = E_{oxidising\ agent}^\theta - E_{reducing\ agent}^\theta$ / $E_{sel}^\theta = E_{oksideermiddel}^\theta - E_{reduseermiddel}^\theta$	

$$I = \frac{Q}{\Delta t}$$

$$n = \frac{Q}{qe} \quad \text{where } n \text{ is the number of electrons}$$



NSC

THE PERIODIC TABLE OF ELEMENTS II DIE PERIODIEKE TABEL VAN ELEMENTE

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18																																	
(I)	(II)	KEY/SLEUTEL										(III)	(IV)	(V)	(VI)	(VII)	(VIII)																																	
Atomic number		Electronegativity		Symbol		Approximate relative atomic mass		Atomic number		Electronegativity		Symbol		Approximate relative atomic mass		Atomic number																																		
1 H 1	3 Li 7	4 Be 9	11 Na 23	19 K 39	37 Rb 85	55 Cs 133	87 Fr 223	2 He 4	10 Ne 20	18 Ar 40	36 Kr 84	54 Xe 131	86 Rn 222	102 No 259	118 Og 294	136 Lv 304	172 Ts 352	118 Og 294																																
2 He 4	10 Ne 20	18 Ar 40	36 Kr 84	54 Xe 131	86 Rn 222	118 Og 294	136 Lv 304	172 Ts 352	118 Og 294	136 Lv 304	172 Ts 352	118 Og 294	136 Lv 304	172 Ts 352	118 Og 294	136 Lv 304	172 Ts 352	118 Og 294																																
21 Sc 45	22 Ti 48	23 V 51	24 Cr 52	25 Mn 55	26 Fe 56	27 Co 59	28 Ni 59	29 Cu 63,5	30 Zn 65	31 Ga 70	32 Ge 73	33 As 75	34 Se 79	35 Br 80	36 Kr 84	37 Rb 85	38 Sr 88	39 Y 89	40 Zr 91	41 Nb 92	42 Mo 96	43 Tc 98	44 Ru 101	45 Rh 103	46 Pd 106	47 Ag 108	48 Cd 112	49 In 115	50 Sn 119	51 Sb 122	52 Te 128	53 I 127	54 Xe 131	55 Cs 133	56 Ba 137	57 La 139	58 Ce 140	59 Pr 141	60 Nd 144	61 Pm 145	62 Sm 150	63 Eu 152	64 Gd 157	65 Tb 159	66 Dy 163	67 Ho 165	68 Er 167	69 Tm 169	70 Yb 173	71 Lu 175
72 Hf 178	73 Ta 181	74 W 184	75 Re 186	76 Os 190	77 Ir 192	78 Pt 195	79 Au 197	80 Hg 201	81 Tl 204	82 Pb 207	83 Bi 209	84 Po 209	85 At 210	86 Rn 222	87 Fr 223	88 Ra 226	89 Ac	90 Th 232	91 Pa 231	92 U 238	93 Np 237	94 Pu 244	95 Am 243	96 Cm 247	97 Bk 247	98 Cf 251	99 Es 252	100 Fm 257	101 Md 288	102 No 289	103 Lr 260	104 Rf 261	105 Db 262	106 Sg 266	107 Bh 264	108 Hs 277	109 Mt 268	110 Ds 271	111 Rg 272	112 Cn 285	113 Nh 284	114 Fl 289	115 Mc 288	116 Lv 293	117 Ts 294	118 Og 294				