

SA's Leading Past Year

Exam Paper Portal


S T U D Y

You have Downloaded, yet Another Great
Resource to assist you with your Studies ☺

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexamapers.co.za





basic education


Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

SENIOR CERTIFICATE EXAMINATIONS/ NATIONAL SENIOR CERTIFICATE EXAMINATIONS **SENIORSERTIFIKAAT-EKSAMEN/** **NASIONALE SENIORSERTIFIKAAT-EKSAMEN**

PHYSICAL SCIENCES: CHEMISTRY (P2)
FISIESE WETENSKAPPE: CHEMIE (V2)

MAY/JUNE/MEI/JUNIE 2024

MARKING GUIDELINES/NASIENRIGLYNE



MARKS/PUNTE: 150

These marking guidelines consist of 20 pages./
Hierdie nasienriglyne bestaan uit 20 bladsye.

Approved
Moek
DBE Moderator

Approved
↓
DBE moderator

Approved
Jabulile
Umahle Ext. Mod

Approved!
Vulindlela.
Umthwisi Ext. moderator



QUESTION 1/VRAAG 1


- 1.1 A ✓✓ (2)
 1.2 C ✓✓ (2)
 1.3 A ✓✓ (2)
 1.4 D ✓✓ (2)
 1.5 D ✓✓ (2)
 1.6 C ✓✓ (2)
 1.7 B ✓✓ (2)
 1.8 A ✓✓ (2)
 1.9 B ✓✓ (2)
 1.10 B ✓✓ (2)
- [20]**

QUESTION 2/VRAAG 2


- 2.1 Organic compounds that consist of hydrogen and carbon only. ✓✓ (2 or 0)
Organiese verbindings wat slegs uit waterstof en koolstof bestaan. (2 or 0) (2)
- 2.2.1 C and/en E ✓ (1)
- 2.2.2 D and/en H ✓✓ (2 or/of 0) (2)
- 2.2.3 A ✓ (1)

2.3	Marking criteria/Nasienkriteria:	IF/INDIEN:
2.3.1	<ul style="list-style-type: none"> Functional group. ✓ <i>Funksionele groep.</i> Whole structure correct. ✓ <i>Hele struktuur korrek.</i> 	<ul style="list-style-type: none"> More than one functional group/wrong functional group: <i>Meer as een funksionele groep/foutiewe funksionele groep:</i> $\frac{0}{2}$ If condensed structural formulae used/Indien gekondenseerde struktuurformules gebruik: <i>Max/Maks. $\frac{1}{2}$</i>

N.
Vraag
1
K



OR/OF




(2)

2.3.2 C_nH_{2n+2} ✓

(1)

2.3.3

**Marking criteria/Nasienkriteria:**

- Functional group $-C\equiv C-$. ✓
- Funksionele groep $-C\equiv C-$.
- Whole structure correct. ✓
- Hele struktuur korrek.

IF/INDIEN

- More than one functional group/wrong functional group:
Meer as een funksionele groep/foutiewe funksionele groep: 0/2
- If condensed structural formulae used/Indien gekondenseerde struktuurformules gebruik:
Max/Maks. 1/2

(2)

2.4.1 3-ethylhex-3-ene ✓✓✓/3-ethyl-3-hexene/3-etielheks-3-een/3-etiel-3-hekseen

Marking criteria:

- Correct stem i.e. hexene. ✓
- Substituent (ethyl) correctly identified. ✓
- IUPAC name completely correct including numbering, sequence, hyphens and commas. ✓

Nasienkriteria:

- Korrekte stam d.i. hekseen. ✓
- Substituent (etiel) korrek geïdentifiseer. ✓
- IUPAC-naam heeltemal korrek insluitende nommering, volgorde, koppeltekens en kommas. ✓

(3)

2.4.2 2,5-dichloro-2,4-dimethylhexane ✓✓✓/ 2,5-dichloro-2,4-dimetielheksaan

Marking criteria:

- Correct stem i.e. hexane. ✓
- All substituents (dichloro and dimethyl) correctly identified. ✓
- IUPAC name completely correct including numbering, sequence, hyphens and commas. ✓

Nasienkriteria:

- Korrekte stam d.i. heksaan. ✓
- Alle substituente (dichloro en dimetiel) korrek geïdentifiseer. ✓
- IUPAC-naam heeltemal korrek insluitende nommering, volgorde, koppeltekens en kommas. ✓

(3)

2.4.3 2,2-dimethylpropanal ✓/dimethylpropanal2,2-dimetielpropanaal/dimetielpropanaal

(2)

NOTE/NOTA:

2,2-dimethyl✓propan-1-al (Max/Maks: 1/2)

2.5

Marking criteria/Nasienkriteria:

- Correct molecular formula: C₇H₁₆ ✓
Korrekte molekulêre formula: C₇H₁₆
- Correct molecular formula of inorganic reactant and products. ✓
Korrekte molekulêre formule vir die anorganiese reaktans en produkte.
- Balancing/Balansering ✓

**Notes/Aantekeninge:**

- Ignore double arrows and phases./Ignoreer dubbelpyle en fases.
- Marking rule 6.3.10/Nasienreël 6.3.10.
- If condensed structural formulae used:/Indien gekondenseerde struktuurformules gebruik: Max/Maks. $\frac{2}{3}$
- **ACCEPT:** multiple coefficients for this exam.
AANVAAR: veelvoude van koëffisiënte vir hierdie eksamen.

(3)
[22]**QUESTION 3/VRAAG 3**

3.1

Marking criteria/Nasienkriteria

If any one of the underlined key phrases in the **correct context** is omitted, deduct 1 mark./Indien enige van die onderstreepte frases in die **korrekte konteks** uitgelaat is, trek 1 punt af.

The underlined phrases must be in the correct context. / Die onderstreepte frases moet in die korrekte konteks wees.

The temperature at which the vapour pressure (of a substance) equals atmospheric pressure. ✓✓


Die temperatuur waarby die dampdruk (van die stof) gelyk is aan atmosferiese druk.

(2)

3.2

C ✓

(1)



N.
Vif
Ha
K

3.3

Marking criteria:

- Compare structures. ✓
- Compare the strength of intermolecular forces. ✓
- Compare the energy required to overcome intermolecular forces. ✓

Nasienkriteria:

- Vergelyk strukture. ✓
- Vergelyk die sterkte van intermolekulêre kragte. ✓
- Vergelyk die energie benodig om intermolekulêre kragte te oorkom. ✓

Accept: IMF for this exam/**Aanvaar:** IMK vir hierdie eksamen**A/CH₃CH₂CH₂CH₂Cl /1-chlorobutane****• Structure:**Longer chain length/larger surface area (over which intermolecular forces act). ✓**• Intermolecular forces:**Stronger/more intermolecular forces/Van der Waals forces/London forces/dipole-dipole forces. ✓**• Energy:**More energy needed to overcome or break intermolecular forces/Van der Waals forces/dipole-dipole forces. ✓**OR****B/CH₃CH(CH₃)CH₂Cl/1-chloro-2-methylpropane****• Structure:**Shorter chain length / branched / compact / more spherical / smaller surface area (over which intermolecular forces act). ✓**• Intermolecular forces:**Weaker/less intermolecular forces/Van der Waals forces/London forces/dipole-dipole forces. ✓**• Energy:**Less energy needed to overcome or break intermolecular forces/Van der Waals forces/dipole-dipole forces. ✓**A/CH₃CH₂CH₂CH₂Cl /1-chlorobutanaan****• Struktuur:**Langer kettinglengte/groter oppervlak (waaroor intermolekulêre kragte werk). ✓**• Intermolekulêre kragte:**Sterker/meer intermolekulêre kragte/Van der Waalskragte/Londonkragte/dipool-dipoolkragte. ✓**• Meer energie benodig om intermolekulêre kragte/Van der Waalskragte/Londonkragte/dipool-dipoolkragte te oorkom/breek.** ✓**OF****B/CH₃CH(CH₃)CH₂Cl/1-chloro-2-metielpropaan****• Struktuur:**Korter kettinglengte / vertak / kompak / meer sferies / kleiner oppervlak (waaroor intermolekulêre kragte werk). ✓**• Intermolekulêre kragte:**Swakker/minder intermolekulêre kragte/Van der Waalskragte/Londonkragte/dipool-dipoolkragte. ✓**• Energie:**Minder energie benodig om intermolekulêre kragte/Van der Waalskragte/Londonkragte/dipool-dipoolkragte te oorkom/breek. ✓

(3)

N.
Vis
Vla
MK

3.4.1 75 (°C) ✓

(1)

3.4.2

Marking criteria:

- Compare the strength of intermolecular forces. ✓
- Compare the energy required to overcome intermolecular forces. ✓

Nasienkriteria:

- Vergelyk die sterkte van intermolekulêre kragte. ✓
- Vergelyk die energie benodig om intermolekulêre kragte te oorkom. ✓

• Intermolecular forces:

C (CH₃CH₂CH₂CH₂OH/butanol) has stronger intermolecular forces than D (CH₃CH₂CH₂CHO/butanal). ✓

• Energy:

More energy needed to overcome or break intermolecular forces. ✓

Accept: Boiling point of C will be more (in relation to C and D/118°C vs 75°C).

OR**• Intermolecular forces:**

D (CH₃CH₂CH₂CHO/butanal) has weaker intermolecular forces than C (CH₃CH₂CH₂CH₂OH/butanol)

• Energy:

Less energy is needed to overcome or break intermolecular forces.

Accept: Boiling point of D will be less (in relation to C and D/118°C vs 75°C).

OR**• Intermolecular forces:**

A (CH₃CH₂CH₂CH₂Cl) is a more polar molecule than D (CH₃CH₂CH₂CHO) increasing the intermolecular forces

• Energy:

More energy is needed to overcome or break intermolecular forces.

Accept: Boiling point of D will be less (in relation to A and D).

OR**• Intermolecular forces:**

Electron density of A (CH₃CH₂CH₂CH₂Cl) is greater than D (CH₃CH₂CH₂CHO) increasing the intermolecular forces

• Energy:

More energy is needed to overcome or break intermolecular forces.

Accept: Boiling point of D will be less (in relation to A and D).

• Intermolekulêre kragte:

C (CH₃CH₂CH₂CH₂OH/butanol) het sterker intermolekulêre kragte as D (CH₃CH₂CH₂CHO/butanaal). ✓

• Meer energie benodig om intermolekulêre kragte te oorkom/breek. ✓
Aanvaar: Kookpunt van D sal minder wees (met betrekking tot C en D)**OF****• Intermolekulêre kragte:**

D (CH₃CH₂CH₂CHO/butanaal) het swakker intermolekulêre kragte as C (CH₃CH₂CH₂CH₂OH/butanol).


• Minder energie benodig om intermolekulêre kragte te oorkom/breek.
Aanvaar: Kookpunt van C sal meer wees (met betrekking tot C en D)

N.

Vuf

(b)

MK



OF

- **Intermolekulêre kragte:**
A ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$) is 'n meer polêre molekule as D wat sterker intermolekulêre kragte tot gevolg het.
- Meer energie benodig om intermolekulêre kragte te oorkom/breek.
Aanvaar: Kookpunt van D sal minder wees (met betrekking tot A en D)

OF

- **Intermolekulêre kragte:**
Elektrondigtheid van A ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$) is groter wat sterker intermolekulêre kragte tot gevolg het.
- Meer energie benodig om intermolekulêre kragte te oorkom/breek.
- Aanvaar: Kookpunt van D sal minder wees (met betrekking tot A en D)

(2)

3.5 Decreases/Neem af ✓

(1)

[10]

QUESTION 4/VRAAG 4

4.1

4.1.1 (Concentrated) sulphuric acid/ $\text{H}_2\text{SO}_4(\text{aq})$ ✓
(Gekonsentreerde) swawelsuur

(1)

4.1.2 Esterification / Condensation ✓ / Veresterung / Esterifikasie / Kondensasie

(1)

4.1.3 ANY TWO/ENIGE TWEE:

- Alcohol/methanol/reactant is flammable/catches fire easily. ✓
Alkohol/metanol/reaktans is vlamaar/slaan maklik aan die brand.
- To heat evenly/A steady/controlled/gradual increase in temperature. ✓
Om eweredig/gekontroleerd/gelydelik te verhit/n Eweredige toename in temperatuur.
- Alcohol/methanol will evaporate too quickly/is volatile.
Alkohol/metanol sal te vinnig verdamp/is vlugtig.

(2)

N.
Vis
Ho
K.



4.1.4

Marking criteria:


- Whole structural formula correct for propanoic acid. ✓
- Whole structural formula correct for methanol. ✓
- Functional group of ester correct. ✓
- Whole structural formula of ester correct. ✓
- H₂O ✓

Nasienkriteria:

- Hele struktuurformule vir propanoësuur korrek. ✓
- Hele struktuurformule vir metanol korrek. ✓
- Funksionele groep van ester korrek. ✓
- Hele struktuurformule van ester korrek. ✓
- H₂O ✓

IF/INDIEN

- Any error e.g. omission of all H atoms, condensed or semi structural formula/*Enige fout bv. weglating van alle H-atome, gekondenseerde of semi-struktuurformule:* Max/Maks. $\frac{2}{5}$ (*Functional group, H₂O/Funksionele groep, H₂O*)
- Any additional reactants or products /*Enige addisionele reaktanse of produkte:* Subtract 1 mark./*Trek 1 punt af.*
- Molecular formulae used:/*Molekuläre formule gebruik:* Max/Maks. $\frac{1}{5}$ (*water*)
- No arrows: The first two structures given are considered as reactants and can be marked/*Geen pyltjie: die eerste twee strukture geskryf, word beskou as reaktanse en kan gemerk word.*



(5)

4.1.5 Methyl ✓ propanoate ✓ /Metielpropanoat (2)

4.2.1 Hydrogen/H₂ ✓ /Waterstof(gas) (1)4.2.2 3,3-dimethyl✓but-1-ene✓/3,3-dimethyl-1-butene
3,3-dimetiel but-1-een/3,3-dimetiel-1-buteen (2)

4.2.3 elimination OR dehydrohalogenation ✓ eliminasie OF dehidrohalogenering (1)

4.2.4 H₂SO₄/H₃PO₄ OR/OF Sulphuric acid/Phosphoric acid ✓
Swawelsuur/Fosforsuur (1)4.2.5 3,3-dimethyl✓butan-2-ol✓/3,3-dimethyl-2-butanol
3,3-dimetiel butan-2-ol/3,3-dimetiel-2-butanol (2)

4.2.6 Addition/hydration ✓ Addisie/hidrasie (1)

4.2.7 Secondary ✓ /Sekondêr (1)

(1)

[20]

QUESTION 5/VRAAG 5

5.1.1 Exothermic/Eksotermies ✓

Lower (potential) energy of the products than reactants. $\Delta H < 0/\Delta H$ negative /
 $\Delta H = -121,7 \text{ kJ}/\text{More energy is released than absorbed.} \checkmark$

*Laer (potensiële) energie van produkte as die reaktanse./ $\Delta H < 0/\Delta H$ negatief /
 $\Delta H = -121,7 \text{ kJ}/\text{Meer energie word afgegee as wat opgeneem is.}$*

(2)

5.1.2 (The number of) particles with sufficient/enough (kinetic) energy (with a catalyst) OR $E_k \geq E_A$ (which can undergo effective collisions.) ✓
(Die hoeveelheid) deeltjies met genoeg/voldoende (kinetiese) energie (met 'n katalisator) OF $E_k \geq E_A$ (om effektiewe botsings te ondergaan).

(1)

5.1.3 $240,8 - 208,2 \checkmark = 32,6 \text{ (kJ)}$ ✓

(2)

IF: only answer award 2 marks//**INDIEN:** slegs antwoord gee 2 punte

5.2

5.2.1 Decreases/Afneem ✓

(1)

5.2.2 Remains the same/Bly dieselfde ✓

(1)

5.2.3 Remains the same/Bly dieselfde ✓

(1)

5.3.1 Concentration (of sulphuric acid/ $H_2SO_4(aq)$)/Konsentrasie (van swawelsuur)✓

(1)

5.3.2 • More (H_2SO_4) particles per unit volume. ✓

- More effective collisions per unit time./Higher frequency of effective collisions. ✓

- Higher reaction rate. ✓

OR

- Less (H_2SO_4) particles per unit volume. ✓

- Less effective collisions per unit time./Lower frequency of effective collisions. ✓

- Lower reaction rate ✓

- Meer (H_2SO_4) deeltjies per eenheid volume. ✓

- Meer effektiewe botsings per eenheidtyd./Hoër frekwensie van effektiewe botsings. ✓

- Hoër reaksietempo. ✓

OF

- Minder (H_2SO_4)-deeltjies per eenheid volume. ✓

- Minder effektiewe botsings per eenheidtyd./Laer frekwensie van effektiewe botsings. ✓

- Laer reaksietempo. ✓

(3)

N.
Vuf
H
K

5.3.3

Marking criteria:

- (a) Substitute $\underline{2,6(60)(40)}$ cm³ OR $\underline{(156)(40)}$ in rate formula ✓
 (b) Substitute $\underline{27\ 000}$ cm³ / $\underline{27}$ dm³ and volume in $n(H_2)$ = $\frac{V}{V_m}$ ✓
 (c) USE mole ratio $n(A\ell) = \frac{2}{3}n(H_2)$ ✓
 (d) Substitution 27 and reacting mole in $n(A\ell) = \frac{m}{M}$ ✓
 (e) Substitution of $\frac{4,05}{5}(100)$ ✓
 (f) Final answer: 83,2 % ✓
 Range: 81 – 83,3 %

Nasienkriteria:

- (a) *Vervang* $\underline{2,6(60)(40)}$ cm³ OF $\underline{(156)(40)}$ in tempo formule ✓
 (b) *Vervang* $\underline{27\ 000}$ cm³ / $\underline{27}$ dm³ en volume in $n(H_2)$ = $\frac{V}{V_m}$ ✓
 (c) *GEBRUIK* molverhouding $n(A\ell) = \frac{2}{3}n(H_2)$ ✓
 (d) *Vervang* 27 en mol gereageer in $n(A\ell) = \frac{m}{M}$ ✓
 (e) *Vervang van* $\frac{4,05}{5}(100)$ ✓
 (f) *Finale antwoord:* 81 % ✓
Gebied: 81 – 83,3 %

OPTION 1/OPSIE 1:

$$\text{Rate/Tempo} = \frac{\Delta V_{H_2}}{\Delta t}$$

$$40 = \frac{\Delta V_{H_2}}{2,6(60)} \quad \checkmark \text{ (a)}$$

$$V(H_2) = 6\ 240 \text{ cm}^3$$

$$n(H_2) = \frac{V}{V_m}$$

$$= \frac{6\ 240}{27\ 000} \quad \checkmark \text{ (b)}$$

$$= 0,23 \text{ mol}$$

$$n(A\ell) = \frac{2}{3} n(H_2)$$

$$n(A\ell) = \frac{2}{3} (0,23) \quad \checkmark \text{ (c)}$$

$$= 0,15 \text{ mol}$$

$$n(A\ell) = \frac{m}{M}$$

$$0,15 = \frac{m}{27} \quad \checkmark \text{ (d)}$$

$$m = 4,05 \text{ g}$$

$$\% \text{ purity/suiwerheid} = \frac{4,05}{5}(100)$$

$$= 81 \% \quad \checkmark \text{ (f)}$$

N.
V.
h.
M.

OPTION 2/OPSIE 2:

$$\text{rate H}_2 = 40 \text{ cm}^3 \cdot \text{s}^{-1}$$

$$\text{Rate in } n(H_2) = \frac{V}{V_m}$$

$$= \frac{40}{27\ 000} \quad \checkmark \text{ (b)}$$

$$= 0,00148 \text{ mol} \cdot \text{s}^{-1}$$

$$\text{Rate}(A\ell) = \frac{2}{3} n(H_2)$$

$$= \frac{2}{3} (0,00148) \quad \checkmark \text{ (c)}$$

$$= 9,88 \times 10^{-4} \text{ mol} \cdot \text{s}^{-1}$$

$$n(A\ell) = \frac{m}{M}$$

$$9,88 \times 10^{-4} = \frac{m}{27} \quad \checkmark \text{ (d)}$$

$$m = 0,0267 \text{ g} \cdot \text{s}^{-1}$$

$$\text{Rate/Tempo} = \frac{\Delta m_{A\ell}}{\Delta t}$$

$$0,0267 = \frac{\Delta m_{A\ell}}{2,6(60)}$$

$$m(A\ell) = 4,16 \text{ g}$$

$$\% \text{ purity/suiwerheid} = \frac{4,16}{5}(100) \quad \checkmark \text{ (e)}$$

$$= 83,2 \% \quad \checkmark \text{ (f)}$$

(6)
[18]

QUESTION 6/VRAAG 6

6.1

Marking criteria/Nasienkriteria:

If any one of the underlined key phrases in the **correct context** is omitted, deduct 1 mark./Indien enige van die onderstreepte frases in die **korrekte konteks** uitgelaat is, trek 1 punt af.

The underlined phrases must be in the correct context. / Die onderstreepte frases moet in die korrekte konteks wees.

When the equilibrium in a closed system is disturbed, the system will re-instate a new equilibrium by favouring the reaction that will cancel/oppose the disturbance. ✓✓

Wanneer die ewewig in 'n geslote sisteem versteur word, sal die sisteem 'n nuwe ewewig instel deur die reaksie te bevoordeel wat die versteuring kanselleer/teenwerk.

IF "isolated" system -1/INDIEN: "geïsoleerde" sisteem -1)

(2)

6.2

(Chemical) equilibrium/Concentrations of reactants and products remain constant./Rate of the forward and reverse reactions are equal. ✓

(Chemiese) ewewig/Konsentrasies van reaktanse en produkte bly konstant./Tempo van voorwaartse en terugwaartse reaksie is gelyk.

(1)

OPTION 1/OPSIE 1:

6.3

Exothermic/Eksotermies ✓



- With an increase in temperature the endothermic reaction is favoured. ✓
- The reverse reaction is favoured./ Equilibrium shifts to the left. / Reactants / [P₂Q] increases OR Products / [PQ₂] decreases ✓
- 'n Toename in temperatuur bevoordeel die endotermiese reaksie.
- Die terugwaartse reaksie word bevoordeel./ Ewewig skuif na links. / Reaktante / [P₂Q] neem toe OF Produkte / [PQ₂] neem af



6.5

Less than/Kleiner as ✓

OPTION 2/OPSIE 2:

Endothermic/Endotermies ✓



- With an increase in temperature the endothermic reaction is favoured. ✓
- The forward reaction is favoured./ Equilibrium shifts to the right. Reactants / [PQ₂] increases OR Products / [P₂Q] decreases ✓
- 'n Toename in temperatuur bevoordeel die endotermiese reaksie.
- Die voorwaartse reaksie word bevoordeel./ Ewewig skuif na regs./ Reaktante / [PQ₂] neem toe OF Produkte / [P₂Q] neem af




Greater than/Groter as ✓

(1)

(2)

(1)

N.
Vis
H
K



6.6

METHOD 1/METODE 1: Using lines/Gebruik lyne**CALCULATIONS USING CONCENTRATION****Marking criteria:**

- Correct K_c expression (formulae in square brackets). ✓✓
(If solid is included deduct 1 mark)
- Substitute 0,49 into K_c expression. ✓
- Substitute equilibrium concentration (0,35) into correct K_c expression. ✓
- Change in concentration/mole ✓
- USE** ratio: $P_2Q : 2PQ_2 = 1 : 2$ ✓
- Substitute 2 dm³ in $n = cV$. ✓
- Final answer = 0,85 (mol) OR 1,11 (mol) OR 3,09 (mol) ✓

Nasienkriteria:

- Korrekte K_c uitdrukking (formules in vierkantige hakies). ✓✓
(Indien vastestof invervang is, trek 1 punt af)
- Vervang 0,49 in K_c -uitdrukking. ✓
- Vervang ewewigkonsentrasie (0,35) in korrekte K_c -uitdrukking. ✓
- Verandering in konsentrasie/mol ✓
- GEBRUIK** verhouding: $P_2Q : PQ_2 = 1 : 2$ ✓
- Vervang 2 dm³ in $n = cV$. ✓
- Finale antwoord = 0,85 (mol) OF 1,11 (mol) OF 3,09 (mol) ✓

OPTION 1/OPSIE 1:

	P ₂ Q	PQ ₂
Initial concentration (mol·dm ⁻³) <i>Aanvangskonsentrasie (mol·dm⁻³)</i>	x	0
Change in concentration (mol·dm ⁻³) <i>Verandering in konsentrasie (mol·dm⁻³)</i>	0,175 ✓ (e)	0,35
Equilibrium concentration (mol·dm ⁻³) <i>Ewewigkonsentrasie (mol·dm⁻³)</i>	✓ (d) x - 0,175	0,35

$$K_c = \frac{[PQ_2]^2}{[P_2Q]} \quad \checkmark \checkmark \text{ (a)}$$

$$0,49 \checkmark \text{ (b)} = \frac{(0,35)^2}{(x - 0,175)} \checkmark \text{ (c)}$$

$$x = 0,425 \text{ mol} \cdot \text{dm}^{-3}$$


$$n(P_2Q) = cV$$

$$= 0,425 \times 2 \checkmark \text{ (f)}$$

$$= 0,85 \text{ mol} \checkmark \text{ (g)}$$

No K_c expression, correct substitution/Geen K_c -uitdrukking, korrekte substitusie: Max./Maks. 6/8

Wrong K_c expression/
Verkeerde K_c -uitdrukking: Max./Maks. 5/8



N.
Vuf
Ia
MK

OPTION2/OPSIE 2:

$$K_c = \frac{[PQ_2]^2}{[P_2Q]}$$

$$\checkmark(b) \quad 0,49 = \frac{(0,35)^2}{P_2Q}$$

$$P_2Q = 0,25 \text{ mol} \cdot \text{dm}^{-3}$$

No K_c expression, correct substitution/Geen K_c -uitdrukking, korrekte substitusie: Max./Maks. 6/8

Wrong K_c expression/
Verkeerde K_c -uitdrukking: Max./Maks. 5/8

	P_2Q	PQ_2
Initial concentration ($\text{mol} \cdot \text{dm}^{-3}$) <i>Aanvangskonsentrasie ($\text{mol} \cdot \text{dm}^{-3}$)</i>	✓ (d) 0,425	0
Change in concentration ($\text{mol} \cdot \text{dm}^{-3}$) <i>Verandering in konsentrasie ($\text{mol} \cdot \text{dm}^{-3}$)</i>	-0,175	✓ (e) 0,35
Equilibrium concentration ($\text{mol} \cdot \text{dm}^{-3}$) <i>Ewewigkonsentrasie ($\text{mol} \cdot \text{dm}^{-3}$)</i>	0,25	0,35

$$n(P_2Q) = cV$$

$$= 0,425(2) \quad \checkmark(f)$$

$$= 0,85 \text{ mol} \quad \checkmark(g)$$

CALCULATIONS USING NUMBER OF MOLES**OPTION 3/OPSIE 3:**

	P_2Q	PQ_2
Initial quantity (mol) <i>Aanvangshoeveelheid (mol)</i>	x	0
Change (mol) <i>Verandering (mol)</i>	✓ (e) 0,35	0,7
Quantity at equilibrium (mol) <i>Hoeveelheid by ewewig (mol)</i>	✓ (d) $x - 0,35$	0,7
Equilibrium concentration ($\text{mol} \cdot \text{dm}^{-3}$) <i>Ewewigkonsentrasie ($\text{mol} \cdot \text{dm}^{-3}$)</i>	✓ (f) $\frac{x - 0,35}{2}$	0,35

$$K_c = \frac{[PQ_2]^2}{[P_2Q]} \checkmark(a)$$


$$\checkmark(b) \quad 0,49 = \frac{(0,35)^2}{\frac{x - 0,35}{2}} \checkmark(c)$$

$$x = 0,85 \text{ mol} \quad \checkmark(g)$$

No K_c expression, correct substitution/Geen K_c -uitdrukking, korrekte substitusie: Max./Maks. 6/8

Wrong K_c expression/
Verkeerde K_c -uitdrukking: Max./Maks. 5/8

N.
Vis
H
K



OPTION 4/OPSIE 4:

$$K_c = \frac{[PQ_2]^2}{[P_2Q]} \quad \checkmark \checkmark \text{ (a)}$$

$$0,49 \checkmark \text{ (b)} = \frac{(0,35)^2}{[P_2Q]} \quad \checkmark \text{ (c)}$$

$$[P_2Q] = 0,25 \text{ mol} \cdot \text{dm}^{-3}$$

Wrong K_c expression/
Verkeerde K_c -uitdrukking: Max./Maks. 5/8

No K_c expression, correct substitution/Geen K_c -
uitdrukking, korrekte substitusie: Max./Maks. 6/8

	P_2Q	PQ_2
Initial quantity (mol) Aanvangshoeveelheid (mol)	$\checkmark \text{ (g)}$ 0,85	0
Change (mol) Verandering (mol)	$\checkmark \text{ (e)}$ -0,35	0,7 $\checkmark \text{ (d)}$
Quantity at equilibrium (mol) Hoeveelheid by ewewig (mol)	0,5	$\checkmark \text{ (f)}$ 0,7
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	0,25	0,35

METHOD 2/METODE 2: Using labels/Gebruik byskrifte**OPTION 1/OPSIE 1:**

	P_2Q	PQ_2
Initial concentration (mol·dm ⁻³) Aanvangskonsentrasie (mol·dm ⁻³)	x	0
Change in concentration (mol·dm ⁻³) Verandering in konsentrasie (mol·dm ⁻³)	$\checkmark \text{ (e)}$ -0,207	0,414 $\checkmark \text{ (d)}$
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	0,35	0,414

$$K_c = \frac{[PQ_2]^2}{[P_2Q]} \quad \checkmark \checkmark \text{ (a)}$$

$$(b) \checkmark \frac{0,49}{(0,35)} = \frac{[PQ_2]^2}{(0,35)} \quad \checkmark \text{ (c)}$$

$$[PQ_2] = 0,414 \text{ mol} \cdot \text{dm}^{-3}$$


No K_c expression, correct
substitution/Geen K_c - uitdrukking,
korrekte substitusie: Max./Maks. 6/8

$$\text{initial } n(P_2Q) = (0,35 + 0,207)(2) \quad \checkmark \text{ (f)}$$

$$= 1,11 \text{ mol} \quad \checkmark \text{ (g)}$$

Wrong K_c expression/
Verkeerde K_c -uitdrukking: Max./Maks. 5/8

N.
Vis
H
MK



Copyright reserved/Kopiereg voorbehou Please turn over/Blaai om asseblief



SA EXAM
PAPERS

OPTION 2/OPSIE 2:

	P ₂ Q	PQ ₂
Initial quantity (mol) Aanvangshoeveelheid (mol)	x	0
Change (mol) Verandering (mol)	0,414	0,828 ✓ (e)
Quantity at equilibrium (mol) Hoeveelheid by ewewig (mol)	0,7	0,828
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	0,35 ✓ (f)	0,414

$$K_c = \frac{[PQ_2]^2}{[P_2Q]} \quad \checkmark \checkmark \text{ (a)}$$

$$(b) \checkmark \quad 0,49 = \frac{[PQ_2]^2}{(0,35)} \quad \checkmark \text{ (c)}$$


$$[PQ_2] = 0,414 \text{ mol} \cdot \text{dm}^{-3}$$

$$x - 0,414 = 0,7 \quad \checkmark \text{ (d)}$$

$$x = 1,11 \text{ mol P}_2\text{Q} \quad \checkmark \text{ (g)}$$

No K_c expression, correct substitution/Geen K_c-uitdrukking, korrekte substitusie:
Max./Maks. 6/8

Wrong K_c expression/
Verkeerde K_c-uitdrukking: Max./Maks. 5/8

METHOD 3/METODE 3: (Equation written as reverse/Vergelyking omgekeerd geskryf)

	PQ ₂	P ₂ Q
Initial quantity (mol) Aanvangshoeveelheid (mol)	x	0
Change (mol) Verandering (mol)	1,4	0,7
Quantity at equilibrium (mol) Hoeveelheid by ewewig (mol)	(d) ✓	x - 1,4
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	$\frac{x - 1,4}{2}$	0,35

$$K_c = \frac{[P_2Q]}{[PQ_2]^2} \quad \checkmark \checkmark \text{ (a)}$$

$$(b) \checkmark \quad 0,49 = \frac{(0,35)}{[PQ_2]^2} \quad \checkmark \text{ (c)}$$

$$\frac{x - 1,4}{2} = 0,845 \quad \checkmark \text{ (f)}$$

$$x = 3,09 \text{ mol P}_2\text{Q} \quad \checkmark \text{ (g)}$$

No K_c expression, correct substitution/Geen K_c-uitdrukking, korrekte substitusie:
Max./Maks. 6/8

Wrong K_c expression/
Verkeerde K_c-uitdrukking: Max./Maks. 5/8

METHOD 4/METODE 4: Reading from graph/Aflees van grafiek**OPTION 1/OPSIE 1**

$$K_c = \frac{[PQ_2]^2}{[P_2Q]} = 0,49 \quad \text{Initial } [P_2Q] = 0$$

$$n = 0 \text{ (mol)} \quad \checkmark \checkmark \checkmark \checkmark \checkmark \checkmark \checkmark \quad (8/8)$$

N.
Vuf
lk
K

OPTION 2/OPSIE 2:

	P ₂ Q	PQ ₂
Initial concentration (mol·dm ⁻³) <i>Aanvangskonsentrasie (mol·dm⁻³)</i>	→ 0	y
Change in concentration (mol·dm ⁻³) <i>Verandering in konsentrasie (mol·dm⁻³)</i>	-0,207	0,414
Equilibrium concentration (mol·dm ⁻³) <i>Ewewigskonsentrasie (mol·dm⁻³)</i>	0,35	0,414

n = 0 (mol) ✓✓✓✓✓✓✓✓ (8/8)

(8)

- 6.7 Pressure was decreased/volume of the container was increased. ✓
Druk is verlaag/volume van die houer is vergroot.

(1)

OPTION 1/OPSIE 1: Using labels/Gebruik byskrifte


- Favours the reaction that increases the number of moles (of gas) ✓/
Bevoordeel die reaksie wat aantal mol (gas) laat toeneem
- [P₂Q] increased/neem toe ✓ OR/OF [PQ₂] decreased/neem af

OPTION 2/OPSIE 2: Using lines/Gebruik lyne

- Favours the reaction that increases the number of moles (of gas) ✓/
Bevoordeel die reaksie wat aantal mol (gas) laat toeneem
- [PQ₂] increased/neem toe ✓ OR/OF [P₂Q] decreased/neem af

(2)
[18]

N.
Vid
Lj
K






QUESTION 7/VRAAG 7

<p>Marking criteria:</p> <ul style="list-style-type: none"> Any formula $c = \frac{m}{MV}$ or $n = \frac{m}{M}$ or $c = \frac{n}{V}$ ✓ Substitute <u>10, 106 and 0,7</u> into formula ✓ Final answer: $0,13 \text{ mol} \cdot \text{dm}^{-3}$ ✓ 	<p>Nasienkriteria:</p> <ul style="list-style-type: none"> Enige formule $c = \frac{m}{MV}$ of $n = \frac{m}{M}$ of $c = \frac{n}{V}$ ✓ Vervang <u>10, 106 and 0,7</u> in formula ✓ Finale antwoord: $0,13 \text{ mol} \cdot \text{dm}^{-3}$ ✓
<p>OPTION 1/OPSIE 1:</p> $\begin{aligned} c &= \frac{m}{MV} \quad \checkmark \\ &= \frac{10}{(106)(0,7)} \quad \checkmark \\ &= 0,13 \text{ mol} \cdot \text{dm}^{-3} \quad \checkmark \end{aligned}$	<p>OPTION 2/OPSIE 2:</p> $\begin{aligned} n &= \frac{m}{M} \quad \text{Any one/Enige een} \quad \checkmark \\ &= \frac{10}{106} \\ &= 0,09 \\ c &= \frac{n}{V} \\ &= \frac{0,09}{0,7} \\ &= 0,13 \text{ mol} \cdot \text{dm}^{-3} \quad \checkmark \end{aligned}$

(3)

7.1.2 Greater than/Groter as ✓

(1)

**OR/OF****OR/OF****OR/OF****Marking criteria/Nasienkriteria:**

- Reactants ✓ Products ✓
Reaktanse ✓ Produkte ✓
- Ignore/Ignoreer → and phases/en fases
- Marking rule 6.3.10/Nasienreël 6.3.10

(2)

7.1.4 P ✓

(Titration of) weak base and a strong acid./The equivalence point is lower than pH 7. ✓

(Titratie van) 'n swak basis en 'n sterk suur./Die ekwivalente punt is laer as 'n pH van 7.

(2)


7.2

7.2.1 Dilute acid contains small amount/number of moles of acid in proportion to the volume of water. ✓✓ (2 or/of 0)

Verdunde sure bevat 'n klein hoeveelheid/getal mol suur in verhouding met die volume water.

(2)

N.
Vis
Ib
MK



Copyright reserved/Kopiereg voorbehou

Please turn over/Blaai om asseblief

7.2.2

Marking criteria:

- (a) USE of ratio:
 $n(\text{KOH})_{\text{reacted}} = 2n(\text{H}_2\text{SO}_4)_{\text{reacted}}$
 $[\text{KOH}]_{\text{reacted}} = 2n[\text{H}_2\text{SO}_4]_{\text{reacted}} \checkmark$
- (b) Subtract: $n(\text{KOH})_{\text{initial}} - n(\text{KOH})_{\text{reacted}} / [\text{KOH}]_{\text{initial}} - [\text{KOH}]_{\text{reacted}} \checkmark \checkmark$
- (c) Divide n by 0,20 dm³ in c = $\frac{n}{V} \checkmark$
- (d) Either formulae: $pH = -\log[\text{H}_3\text{O}^+] / pH = -\log[\text{H}^+] / pOH = -\log[\text{OH}^-]$ AND $[\text{H}_3\text{O}^+][\text{OH}^-] = 10^{-14} / pH + pOH = 14 \checkmark$
- (e) Substitute calculated $[\text{OH}^-]$ in $[\text{H}_3\text{O}^+][\text{OH}^-] / \text{in } pOH = -\log[\text{OH}^-] \checkmark$
- (f) Substitute calculated $[\text{H}_3\text{O}^+]$ in pH formula/ pOH in pH + $pOH = 14 \checkmark$
- (g) Final answer: 12,3 \checkmark

Nasienkriteria:

- (a) GEBRUIK verhouding:
 $n(\text{KOH})_{\text{gereageer}} = 2n(\text{H}_2\text{SO}_4)_{\text{gereageer}}$
 $[\text{KOH}]_{\text{gereageer}} = 2n[\text{H}_2\text{SO}_4]_{\text{gereageer}} \checkmark$
- (b) Aftrek: $n(\text{KOH})_{\text{aanvanklik}} - n(\text{KOH})_{\text{gereageer}} / [\text{KOH}]_{\text{aanvanklik}} - [\text{KOH}]_{\text{gereageer}} \checkmark \checkmark$
- (c) Deel n deur 0,20 dm³ in c = $\frac{n}{V} \checkmark$
- (d) Enige een v formules: $pH = -\log[\text{H}_3\text{O}^+] / pH = -\log[\text{H}^+] / pOH = -\log[\text{OH}^-]$ EN $[\text{H}_3\text{O}^+][\text{OH}^-] = 10^{-14} / pH + pOH = 14 \checkmark$
- (e) Vervang berekende $[\text{OH}^-]$ in $[\text{H}_3\text{O}^+][\text{OH}^-] / \text{in } pOH = -\log[\text{OH}^-] \checkmark$
- (f) Vervang berekende $[\text{H}_3\text{O}^+]$ in pH formule/ pOH in pH + $pOH = 14 \checkmark$
- (g) Finale antwoord: 12,3 \checkmark

OPTION 1/OPSIE 1:

$$\begin{aligned} n(\text{KOH})_{\text{reacted}} &= 2n(\text{H}_2\text{SO}_4)_{\text{reacted}} \\ &= 2(0,01) \checkmark(a) \\ &= 0,02 \\ n(\text{KOH})_{\text{excess}} &= 0,024 - 0,02 \checkmark \checkmark(b) \\ &= 0,004 \text{ mol} \\ [\text{OH}^-] &= \frac{n}{V} \\ &= \frac{0,004}{0,20} \checkmark(c) \\ &= 0,02 \text{ mol} \cdot \text{dm}^{-3} \\ [H_3O^+][OH^-] &= 10^{-14} \checkmark(d) \\ [H_3O^+] (0,02) &= 1 \times 10^{-14} \checkmark(e) \\ [H_3O^+] &= 5 \times 10^{-13} \text{ mol} \cdot \text{dm}^{-3} \\ pH &= -\log[H_3O^+] \\ &= -\log(5 \times 10^{-13}) \checkmark(f) \\ &= 12,3 \checkmark(g) \end{aligned}$$

OPTION 2/OPSIE 2:

$$\begin{aligned} [\text{KOH}] &= \frac{n}{V} \\ &= \frac{0,024}{0,20} \checkmark(c) \\ &= 0,12 \text{ mol} \cdot \text{dm}^{-3} \\ [\text{H}_2\text{SO}_4] &= \frac{n}{V} \\ &= \frac{0,01}{0,20} \\ &= 0,05 \text{ mol} \cdot \text{dm}^{-3} \\ [\text{KOH}]_{\text{reacted}} &= 2[\text{H}_2\text{SO}_4]_{\text{reacted}} \checkmark(a) \\ &= 2(0,05) \\ &= 0,1 \text{ mol} \cdot \text{dm}^{-3} \\ [\text{KOH}]_{\text{excess}} &= 0,12 - 0,1 \checkmark \checkmark(b) \\ &= 0,02 \text{ mol} \cdot \text{dm}^{-3} \\ [H_3O^+][OH^-] &= 10^{-14} \checkmark(d) \\ [H_3O^+] (0,02) &= 1 \times 10^{-14} \checkmark(e) \\ [H_3O^+] &= 5 \times 10^{-13} \text{ mol} \cdot \text{dm}^{-3} \\ pH &= -\log[H_3O^+] \\ &= -\log(5 \times 10^{-13}) \checkmark(f) \\ &= 12,3 \checkmark(g) \end{aligned}$$

OPTION 3/OPSIE 3

$$\begin{aligned} pOH &= -\log[\text{OH}^-] \\ pOH &= -\log(0,02) \checkmark(e) \\ pOH &= 1,7 \\ pH + pOH &= 14 \\ pH + 1,7 &= 14 \checkmark(f) \\ pH &= 12,3 \checkmark(g) \end{aligned}$$

N.
Vuf
Iq
MK

Any one/Enige een $\checkmark(d)$ (8)
[18]

QUESTION 8/VRAAG 8

- 8.1 Aluminium/Aℓ ✓ (1)
- 8.2 0,325 (mol·dm⁻³) ✓✓
Range/Gebied: 0,32 – 0,33 (mol·dm⁻³) (2)
- 8.3 Decreases / Neem af✓
M²⁺ is reduced/ M²⁺ used up/M²⁺ is the oxidising agent. ✓
M²⁺ word gereduseer/ M²⁺ opgebruik/M²⁺ is die oksideermiddel. (2)
- 8.4 M ✓ (1)

OPTION 1/OPTION 1 $E_{\text{cell}}^{\theta} = E_{\text{reduction}}^{\theta} - E_{\text{oxidation}}^{\theta}$ ✓ $2 \checkmark \checkmark = E_{\text{cathode}}^{\theta} - (-1,66) \checkmark$ $E_{\text{cathode}}^{\theta} = 0,34 \text{ (V)} \checkmark$ M is copper/Cu/koper ✓	NOTE/LET WEL <ul style="list-style-type: none"> Accept any other correct formula from the data sheet. /Aanvaar enige ander korrekte formule vanaf gegewensblad. Any other formula using unconventional abbreviations, e.g. $E_{\text{cell}}^{\theta} = E_{\text{OA}}^{\theta} - E_{\text{RA}}^{\theta}$ followed by correct substitutions:/Enige ander formule wat onkonvensionele afkortings gebruik, bv. $E_{\text{sel}}^{\theta} = E_{\text{OM}}^{\theta} - E_{\text{RM}}^{\theta}$ gevvolg deur korrekte vervangings 5/6
OPTION 2/OPSIE 2 $\begin{cases} \text{M}^{2+}(\text{aq}) + 2e^- \rightarrow \text{M}(\text{aq}) \\ \text{Al}^{\text{s}} \rightarrow \text{Al}^{3+}(\text{aq}) + 3e^- \end{cases} \quad E = +x \text{ V}$ $2\text{Al}^{\text{s}} + 3\text{M}^{2+}(\text{aq}) \rightarrow 2\text{Al}^{3+}(\text{aq}) + 3\text{M}(\text{s}) \quad E = 2,00 \text{ (V)} \checkmark \checkmark$ $x = 0,34 \text{ (V)} \checkmark$ M is copper/Cu/koper ✓	

- 8.6.1 Magnesium/Mg ✓ (1)
- 8.6.2 Al³⁺ is a stronger oxidising agent than Mg²⁺✓, therefore, Mg will be oxidised ✓ (to Mg²⁺)./
Mg²⁺ is a weaker oxidising agent than Al³⁺✓, therefore, Mg will be oxidised ✓ (to Mg²⁺).


Al³⁺ is 'n sterker oksideermiddel as Mg²⁺, daarom sal Mg geoksideer word (tot Mg²⁺)./
Mg²⁺ is 'n swakker oksideermiddel as Al³⁺, daarom sal Mg geoksideer word (tot Mg²⁺).

ACCEPT/AANVAAR:

Mg ion and Al ion/Mg ion en Al ion

(2)
[15]



N.
Visf
H
K



QUESTION 9/VRAAG 9

9.1 Electrical to chemical (energy)/Elektriese na chemiese (energie) ✓ (1)

9.2 P ✓ (1)

9.3 $\text{Ag}^+ + \text{e}^- \rightarrow \text{Ag}$ ✓✓**Marking criteria/Nasienkriteria:**

Ignore if charge omitted on electron./Ignoreer indien lading weggelaat op elektron.

(2)

9.4

Marking criteria:(a) Substitute 3,25 and 108 in the formula $n = \frac{m}{M}$ ✓(b) Substitute $6,02 \times 10^{23}$ in $n(\text{e}^-) = \frac{N}{N_A}$ ✓(c) Substitute 0,03 mol in $n(\text{e}^-) = \frac{N}{N_A}$ ✓(Substitute 96 500 in formula $Q = nF$)

(d) Substitute 30(60) OR 1 800 ✓

(e) Final answer: 1,61 A ✓

Nasienkriteria:(a) Vervang 3,25 en 108 in die formule $n = \frac{m}{M}$ ✓(b) Vervang $6,02 \times 10^{23}$ in $n(\text{e}^-) = \frac{N}{N_A}$ ✓(c) Vervang 0,03 mol in $n(\text{e}^-) = \frac{N}{N_A}$ ✓(Vervang 96 500 in formule $Q = nF$)

(d) Vervang 30(60) OF 1 800 ✓

(e) Finale antwoord: 1,61 A ✓

OPTION 1/OPSIE 1:

$$\begin{aligned} n(\text{Ag}) &= \frac{m}{M} \\ &= \frac{3,25}{108} \quad \checkmark(a) \\ &= 0,03 \text{ mol} \end{aligned}$$

$$\begin{aligned} n(\text{e}^-) &= \frac{N}{N_A} \\ (c) \checkmark \quad 0,03 &= \frac{N}{6,02 \times 10^{23}} \quad \checkmark(b) \\ N \text{ e}^- &= 1,81 \times 10^{22} \end{aligned}$$

$$\begin{aligned} N \text{ e}^- &= \frac{Q}{e} \text{ OF/OR } \frac{Q}{q_e} \\ 1,81 \times 10^{22} &= \frac{Q}{1,6 \times 10^{-19}} \\ Q &= 2889,6 \text{ C} \end{aligned}$$

$$\begin{aligned} I &= \frac{Q}{\Delta t} \\ &= \frac{2889,6}{30(60)} \quad \checkmark(d) \\ &= 1,61 \text{ A} \quad \checkmark(e) \end{aligned}$$


OPTION 2/OPSIE 2:

$$\begin{aligned} n(\text{Ag}) &= \frac{m}{M} \\ &= \frac{3,25}{108} \quad \checkmark(a) \\ &= 0,03 \text{ mol} = n \text{ e}^- \end{aligned}$$

$$\begin{aligned} \checkmark(b) \quad Q &= 0,03 \times 96 500 \quad \checkmark(c) \\ &= 2895 \text{ C} \\ I &= \frac{Q}{\Delta t} \\ &= \frac{2895}{30(60)} \quad \checkmark(d) \\ &= 1,61 \text{ A} \quad \checkmark(e) \end{aligned}$$

N.
Vuf
H
K

Copyright reserved/Kopiereg voorbehou

**TOTAL/TOTAAL:**(5)
[9]

150