

SA's Leading Past Year

Exam Paper Portal

S T U D Y

You have Downloaded, yet Another Great
Resource to assist you with your Studies ☺

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexamapers.co.za



SA EXAM
PAPERS

SA EXAM PAPERS
Proudly South African





basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

NATIONAL SENIOR CERTIFICATE *NASIONALE SENIOR SERTIFIKAAT*

GRADE 12/GRAAD 12

TECHNICAL MATHEMATICS P1/TEGNIESE WISKUNDE VI

NOVEMBER 2024

MARKING GUIDELINES/NASIENRIGLYNE

MARKS/PUNTE: 150

MARKING CODES/NASIENKODES	
A	Accuracy/ <i>Akkuraatheid</i>
CA	Consistent accuracy/ <i>Volgehoue akkuraatheid</i>
M	Method/ <i>Metode</i>
R	Rounding/ <i>Afronding</i>
NPR	No penalty for rounding/ <i>Geen penalisering vir afronding nie</i>
NPU	No penalty for units omitted/ <i>Geen penalisering vir eenhede weggelaat nie</i>
S	Simplification/ <i>Vereenvoudiging</i>
SF	Substitution in correct formula/ <i>Vervanging in korrekte formule</i>

These marking guidelines consist of 19 pages.
Hierdie nasienriglyne bestaan uit 19 bladsye.



SA EXAM PAPERS

NOTE:

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- Consistent accuracy (CA) applies in all aspects of the marking guidelines where indicated.
- No penalty for rounding (NPR) for ALL questions.
- # Shows questions where a Tolerance Range will be applied:
Q 2.2 ; Q 4.1.5 ; Q 5.4 & Q 9.2.2

LET WEL:

- *Indien 'n kandidaat 'n vraag TWEE keer beantwoord, sien slegs die EERSTE poging na.*
- *Volgehoue akkuraatheid (CA) is deurgaans op alle aspekte van die nasienriglyne van toepassing soos aangedui.*
- *Geen penalisering vir afronding (NPR) vir ALLE vrae nie.*
- *# Toon vrae waar Toleransie wydte (Verdraagsaamheids-omvang) toegepas word: V 2.2 ; V 4.1.5 ; V 5.4 & V 9.2.2*

QUESTION/VRAAG 1

1.1.1	$x(2x + 7) = 0$ $x = 0 \text{ or } of -\frac{7}{2}$ OR/OF $x = -3,5$	$\checkmark 0$ $\checkmark -\frac{7}{2}$ OR/OF $-3,5$ A (2)
1.1.2	$3x^2 + x = 6 + 5x$ $3x^2 - 4x - 6 = 0$ OR/OF $0 = -3x^2 + 4x + 6$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $= \frac{-(-4) \pm \sqrt{(-4)^2 - 4(3)(-6)}}{2(3)}$ OR/OF $= \frac{-(4) \pm \sqrt{(4)^2 - 4(-3)(6)}}{2(-3)}$ $= \frac{4 \pm \sqrt{88}}{6}$ $\therefore x \approx 2,23 \text{ or } of x \approx -0,90$ Refer to the Addendum/ Verwys na die Addendum	\checkmark std form/vorm A \checkmark SF CA $\checkmark x \approx 2,23$ CA $\checkmark x \approx -0,90$ CA NPR (4)



<p>1.1.3 $x^2 + 3x - 10 \leq 0$</p> $(x - 2)(x + 5) \leq 0 \quad \text{OR/OF} \quad \frac{-(3) \pm \sqrt{(3)^2 - 4(1)(-10)}}{2(1)}$ <p>Critical values/Kritieke waardes: 2 and/en -5</p> $\therefore -5 \leq x \leq 2 \quad \text{OR/OF} \quad x \in [-5; 2] \quad \text{OR/OF} \quad x \geq -5 \text{ and/en } x \leq 2$ <p>OR/OF</p>  <p>OR/OF</p> <p>Accept: From -5 to 2 / Aanvaar: Vanaf -5 tot 2</p>	<p>✓ factors/faktore/SF A</p> <p>✓ both critical values/kritieke waardes CA</p> <p>✓ correct notation/korrekte notasie/correct graphical solution/korrekte grafiese oplossing A</p> <p>AO: Full Marks/Volpunte (3)</p>
---	---



1.2	$y - x = 2$ and/en $x^2 + y^2 = 20$ $x = y - 2$ $(y - 2)^2 + y^2 = 20$ $y^2 - 4y + 4 + y^2 = 20$ $2y^2 - 4y - 16 = 0$ $2(y - 4)(y + 2) = 0$ OR/OF $y = \frac{-(4) \pm \sqrt{(-4)^2 - 4(2)(-16)}}{2(2)}$ $\therefore y = 4$ or/of $y = -2$ $\therefore x = 4 - 2 = 2$ or/of $x = -2 - 2 = -4$	✓ subject/onderwerp ✓ subst./vervang ✓ std form/vorm ✓ Factors/Faktore/SF ✓ both y-values/beide y-wrdes ✓ both x-values/beide x-wrdes	A CA CA CA CA
	OR/OF	OR/OF	
	$y = x + 2$ $x^2 + (x + 2)^2 = 20$ $x^2 + x^2 + 4x + 4 = 20$ $2x^2 + 4x - 16 = 0$ $2(x - 2)(x + 4) = 0$ OR/OF $x = \frac{-(4) \pm \sqrt{(4)^2 - 4(2)(-16)}}{2(2)}$ $\therefore x = 2$ or/of $x = -4$ $\therefore y = 2 + 2 = 4$ or/of $y = -4 + 2 = -2$	✓ subject/onderwerp ✓ subst./vervang ✓ std form/vorm ✓ Factors/Faktore/SF ✓ both x-values/beide x-wrdes ✓ both y-values/beide y-wrdes	A CA CA CA CA CA
	Refer to the Addendum/ Verwys na die Addendum		(6)



<p>1.3.1</p> $CR = \frac{CV + SV}{SV}$ $SV \times CR = CV + SV$ $CV = SV \times CR - SV \quad \text{OR/OF} \quad CV = SV \times (CR - 1)$ <p style="text-align: center;">OR/OF</p> $CR = \frac{CV}{SV} + \frac{SV}{SV}$ $CR = \frac{CV}{SV} + 1$ $CV = SV \times (CR - 1)$	<p>✓ multiplying by SV/ <i>vermenigvuldiging met SV</i></p> <p>A</p> <p>✓ CV subject/<i>onderwerp</i></p> <p>CA</p> <p style="text-align: center;">OR/OF</p> <p>✓ dividing/<i>verdeling</i></p> <p>A</p> <p>✓ CV subject/<i>onderwerp</i></p> <p>CA</p> <p style="text-align: right;">(2)</p>
<p>1.3.2</p> $CV = SV \times CR - SV$ $= 48 \text{ cm}^3 \times \frac{9,5}{1} - 48 \text{ cm}^3$ $= 408 \text{ cm}^3$ <p style="text-align: center;">OR/OF</p> $CV = SV (CR - 1)$ $= 48 \text{ cm}^3 \left(\frac{9,5}{1} - 1 \right)$ $= 408 \text{ cm}^3$ <p style="text-align: center;">OR/OF</p> $CR = \frac{CV + SV}{SV}$ $\frac{9,5}{1} = \frac{CV + 48 \text{ cm}^3}{48 \text{ cm}^3}$ $\therefore CV = 408 \text{ cm}^3$	<p>✓ SF</p> <p>CA</p> <p>✓ S</p> <p>CA</p> <p style="text-align: center;">OR/OF</p> <p>✓ SF</p> <p>CA</p> <p>✓ S</p> <p>CA</p> <p style="text-align: center;">OR/OF</p> <p>✓ SF</p> <p>A</p> <p>✓ S</p> <p>CA</p> <p style="text-align: right;">(2)</p>



1.4	$1110_2 = 14$	✓ 14	A (1)
1.5	$1110_2 \times 35$ $= 14 \times 35 = 490 = 111101010_2$ OR/OF $1110_2 \times 100011_2 = 111101010_2$ Refer to the Addendum /Verwys na die Addendum	✓ 490 ✓ 111101010_2 OR/OF ✓ 100011_2 ✓ 111101010_2 AO: Full marks/Volpunte	CA CA A CA (2)
			[22]

QUESTION/VRAAG 2


2.1.1	$p = 3$	✓ 3	A (1)
2.1.2	$1 - 7p < 0$ $p > \frac{1}{7}$ OR/OF $p > 0,14$ AO: Full marks/Volpunte	✓ $\Delta < 0$ ✓ $p > \frac{1}{7}$ OR/OF $p > 0,14$ AO: Full marks/Volpunte	A CA (2)
2.2	$3(x + 1) = x^2 + t$ $x^2 - 3x - 3 + t = 0$ OR/OF $-x^2 + 3x + 3 - t = 0$ $(-3)^2 - 4(1)(-3+t) \geq 0$ OR/OF $(3)^2 - 4(-1)(3-t) \geq 0$ $9 + 12 - 4t \geq 0$ $-4t \geq -21$ $\therefore t \leq \frac{21}{4}$ OR/OF 5,25	✓ standard form / standaardvorm ✓ SF ✓ $\Delta \geq 0$ ✓ value(s) of/ waardes van t	A CA A CA (4)
			[7]



QUESTION/VRAAG 3

3.1.1	$27^{\frac{2}{3}} = 9$ OR/OF 3^2	✓ 9 OR/OF 3^2 A (1)
3.1.2	$\begin{aligned} & (1 + \sqrt{3})^2 - \sqrt{12} \\ &= 1 + 2\sqrt{3} + 3 - 2\sqrt{3} \\ &= 4 \\ &\text{Refer to the Addendum/ Verwys na die Addendum} \end{aligned}$	✓ expanding/ uitbreiding ✓ $2\sqrt{3}$ ✓ S AO : 1 mark/ punt (3)
3.1.3	$\log_p p = 1$	✓ 1 A (1)
3.1.4	$\begin{aligned} & \log_3 81 - \log_2 \sin 30^\circ - \log_5 \sqrt{5} \\ &= \log_3 3^4 - \log_2 \frac{1}{2} - \log_5 5^{\frac{1}{2}} \text{ OR/OF } \frac{\log 3^4}{\log 3} - \frac{\log 2^{-1}}{\log 2} - \frac{\log 5^{\frac{1}{2}}}{\log 5} \\ &= 4 + 1 - \frac{1}{2} \\ &= 4\frac{1}{2} \text{ OR/OF } \frac{9}{2} \text{ OR/OF } 4,5 \end{aligned}$	✓ 4 ✓ + 1 ✓ $-\frac{1}{2}$ ✓ S AO : 1 mark/ punt (4)



3.2	$5^{x+2} - 5^x = \frac{24}{5}$ $5^x \times 5^2 - 5^x = \frac{24}{5} \quad \text{OR/OF} \quad 5^x \times 25 - 5^x = \frac{24}{5}$ $5^x (5^2 - 1) = \frac{24}{5} \quad \text{OR/OF} \quad 5^x (24) = \frac{24}{5}$ $5^x = \frac{1}{5} = 5^{-1}$ <p>$\therefore x = -1$</p> <p style="text-align: center;">OR/OF</p> $5^{x+2} - 5^x = \frac{24}{5}$ $5^{x+3} - 5^{x+1} = 24$ $5^x \times 5^3 - 5^x \times 5^1 = 24 \quad \text{OR/OF} \quad 5^x \times 125 - 5^x \times 5 = 24$ $5^x (125 - 5) = 24 \quad \text{OR/OF} \quad 5^x (120) = 24$ $5^x = \frac{1}{5} = 5^{-1}$ <p>$\therefore x = -1$</p>	✓ exp.prop./eksp.einsk. ✓ S ✓ value of/waarde van x OR/OF ✓ exp.prop./eksp.einsk. ✓ S ✓ value of/waarde van x AO: Full marks/Volpunte (3)	A CA CA CA CA CA CA CA CA CA
	$z_1 = 1 - i$	✓ form/vorm	A (1)
	$\bar{z}_1 = 1 + i$	✓ conjugate/konjugeerde	CA (1)
	 <p style="text-align: center;">OR/OF</p>	✓ Real part/Reële deel ✓ Imaginary part/Imaginäre deel ✓ terminal arm or point where Re and Im intersect/terminale arm of punt waar Re en Im sny	CA CA CA
		OR/OF	





QUESTION 4/VRAAG 4

4.1.1	$y = -1$	✓ equation / vergelyking A (1)
4.1.2	$-5 \leq x \leq 5$ OR/OF $x \in [-5; 5]$ OR/OF $x \leq 5$ and / en $x \geq -5$ OR/OF Accept: From -5 to 5 / Aanvaar: Vanaf -5 tot 5	✓ critical values / kritieke waardes A ✓ notation / notasie A (2)
4.1.3	$0 = 3^x - 1$ $3^x = 1$ $3^x = 3^0$ $x = 0$ Accept/Aanvaar: $(0; 0)$	✓ subst / vervang. $y = 0$ A ✓ $x = 0$ A AO: Full marks/Volpunte (2)
4.1.4	$y = 3^0 - 1$ $y = 0$ Accept/Aanvaar: $(0; 0)$	✓ subst. / vervang $x = 0$ A ✓ $y = 0$ CA AO: Full marks/Volpunte (2)
4.1.5	<p>For/vir h :</p> <ul style="list-style-type: none"> ✓ shape / vorm A ✓ all intercepts on axes / alle afsnitte op die asse A <p>For/ vir f:</p> <ul style="list-style-type: none"> ✓ shape / vorm A ✓ asymptote / asimptoot CA ✓ intercept(s) / afsnitte CA 	(5)



4.1.6	$-5 \leq x \leq 0$ or/of $x = 5$ OR/OF $x \in [-5 ; 0]$ or/of $x = 5$ OR/OF $x \geq -5$ and/en $x \leq 0$ or/of $x = 5$ Accept: From -5 to 0 or $x = 5$ /Aanvaar: Vanaf -5 tot 0 of $x = 5$	✓ $-5 \leq x \leq 0$ ✓ $x = 5$	CA A
4.2	$g(x) = a(x - p)^2 + q$ $g(x) = a(x - 1)^2 - 4$ $4 = a(3 - 1)^2 - 4$ $8 = 4a$ $a = 2$ $g(x) = 2(x - 1)^2 - 4 = 2x^2 - 4x - 2$ Refer to the Addendum/ Verwys na die Addendum	✓ subst./ vervang (1 ; -4) ✓ subst./vervang (3 ; 4)	A A
		✓ value of / waarde van a	CA
		✓ equation in reqd form/ vergelyking in voorgesk. vorm	CA (4)
4.3.1	$x = 0$ $y = 2$	✓ vertical- asymptote / vertikale-asimptoot ✓ horizontal- asymptote / horizontale-asimptoot	A A
		AO: Full marks/Volpunte	(2)
4.3.2	$p(x) = x + 2$ $4 = k + 2$ $k = 2$	✓ Subst./vervang ✓ k value /waarde	A CA
		AO: Full marks/Volpunte	(2)
4.3.3	$x = 2$ Accept/Aanvaar: $(2; 0)$	✓ x value / waarde	CA (1)
4.3.4	$h(x) = \frac{a}{x} + q = \frac{a}{x} + 2$ $0 = \frac{a}{2} + 2$ $a = -4$ $\therefore h(x) = -\frac{4}{x} + 2$	✓ Subst./vervang (2; 0) ✓ a value/ waarde	CA CA
			(2)



QUESTION/VRAAG 5**Note: ZERO MARKS if Incorrect Formula is used****Let wel: GEEN PUNTE indien Verkeerde Formule gebruik word.**

5.1	$i_{eff} = \left(1 + \frac{i}{m}\right)^m - 1$ $1 + 9,1\% = \left(1 + \frac{i}{4}\right)^4$ $1 + \frac{i}{4} = \sqrt[4]{1,091}$ $i = 4(\sqrt[4]{1,091} - 1)$ $\approx 8,80\% \quad \text{OR/OF} \quad \approx 0,088$	✓ F ✓ SF ✓ S ✓ 8,80% OR/OF 0,088 CA CA CA (4)
5.2	$A = P(1 + i)^n$ $= 50\ 000(1 + 0,03)^5$ $\approx 57\ 963,70$ OR/OF $A_1 = 50\ 000(1,03) = 51\ 500$ $A_2 = 51\ 500(1,03) = 53\ 045$ $A_3 = 53\ 045(1,03) = 54\ 636,35$ $A_4 = 54\ 636,35(1,03) = 56\ 275,4405$ $A_5 = 56\ 275,4405(1,03) \approx 57\ 963,70$ Accept/Aanvaar: $\approx 57\ 963 \quad \text{OR/OF} \quad \approx 57\ 964$	✓ F ✓ SF ✓ S OR/OF ✓ SF ✓ 51 500 OR/OF ✓ S (3)
5.3.1	$A = 260\ 000 \times 0,25 = R65\ 000$	✓ value / waarde (1)
5.3.2	$A = P(1 - i)^n$ $65\ 000 = 260\ 000(1 - 0,14)^n$ $(0,86)^n = 0,25 \quad \text{OR/OF} \quad \left(\frac{65\ 000}{260\ 000}\right) = (1 - 0,14)^n$ $n \log 0,86 = \log 0,25$ $n = \frac{\log 0,25}{\log 0,86} \quad \text{OR/OF} \quad n = \log_{(1 - 0,14)}\left(\frac{65\ 000}{260\ 000}\right)$ $n \approx 9,19 \text{ years / jaar}$ Accept/Aanvaar: $n \approx 9 \text{ years / jaar} \quad \text{OR/OF}$ $n \approx 10 \text{ years / jaar}$ Refer to the Addendum/ Verwys na die Addendum	✓ F ✓ SF ✓ log form / vorm ✓ value of / waarde van n CA CA CA (4)

<p>5.4</p> $i = \frac{10\%}{12}$ $i = \frac{8\%}{4}$	$A = 20\ 000 \left(1 + \frac{10\%}{12}\right)^{1,5 \times 12} \approx R\ 23\ 222,25$ $\approx 23\ 222,25 \left(1 + \frac{8\%}{4}\right)^{1,5 \times 4} \approx R\ 26\ 152,03$ $\therefore 26\ 152,03 - 3\ 000$ $\approx R\ 23\ 152,03$ $\approx 23\ 152,03 \left(1 + \frac{8\%}{4}\right)^{1 \times 4}$ $\approx R\ 25\ 060,49$ <p>OR/OF</p> $A = 20\ 000 \left(1 + \frac{10\%}{12}\right)^{1,5 \times 12} \left(1 + \frac{8\%}{4}\right)^{2,5 \times 4}$ $- 3\ 000 \left(1 + \frac{8\%}{4}\right)^{1 \times 4}$ $\approx R\ 25\ 060,49$ <p>OR/OF</p> $A = \left[20\ 000 \left(1 + \frac{10\%}{12}\right)^{1,5 \times 12} \left(1 + \frac{8\%}{4}\right)^{1,5 \times 4} - 3\ 000 \right]$ $\times \left(1 + \frac{8\%}{4}\right)^{1 \times 4}$ $\approx R\ 25\ 060,49$	$\checkmark 20\ 000 \left(1 + \frac{10\%}{12}\right)^{1,5 \times 12} \quad \textbf{A}$ $\checkmark 23\ 222,25 \left(1 + \frac{8\%}{4}\right)^{1,5 \times 4} \quad \textbf{CA}$ $\checkmark \mathbf{M} \quad \textbf{A}$ $\checkmark 23\ 152,03 \left(1 + \frac{8\%}{4}\right)^{1 \times 4} \quad \textbf{CA}$ $\checkmark \approx R\ 25\ 060,49 \quad \textbf{CA}$ <p>OR/OF</p> $\checkmark 20\ 000 \left(1 + \frac{10\%}{12}\right)^{1,5 \times 12} \quad \textbf{A}$ $\checkmark \left(1 + \frac{8\%}{4}\right)^{2,5 \times 4} \quad \textbf{A}$ $\checkmark \mathbf{M} \quad \textbf{A}$ $\checkmark \left(1 + \frac{8\%}{4}\right)^{1 \times 4} \quad \textbf{A}$ $\checkmark \approx R\ 25\ 060,49 \quad \textbf{CA}$ <p>(5)</p>
		[17]



QUESTION/VRAAG 6

6.1	$\begin{aligned} f(x) &= 9x - 6 \\ f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{9(x+h) - 6 - (9x - 6)}{h} \\ &= \lim_{h \rightarrow 0} \frac{9x + 9h - 6 - 9x + 6}{h} \\ &= \lim_{h \rightarrow 0} \frac{9h}{h} \\ &= \lim_{h \rightarrow 0} (9) \\ \therefore f'(x) &= 9 \end{aligned}$	✓ definition/definisie ✓ SF ✓ S ✓ S ✓ 9 Penalty: 1 mark for incorrect notation/ Penaliseer : 1 punt vir foutiewe notasie AO: 1mark/punt (5)
6.2	$\begin{aligned} f(x) &= 11\pi^2 \\ f'(x) &= 0 \end{aligned}$	✓ 0 A (1)
6.3.1	$\begin{aligned} y &= x \left(3 + \frac{3}{x^5} \right) \\ &= 3x + 3x^{-4} \end{aligned}$	✓ $3x$ ✓ $3x^{-4}$ or/of $\frac{3}{x^4}$ (2)
6.3.2	$\frac{dy}{dx} = 3 - 12x^{-5}$	✓ 3 ✓ $-12x^{-5}$ or/of $-\frac{12}{x^5}$ (2)
6.4.1	$\sqrt[5]{x^8} = x^{\frac{8}{5}}$	✓ $x^{\frac{8}{5}}$ A (1)
6.4.2	$\begin{aligned} D_x &\left[x^{\frac{8}{5}} - 5x^{12} \right] \\ &= \frac{8}{5}x^{\frac{3}{5}} - 60x^{11} \end{aligned}$	✓ $\frac{8}{5}x^{\frac{3}{5}}$ ✓ $-60x^{11}$ A (2)



6.5.1	$g(x) = -x^3 + 6x^2$ $g'(x) = -3x^2 + 12x$	$\checkmark -3x^2$ $\checkmark 12x$ (2)
6.5.2	$g'(-2) = -3(-2)^2 + 12(-2)$ $= -36$	\checkmark Sub. into derivative / vervang in afgeleide CA $\checkmark -36$ CA AO: Full marks/Volpunte (2)
6.5.3	$-3x^2 + 12x = -36$ $-3x^2 + 12x + 36 = 0$ $-3(x-6)(x+2) = 0$ OR/OF $x = \frac{-(12) \pm \sqrt{(12)^2 - 4(-3)(36)}}{2(-3)}$ $\therefore x = 6$ $y = g(6) = -(6)^3 + 6(6)^2 = 0$ The other point is /Die ander punt is $(6 ; 0)$	\checkmark equating / gelykstelling CA \checkmark factors/formula/faktore CA \checkmark x-coordinate of other point / $x\text{-koördinaat van ander punt}$ CA \checkmark y-coordinate of other point / $y\text{-koördinaat van ander punt}$ CA (4)
		[21]



QUESTION/VRAAG 7

7.1	OD = 60 units/ eenhede	✓ length/ lengte A (1)
7.2	x -intercepts / afsnitte $y = 0$ $(x + 2)(x^2 - x - 30) = 0$ $(x + 2)(x - 6)(x + 5) = 0 \quad \text{OR/OF}$ $x + 2 = 0 \quad \text{or/of} \quad x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(1)(-30)}}{2(1)}$ $\therefore x = -2 \quad \text{or/of} \quad x = 6 \quad \text{or/of} \quad x = -5$ $\therefore A(-5; 0) \text{ and/en } C(6; 0)$	✓ quadratic factor/ kwadratiese faktor A ✓ factors/ faktore / SF CA ✓ coordinates of/koördinate van A CA ✓ coordinates of/koördinate van C CA AO: Full marks/Volpunte (4)
7.3	$f'(x) = 3x^2 + 2x - 32 = 0$ $x = \frac{-(2) \pm \sqrt{(2)^2 - 4(3)(-32)}}{2(3)}$ $\therefore x \approx -3,62 \quad \text{or/of} \quad x \approx 2,95$ $f(2,95) = (2,95)^3 + (2,95)^2 - 32(2,95) - 60 \approx -120,03$ $\therefore G(-3,62; -120,03)$	✓ derivative/afgeleide A ✓ equating derivative to 0/ stel afgeleide gelyk aan 0 A ✓ SF CA ✓ both values of/beide waardes van x CA ✓ correct value of / korrekte waarde van y CA ✓ coordinates of/koördinate van G CA NPR (6)
7.4.1	$-5 \leq x \leq -2$ OR/OF $x \in [-5; -2]$ OR/OF $x \geq -5 \text{ and/en } x \leq -2$ Accept: From -5 to -2 / Aanvaar: Vanaf -5 tot -2	✓ critical values/kritieke waardes CA ✓ correct notation/korrekte notasie A (2)
7.4.2	$-3,62 < x < 2,95$ OR/OF $x \in (-3,62; 2,95)$ OR/OF $x > -3,62 \text{ and/en } x < 2,95$ Accept: Between $-3,62$ and $2,95$ / Aanvaar: Tussen $-3,62$ en $2,95$	✓ critical values/kritieke waardes CA ✓ correct notation/korrekte notasie A NPR (2)
		[15]



QUESTION/VRAAG 8

8.1	R10 000 Refer to the Addendum/ Verwys na die Addendum	✓ R10 000 A (1)
8.2	$P'(x) = -60x^2 + 6 000$	✓ derivative/ afgeleide A (1)
8.3	For maximum / Vir maksimum: $P'(x) = 0$ $-60x^2 + 6 000 = 0$ $-60(x+10)(x-10) = 0$ OR/OF $x = \frac{-(0) \pm \sqrt{(0)^2 - 4(-60)(6 000)}}{2(-60)}$ OR/OF $x^2 = \frac{6 000}{60}$ $x = 10 \text{ or/of } x \neq -10$ $P(10) = -20(10)^3 + 6 000(10) - 10 000$ $= R30 000$	✓ equating derivative to/ stel afgeleide aan 0 A ✓ factors/faktore /SF CA ✓ correct value of x / korrekte waarde vir x CA ✓ SF CA ✓ Maximum Profit / maksimum wins CA NPU (5)
		[7]



QUESTION/VRAAG 9

9.1.1	$\int -\frac{6}{x} dx = -6 \ln x + C$	✓ $-6 \ln x$ ✓ C (2)
9.1.2	$\begin{aligned} & \int (3x - 4)(x + 2) dx \\ &= \int (3x^2 + 2x - 8) dx \\ &= x^3 + x^2 - 8x + C \end{aligned}$	✓ S ✓ x^3 ✓ x^2 ✓ $-8x + C$ (4)
9.2.1	$\int 2^x dx = \frac{2^x}{\ln 2} + C$	✓ $\frac{2^x}{\ln 2} + C$ (1)
9.2.2	$\begin{aligned} A &= \int_{-2}^0 f(x) dx = \int_{-2}^0 2^x dx \\ &= \left[\frac{2^x}{\ln 2} \right]_{-2}^0 \\ &= \left[\frac{2^0}{\ln 2} \right] - \left[\frac{2^{-2}}{\ln 2} \right] = \frac{3}{4 \ln 2} \\ &= \frac{3}{\ln 2} \quad \text{OR/OF } \approx 1,08 \text{ units}^2 / \text{eenhede}^2 \end{aligned}$	✓ Area notation using integrals/Area-notasie met gebruik van integrale ✓ SF ✓ A value / waarde $\begin{aligned} B &= \int_2^3 f(x) dx = \int_2^3 2^x dx = \left[\frac{2^x}{\ln 2} \right]_2^3 \\ &= \left[\frac{2^3}{\ln 2} \right] - \left[\frac{2^2}{\ln 2} \right] = \frac{4}{\ln 2} \\ &= \frac{4}{\ln 2} \quad \text{OR/OF } \approx 5,77 \text{ units}^2 / \text{eenhede}^2 \end{aligned}$ ✓ SF ✓ B value / waarde $\therefore \frac{B}{A} = \frac{4}{\ln 2} - \frac{4}{\ln 2} = \frac{16}{3} \quad \text{OR/OF } \therefore \frac{B}{A} = \frac{5,77}{1,08} = 5,34$ ✓ M

OR/OF

SA EXAM PAPERS



	$\begin{aligned} B - 4A &= \frac{4}{\ln 2} - 4 \left(\frac{\frac{3}{4}}{\ln 2} \right) \textbf{OR/OF} \approx 5,77 - 4(1,08) \\ &= \frac{1}{\ln 2} \qquad \qquad \qquad \approx 1,45 \end{aligned}$ <p style="text-align: center;">OR/OF</p> $\begin{aligned} B > 4A &\quad \frac{4}{\ln 2} > 4 \left(\frac{\frac{3}{4}}{\ln 2} \right) \textbf{OR/OF} \quad 5,77 > 4(1,08) \\ &\quad \frac{4}{\ln 2} > \frac{3}{\ln 2} \qquad \qquad \qquad 5,77 > 4,32 \end{aligned}$ <p style="text-align: center;">∴ The learner's claim is NOT VALID / Die leerder se bewering is NIE GELDIG NIE.</p>	
		✓Conclusion / Gevolgtrekking CA (7)
		[14]
	TOTAL/TOTAAL	[150]

