

# SA's Leading Past Year

## Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ [www.saexampapers.co.za](http://www.saexampapers.co.za)



# SA EXAM PAPERS

SA EXAM PAPERS  
Proudly South African



Province of the  
**EASTERN CAPE**  
EDUCATION

Iphondo leMpuma Kapa: Isebe leMfundo  
Provinsie van die Oos Kaap: Departement van Onderwys  
Porafense ya Kapa Botjhabelo: Lelapha la Thuto

**NATIONAL  
SENIOR CERTIFICATE/  
NACIONALE  
SENIOR SERTIFIKAAT**

**GRADE/GRAAD 12**

**JUNE/JUNIE 2025**

**PHYSICAL SCIENCES: CHEMISTRY P2/  
FISIESE WETENSKAPPE: CHEMIE V2  
MARKING GUIDELINE/NASIENRIGLYN**

**MARKS/PUNTE: 150**

---

This marking guideline consists of 15 pages./  
*Hierdie nasienriglyn bestaan uit 15 bladsye.*

---



**SA EXAM PAPERS**

Proudly South African

**QUESTION/VRAAG 1**

- |      |      |             |
|------|------|-------------|
| 1.1  | B ✓✓ | (2)         |
| 1.2  | D ✓✓ | (2)         |
| 1.3  | D ✓✓ | (2)         |
| 1.4  | B ✓✓ | (2)         |
| 1.5  | C ✓✓ | (2)         |
| 1.6  | C ✓✓ | (2)         |
| 1.7  | B ✓✓ | (2)         |
| 1.8  | D ✓✓ | (2)         |
| 1.9  | C ✓✓ | (2)         |
| 1.10 | B ✓✓ | (2)         |
|      |      | <b>[20]</b> |



## QUESTION/VRAAG 2

- 2.1 A bond or an atom or a group of atoms that determine(s) the physical and chemical properties of a group of organic compounds. ✓✓

*in Binding of 'n atoom of 'n groep atome wat die fisiese en chemiese eienskappe van 'n groep organiese verbindings bepaal.* (2)

- 2.2 2.2.1 A ✓ (1)

- 2.2.2 C ✓ (1)

- 2.2.3 B ✓ (1)

- 2.3 2.3.1 3-methylbutanal ✓✓  
3-metielbutanaal

**Marking criteria/Nasienkriteria:**

- Butanal / butanaal ✓
- Whole name correct / Hele naam korrek ✓

(2)

- 2.3.2 4-methylpent-2-yne ✓✓  
4-metieipent-2-yn

**Marking criteria/Nasienkriteria:**

- Pent-2-yne / pent-2-yn ✓
- Whole name correct / Hele naam korrek ✓✓

**OR/OF**

- 4-methyl-2-pentyne ✓✓  
4-metiel-2-pentyn

(2)


- 2.3.3 2,3-dibromo-4-ethylhexane ✓✓✓  
2,3-dibromo-4-etielheksaan

**Marking criteria/Nasienkriteria:**

- Hexane / heksaan ✓
- dibromo and ethyl / dibromo en etiel ✓
- Whole name correct / Hele naam korrek ✓

(3)

- 2.4 2.4.1

**Markingcriteria  
/Nasienkriteria**


- Correct functional group / Funksionele groep korrek ✓  
1/2
- Whole structure correct/ Hele struktuur korrek ✓ 2/2

(2)

- 2.4.2 Esterification / Condensation / Esterifikasie / Kondensasie ✓ (1)



2.4.3

**Marking criteria/Nasienkriteria**

- Correct functional group / *Funksionele groep korrek* ✓ 1/2
- Whole structure correct/ *Hele struktuur korrek* ✓ 2/2

(2)

2.5 2.5.1  $43,24 = \frac{M}{74} \times 100$  ✓

$$M = 31,9976$$

$$M = 32 \text{ g} \cdot \text{mol}^{-1}$$

$$M(\text{O}) = 16 \text{ g} \cdot \text{mol}^{-1}$$




Organic compound is either carboxylic acid or ester.

*Organiese verbinding is 'n karboksiesuur of 'n ester.*General formula / *Algemene formule*:  $\text{C}_n\text{H}_{2n}\text{O}_2$ 



$$12 \times n + 2 \times n + 32 = 74$$
 ✓

$$n = 3$$



(3)

2.5.2

**AND/EN****Marking criteria/Nasienkriteria**

- Correct functional group / *Funksionele groep korrek* ✓ 1/2
- Whole structure correct/ *Hele struktuur korrek* ✓ 2/2

(4)

**[24]**

## QUESTION/VRAAG 3

**Marking criteria/ Nasienkriteria**

If any of the underlined key words/ phrases in the **correct context** are omitted:  
- 1 mark per word/phrase.

Indien enige van die sleutelwoorde/ frases in die **korrekte konteks** weggelaat word: - 1 punt per woord/frase.

- 3.1 Boiling point is the temperature at which the vapour pressure of a substance /liquid equal the atmospheric pressure ✓✓

*Kookpunt is die temperatuur waarteen die dampdruk van 'n stof/vloeistof gelyk aan die atmosferiesedruk is*

(2)

- 3.2 Same functional group / Same homologous series ✓

*Dieselfde funksionele groep / Dieselfde homoloëreeks*

Accept same type of intermolecular forces/Aanvaar dieselfde tipe van intermolekulêre kragte

(1)

- 3.3 **Marking criteria**

- The boiling point increases down the table
- Chain length / molecular size increases down the table
- Increase in the strength of the London forces/Dispersion forces/Induced dipole forces / down the table
- Relate the strength of London forces /dispersion forces/induced dipole to energy involved

**Nasienkriteria**

- *Die kookpunt styg teen die tafel af*
- *Kettinglengte / molekulêre grootte neem teen die tabel af*
- *Toename in die sterkte van die Londonkragte /Dispersiekragte /Geïnduseerde dipoolkragte / teen die tabel af*
- *Verband hou met die sterkte van die Londonkragte/ verspreidingskrag/ geïnduseerde dipoolkragte met die energie betrokke*

- The trend shows an increase in the boiling point down the table ✓
- Increase in the surface area/chain length/molecular size down the table ✓
- Strength of London forces/dispersion forces/induced dipole forces / increases ✓
- More energy is needed to overcome intermolecular forces ✓

- *Die tendens toon 'n toename in die kookpunt teen die tabel af*
- *Toename in die oppervlakte/ketting lengte/molekulêre grootte teen die tabel af*
- *Sterkte van die Londonkragte/verspreidingskragte/geïnduseerde dipoolkragte neem toe*
- *Meer energie word benodig om die intermolekulêre kragte te oorkom*

(4)

- 3.4 Decreases. ✓ The boiling point increases. ✓

*Afneem. Die kookpunt neem toe.*

(2)



- 3.5 2-methylpropan-1-ol / 2-methyl-1-propanol ✓✓  
2-metielpropan-1-ol/ 2-metiel-1-propanol

**Marking criteria/Nasienkriteria:**

- Propan-1-ol ✓
- Whole name correct / Hele naam korrek ✓

(2)

3.6

**Marking criteria**

- Compare the boiling point of the alcohol to the alcohol with 3-carbon atom alcohol
- Compare the surface area/chain length/molecular size of the alcohol to the alcohol with 3-carbon atoms
- Compare the boiling point of the alcohol to the alcohol with 4-carbon atoms
- Compare the surface area/chain length of the alcohol to the alcohol with 4-carbon atoms

**Nasienkriteria**

- Vergelyk die kookpunt van die alkohol met die 3-koolstof atoom alkohol
- Vergelyk die oppervlakte/kettinglengte/molekulêre grootte van die alkohol met die 3-koolstof atoom alkohol
- Vergelyk die kookpunt van die alkohol met die 4-koolstof atoom alkohol
- Vergelyk die oppervlakte/kettinglengte van die alkohol met die 4-koolstof atoom alkohol

- The boiling point of the unknown alcohol (2-methylpropan-1-ol) is higher than that of the alcohol with 3-carbon atoms (propan-1-ol) ✓
- Surface area / chain length / molecular size is greater than the alcohol with 3-carbon atoms (propan-1-ol) ✓
- The boiling point of the unknown alcohol (2-methylpropan-1-ol) is smaller than that of the straight chain alcohol with 4-carbon atoms (butan-1-ol) ✓
- Surface area / chain length is smaller than the straight chain alcohol with 4-carbon atoms (butan-1-ol) ✓
- Die kookpunt van die onbekende alkohol (2-metielpropan-1-ol) is hoër as dié alkohol met 3-koolstofatome (propan-1-ol)
- Oppervlakte / kettinglengte / molekulêre grootte is groter as die alkohol met 3-koolstof atome (propan-1-ol)
- Die kookpunt van die onbekende alkohol (2-metielpropan-1-ol) is laer as die reguitketting alkohol met 4-koolstofatome (butan-1-ol)
- Oppervlakte / kettinglengte is kleiner as die reguitketting alkohol met 4-koolstof atome (butan-1-ol)

(4)  
[15]

## QUESTION/VRAAG 4

4.1 4.1.1 Breaking down of long chain hydrocarbon molecules into more useful shorter chains ✓✓ (2 or 0)  
*Die afbreek van lang kettingkoolwaterstof-molekules in meer bruikbare, molekule korter kettings (2 of 0)* (2)

4.1.2  $C_2H_4$  ✓✓ (2)

4.1.3 Used as a fuel / Word as 'n brandstof gebruik ✓ (1)


4.1.4  $C_5H_{12} + 8 O_2 \checkmark \longrightarrow 5 CO_2 + 6 H_2O \checkmark$  (✓ bal.)  
 Reactants, ✓ products ✓ and balancing ✓  
*Reaktanse, produkte en balansering* (3)

4.2 4.2.1 2-bromopentane / 2-bromopentaaan ✓✓ (2)

4.2.2 Substitution/ Hydrolysis of haloalkanes / *Substitusie / Hydrolise van haloalkane* ✓ (1)

4.2.3 Diluted strong base / diluted NaOH ✓  
Verdunde sterk basis / verdunde NaOH (1)

4.2.4




**Marking criteria/Nasienkriteria**

- Correct functional group / *Funksionele groep korrek* ✓ 1/2
- Whole structure correct/ *Hele struktuur korrek* ✓ 2/2

(2)

4.2.5 Dehydration / *Dehidrasie / Dihidratering* ✓ (1)

4.2.6



**Marking criteria/Nasienkriteria**

- Correct functional group / *Funksionele groep korrek* ✓ 1/2
- Whole structure correct/ *Hele struktuur korrek* ✓ 2/2

(2)

[17]



## QUESTION/VRAAG 5

5.1 EXOTHERMIC / EKSOTERMIES. ✓  $\Delta H < 0$  ✓ (2)

5.2 Carbon dioxide (CO<sub>2</sub>) gas escapes / Koolstof dioksied (CO<sub>2</sub>) gas ontsnap ✓✓ (2)

5.3 189 g ✓ Accept/Anvaar (188,5 – 189) (1)

5.4 5.4.1 11 (minutes / minute) accept/ aanvaar 10 (minutes / minute) ✓ (1)

5.4.2 EQUAL TO ✓

The amount of CaCO<sub>3</sub> consumed is equal to the amount of CO<sub>2</sub> produced / the mol ratio of CaCO<sub>3</sub> to CO<sub>2</sub> is 1 : 1. ✓

The time taken for consumption and production is the same. ✓

GELYK AAN

Die hoeveelheid CaCO<sub>3</sub> verbruik is gelyk aan die hoeveelheid CO<sub>2</sub> vervaardig / die molverhouding van CaCO<sub>3</sub> tot CO<sub>2</sub> is 1 : 1.

Die tyd geneem vir die verbruik en vervaardiging is dieselfde. (3)

5.4.3 **Marking criteria / Nasienkriteria**

- (a) Subtract / Aftrek 200 – 185
- (b) Subst. into / Vervanging in  $n = m/M$
- (c) Formula / Formule  $V = nV_m$
- (d) Subst. into/ Vervanging in  $V = nV_m$
- (e) Final answer/ Finale antwoord

$$n = \frac{M}{m}$$

$$n = \frac{200 - 185 \checkmark (a)}{44} \checkmark (b)$$

$$n = 0,34 \text{ mol}$$

$$V = nV_m \checkmark (c)$$

$$V = (0,34)(24000) \checkmark (d)$$

$$V = 8\,160 \text{ (cm}^3\text{)} \checkmark (e)$$

$$\text{Range} = 8160 - 8181,82 \quad (5)$$



5.4.4 **Marking criteria/Nasienkriteria**

- (a) Use of mol ratio / *Gebruik van molverhouding*  $\text{CO}_2 : \text{CaCO}_3$
- (b) Subst. into/ *Vervanging in*  $n = m/M$
- (c) Subst. into rate equation/ *Vervanging in tempo vergelyking*
- (d) Final answer / *Finale antwoord*

**Positive marking from/Positiewe nasien van 5.4.3**

$$n(\text{CO}_2) = n(\text{CaCO}_3) = 0,34 \text{ mol} \checkmark \text{ (a)}$$

$$m = nM$$

$$m = (0,34)(100) \checkmark \text{ (b)}$$

$$m = 34 \text{ g}$$

$$\frac{\text{rate/}}{\text{tempo}} = - \frac{\Delta m}{\Delta t}$$

$$\frac{\text{rate/}}{\text{tempo}} = - \frac{0 - 34}{11} \checkmark \text{ (c)}$$

$$m = 3,09 \text{ g} \cdot \text{min}^{-1} \checkmark \text{ (d)} \quad (4)$$

5.5 5.5.1 **Marking criteria for hypothesis**

- Statement regarding the correct independent and dependent variable
- Statement is measurable

**Nasienkriteria vir hipotese**

- *Stelling om die onafhanklike en afhanklike veranderlike is korrek*
- *Stelling is meetbaar*

A(n) decrease / increase in temperature will decrease / increase the reaction rate.

*'n Afname/ toename in temperatuur sal die reaksietempo laat afneem/ toeneem.* (2)

5.5.2 INCREASES / TOENEEM  $\checkmark$  (1)

- 5.5.3
- Higher temperature means the average kinetic energy of the particles increases  $\checkmark$
  - More particles will have sufficient kinetic energy / more particles will have kinetic energy equal to or higher than the activation energy  $\checkmark$
  - More effective collisions per unit time / Frequency of the effective collisions increases  $\checkmark$



- Hoër temperatuur beteken die gemiddelde kinetiese energie van die deeltjies neem toe
- Meer deeltjies het genoeg kinetiese energie / meer deeltjies het kinetiese energie gelyk of hoër as die aktiveringsenergie
- Meer effektiewe botsings per tydseenheid / Frekwensie van die effektiewe botsings neem toe

(3)  
[24]

## QUESTION/VRAAG 6

6.1.1

**Marking criteria/Nasienkriteria**

If any of the underlined key words/phrases in the **correct context** are omitted:  
- 1 mark per word/phrase.

Indien enige van die sleutelwoorde/frases in die **korrekte konteks** weggelaat word: - 1 punt per woord/frase.

When the equilibrium in a closed system is disturbed, the system will re-instate a new equilibrium by favouring the reaction that will oppose/cancel the disturbance. ✓✓

Wanneer die ewewig in 'n geslote sisteem versteur word, sal die sisteem 'n nuwe ewewig deur die reaksie wat die versteuring teenwerk/kanseleer, te bevoordeel. (2)

6.1.2

(a) NO EFFECT / GEEN EFEEK ✓ (1)

(b) INCREASES / TOENEEM ✓ (1)


(c) DECREASES / AFNEEM ✓

- An increase in pressure favours the reaction that produces the least number of gaseous moles. ✓
- Mole ratio of reactants : products 1 : 2 ✓
- The reverse reaction is favoured / the equilibrium position shifted towards left. ✓
- 'n Toename in druk bevoordeel die reaksie wat die minste gasmol produseer.
- Die molverhouding reaktanse : produkte 1 : 2
- Die terugwaartse reaksie word bevoordeel / die ewewigsposisie verskuif links. (4)

6.2

6.2.1  $2\text{NO}_2(\text{g}) \rightarrow \text{N}_2\text{O}_4(\text{g})$  ✓✓ (2)

6.2.2

**Marking criteria/****Nasienkriteria**

Both graphs rates increases vertically and reached new equilibrium ✓

Increase in the reverse reaction rate is higher than the forward rate ✓

*Beide grafieke se tempo neem vertikaal toe en bereik 'n nuwe ewewig*

*Toename in die tempo van die terugwaartse reaksie is hoër as die tempo van die voortwaartse reaksie*

(2)

$$6.3 \quad 6.3.1 \quad K_c = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]} \quad \checkmark$$

$$K_c = \frac{(0,0457)^2}{(0,448)} \quad \checkmark$$

$$K_c = 4,66 \times 10^{-3} \quad \checkmark \quad (3)$$

6.3.2 LOW YIELD. ✓  $K_c$  is less than 1. ✓  
LAE OPBRENGS.  $K_c$  is kleiner as 1.

(2)

6.3.3 **POSITIVE MARKING FROM / POSITIEWE NASIEN VANAF 6.3.1**  
**MARKING CRITERIA**

- Subst.  $K_c$  value and correct  $\text{N}_2\text{O}_4$  concentration into the correct  $K_c$  expression ✓
- Determine the change in concentration of  $\text{NO}_2$  ✓
- Correct ratio  $\text{N}_2\text{O}_4$  and  $\text{NO}_2$  ✓
- Determine the initial concentration  $\text{N}_2\text{O}_4$  ✓
- Subst. change in concentration and initial concentration of  $\text{N}_2\text{O}_4$  in % change formula ✓
- Final answer ✓

**NASIENKRITERIA**

- Vervang die  $K_c$ -waarde en korrekte  $\text{N}_2\text{O}_4$  konsentrasie in die korrekte  $K_c$ -uitdrukking
- Bepaal die verandering in die konsentrasie van  $\text{NO}_2$
- Korrekte verhouding tussen  $\text{N}_2\text{O}_4$  en  $\text{NO}_2$
- Bepaal die aanvanklike konsentrasie van  $\text{N}_2\text{O}_4$
- Vervanging in verandering in konsentrasie en die aanvanklike konsentrasie van  $\text{N}_2\text{O}_4$  in die % verandering formule
- Finale antwoord



SA EXAM PAPERS

$$K_c = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]}$$

$$4,66 \times 10^{-3} = \frac{[\text{NO}_2]^2}{(0,491)} \quad (\text{a}) \checkmark$$

$$[\text{NO}_2] = 0,0478 \text{ mol} \cdot \text{dm}^{-3}$$

|                                                        | $\text{N}_2\text{O}_4$ | $2 \text{NO}_2 (\text{g})$ |
|--------------------------------------------------------|------------------------|----------------------------|
| Initial conc/<br><i>Aanvangskonsentrasie</i>           | 0,5 (d) $\checkmark$   | 0,03                       |
| Change in conc./<br><i>Verandering in konsentrasie</i> | $8,9 \times 10^{-3}$   | 0,0178 (b) $\checkmark$    |
| Equilibrium conc./<br><i>Ewewigskonsentrasie</i>       | 0,491                  | 0,0478                     |

(c) ratio/  
verhouding  $\checkmark$

$$\% [\text{N}_2\text{O}_4] = \frac{8,9 \times 10^{-3}}{0,5} \times 100 \% \quad (\text{e}) \checkmark$$

$$\% [\text{N}_2\text{O}_4] = 1,78 \% \quad (\text{f}) \checkmark$$

(6)  
[23]



**QUESTION/VRAAG 7**

- 7.1 7.1.1 An acid is a proton ( $H^+$ ) donor ✓✓ (2 or 0)  
*'n Suur is proton ( $H^+$ ) skenker (2 of 0)* (2)
- 7.1.2  $CH_3COOH + H_2O \rightleftharpoons CH_3COO^- + H_3O^+$  ✓  
 Ignore phases / *Ignoreer fases* (2)
- 7.1.3 Weak acid / *Swaksuur* ✓  
 $K_a$  value is less than 1 /  *$K_a$ -waarde is kleiner as 1* ✓ (2)
- 7.1.4 *Methanoic acid* ✓  
 It has the larger  $K_a$ -value / Ionises to a greater extent / It is the stronger acid. ✓  
 The higher the concentration of the  $H_3O^+$  the lower the pH-value. ✓  
*Metanoësuur*  
*Dit is het die groter  $K_a$ -waarde / Ioniseer tot 'n groter mate / Dit is die sterker suur.*  
*Hoe hoër die konsentrasie van  $H_3O^+$  hoe kleiner is is die pH-waarde* (3)
- 7.2 7.2.1 Burette / *Buret* ✓ (1)
- 7.2.2 To ensure reliable results / accurate results ✓  
*Om betroubare resultate / akkurate resultate te verseker* (1)
- 7.2.3 8,3–10 ✓ (1)
- 7.2.4  $C_2O_4^{2-} + H_2O \rightarrow HC_2O_4^- + OH^-$  ✓  
 (reactants / *reaktanse* and / *en* products / *produkte*)  
 Excess  $OH^-$  is produced / *Oormaat  $OH^-$  is geproduseer* ✓ (3)
- 7.3 7.3.1  $n = cV$  ✓  
 $n = (0,1)(25 \times 10^{-3})$  ✓  
 $n = 2,5 \times 10^{-3} \text{ mol}$  ✓ (3)



7.3.2

|                                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                                                                                                                                                                                                                 |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>OPTION 1</b><br><b>Marking criteria</b><br>(a) pH formula<br>(b) Subst. into pH-formula<br>(c) Subst. into $K_w = [\text{H}_3\text{O}^+][\text{OH}^-]$<br>(d) Final answer                                                                                                                                                                                                                              | <b>OPTION 2</b><br><b>Marking criteria</b><br>(a) Formula $\text{pH} + \text{pOH} = 14$<br>(b) Subst. into $\text{pH} + \text{pOH} = 14$<br>(c) Subst. into pOH formula<br>(d) Final answer                                                                                     |
| <b>OPSIE 1</b><br><b>Nasienkriteria</b><br>(a) pH-formule<br>(b) Vervang in pH-formule<br>(c) Vervang in $K_w = [\text{H}_3\text{O}^+][\text{OH}^-]$<br>(d) Finale antwoord                                                                                                                                                                                                                                | <b>OPSIE 2</b><br><b>Nasienkriteria</b><br>(a) Formule $\text{pH} + \text{pOH} = 14$<br>(b) Vervang in $\text{pH} + \text{pOH} = 14$<br>(c) Vervang in pOH-formule<br>(d) Finale antwoord                                                                                       |
| $\text{pH} = -\log[\text{H}_3\text{O}^+] \checkmark$ (a)<br>$12,52 = -\log[\text{H}_3\text{O}^+] \checkmark$ (b)<br>$[\text{H}_3\text{O}^+] = 3,02 \times 10^{-13} \text{ mol}\cdot\text{dm}^{-3}$<br><br>$K_w = [\text{H}_3\text{O}^+][\text{OH}^-]$<br>$1 \times 10^{-14} = (3,02 \times 10^{-13})[\text{OH}^-] \checkmark$ (c)<br>$[\text{OH}^-] = 0,033 \text{ mol}\cdot\text{dm}^{-3} \checkmark$ (d) | $\text{pH} + \text{pOH} = 14 \checkmark$ (a)<br>$12,52 + \text{pOH} = 14 \checkmark$ (b)<br>$\text{pOH} = 1,48$<br><br>$\text{pOH} = -\log[\text{OH}^-]$<br>$1,48 = -\log[\text{OH}^-] \checkmark$ (c)<br>$[\text{OH}^-] = 0,033 \text{ mol}\cdot\text{dm}^{-3} \checkmark$ (d) |

(4)

7.3.3

**POSITIVE MARKING FROM QUESTION 7.3.1 AND 7.3.2**  
**POSITIEWE NASIEN VAN VRAAG 7.3.1 EN 7.3.2**

**Marking criteria**

- (a) Subst. 0,033 and V into  $n = cV$   
 (b) Subst. 0,033V, 0,1V and n equivalence point in  
 $n_{\text{excess}} = n_{\text{totaal}} - n_{\text{ep}}$   
 (c) Subst. into  $V$  (total volume) =  $V$  (volume titrated) +  $V$  (acid)  
 (d) Subst. eq (2) into (1)  
 (e) Final answer

**Nasienkritria**

- (a) Vervang 0,033 en V in  $n = cV$   
 (b) Vervang 0,033V, 0,1V en n ekwivalente punt in  
 $n_{\text{oormaat}} = n_{\text{totaal}} - n_{\text{ekw}}$   
 (c) Vervang in  $V$  (totale volume) =  $V$  (volume getitreer) +  $V$  (suur)  
 (d) Vervang vergelyking (2) in (1)  
 (e) Finale antwoord

$$n(\text{excess}) = n(\text{total}) - n(\text{equivalence point})$$

$$n(\text{oormaat}) = n(\text{totaal}) - n(\text{ekwivalente punt})$$

$$cV(\text{total volume}) = cV(\text{volume titrated}) - n(\text{equivalence point})$$

$$cV(\text{totaal volume}) = cV(\text{volume getitreer}) - n(\text{ekwivalente punt})$$

$$0,033V(\text{total volume}) \checkmark \text{ (a)} = 0,1V(\text{volume titrated}) - 2,5 \times 10^{-3} \dots (1) \checkmark$$

(b)

$$V(\text{total volume}) = V(\text{volume titrated}) + V(\text{acid})$$

$$V(\text{totale volume}) = V(\text{getitreer}) + V(\text{suur})$$



$$V \text{ (total volume)} = V \text{ (volume titrated)} + 25 \times 10^{-3} \dots (2) \text{ (c)} \checkmark$$

Subst. equation (2) into (1) / Vervang vergelyking (2) in (1)

$$0,033 [\text{Volume titrated} + 25 \times 10^{-3}] \checkmark \text{ (d)} = 0,1 \text{ Volume titrated} - 2,5 \times 10^{-3}$$

$$\text{Volume titrated} / \text{volume getitreer} = 0,050 \text{ dm}^3$$

$$\text{Volume titrated} / \text{volume getitreer} = 50 \text{ cm}^3 \checkmark \text{ (e)} \quad (5)$$

[27]

**TOTAL/TOTAAL: 150**

