

# SA's Leading Past Year

## Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ [www.saexampapers.co.za](http://www.saexampapers.co.za)





**GAUTENG PROVINCE**

EDUCATION  
REPUBLIC OF SOUTH AFRICA

# **JUNE EXAMINATION GRADE 12**

**2025**

## **MARKING GUIDELINES**

**AGRICULTURAL SCIENCES**

**(PAPER 1)**

**11 pages**

**SECTION A****QUESTION 1**

- 1.1 1.1.1 C ✓✓
- 1.1.2 D ✓✓
- 1.1.3 C ✓✓
- 1.1.4 A ✓✓
- 1.1.5 C ✓✓
- 1.1.6 B ✓✓
- 1.1.7 D ✓✓
- 1.1.8 C ✓✓
- 1.1.9 A ✓✓
- 1.1.10 B ✓✓ (10 x 2) (20)
- 1.2 1.2.1 Both A and B ✓✓
- 1.2.2 A only ✓✓
- 1.2.3 Both A and B ✓✓
- 1.2.4 B only ✓✓
- 1.2.5 None ✓✓ (5 x 2) (10)
- 1.3 1.3.1 Peristalsis ✓✓
- 1.3.2 Fodder/Feed flow ✓✓
- 1.3.3 Bunching ✓✓
- 1.3.4 Cryptorchidism ✓✓
- 1.3.5 Hydrocephalus ✓✓ (5 x 2) (10)





- 1.4 1.4.1 Amylase/Ptyalin ✓
- 1.4.2 Cafeteria style/Free choice/Ad lib ✓
- 1.4.3 Foot and Mouth Disease/FMD ✓
- 1.4.4 Mummification ✓
- 1.4.5 Colostrum/Beestings ✓ (5 x 1) (5)

**TOTAL SECTION A: 45**

## SECTION B

### QUESTION 2: ANIMAL NUTRITION

#### 2.1 The digestive system of a ruminant

##### 2.1.1 The name of the animal

Cattle/sheep/goat ✓ (1)

##### 2.1.2 Processes illustrated by arrows

**A** – Swallowing/peristalsis/ingestion/intake of food ✓

**B** – Regurgitation/retro-peristalsis ✓

**C** – Re-swallowing/peristalsis ✓ (3)

##### 2.1.3 TWO advantages of regurgitation/retro-peristalsis

- Breaks down food into finer particles ✓
  - Increases surface area ✓
  - Stimulates secretion of saliva ✓
  - Maintains rumen pH level ✓
  - Improves the mixing of food ✓
  - Helps with the forming of bolus ✓
- (Any 2) (2)

#### 2.2 The schematic representation of the components of feed:

##### 2.2.1 Identification of substances

**A** – Dry matter ✓

**B** – Inorganic matter/minerals/elements ✓

**C** – Vitamins ✓ (3)

##### 2.2.2 Difference between oil and fat

**Oil** – unsaturated/liquid at room temperature/plant origin/ Double bonds ✓

**Fat** – saturated/ solid at room temperature/animal origin/ Single bonds ✓ (2)





## 2.2.3 The end-products of digestion in pigs

- (a) **Carbohydrates** – Glucose/energy ✓ (1)
- (b) **Protein** – Amino acids ✓ (1)

## 2.2.4 Two functions of water in digestion

- Acts as a solvent/assists in the absorption of nutrients ✓
  - Moisturises the swallowed food ✓
  - Provides a suitable environment for micro-organisms ✓
  - Helps with the breaking down of food ✓
  - Prevents constipation ✓
  - Transports nutrients ✓
  - Facilitates the excretion of waste products ✓
- (Any 2) (2)

## 2.3 Minerals, vitamins and their deficiency symptoms

## 2.3.1 Missing information in A, B and C

- A** – Zinc/Zn ✓
- B** – Metritis/inflammation of the udder ✓
- C** – Stiff lamb/muscle dystrophy/white muscle ✓ (3)

## 2.3.2 Methods used to supplement the following

- (a) **Vitamin A** – Injections/supplementary rations ✓ (1)
- (b) **Soluble minerals** – Dissolve minerals in drinking water/dosing ✓ (1)





## 2.4 Calculation of digestibility co-efficient

$$\begin{aligned}
 2.4.1 \quad \text{DM of hay} &= \frac{8 \times 30}{100} \\
 &= 2,4 \text{ kg} \\
 &30 - 2,4 \\
 &= 27,6 \text{ kg} \checkmark
 \end{aligned}$$

$$\begin{aligned}
 \text{DC} &= \frac{\text{Dry matter intake (kg)} - \text{Dry mass manure (kg)}}{\text{Dry matter intake (kg)}} \times \frac{100}{1} \checkmark \\
 &= \frac{27,6 \text{ kg} - 12 \text{ kg}}{27,6 \text{ kg}} \times 100 \checkmark \\
 &= 56,52 \checkmark \% \checkmark \text{ or } 57 \checkmark \% \checkmark
 \end{aligned}$$

(5)

2.4.2 56,52% ✓ of feed has been digested and absorbed ✓

OR

43,48% ✓ was excreted and not digested and absorbed ✓

(Any 1) (2)

## 2.5 Nutritive ratio

## 2.5.1 The percentage of digestible non-nitrogenous substances

$$75\% - 20\% \checkmark = 55\% \checkmark \quad (2)$$

## 2.5.2 Calculation of the nutritive ratio

$$\begin{aligned}
 \text{NR} &= 1: \frac{\% \text{ Digestible non-nitrogen substances}}{\% \text{ Digestible protein}} \checkmark \\
 &\text{or} \\
 \text{NR} &= \frac{1: \% \text{ TDN} - \% \text{ DP}}{\% \text{ DP}}
 \end{aligned}$$

$$\begin{aligned}
 &= 1 : \frac{75\% - 20\%}{20\%} \checkmark \\
 &\text{or} \\
 &\frac{55\%}{20\%}
 \end{aligned}$$

$$= 1 : 2,75 \checkmark \text{ OR } 1:3 \checkmark$$

(3)

## 2.6 Fodder flow plan

## Calculation of feed required for a year

$$\begin{aligned}
 &15 \times 30 \text{ animals} \times 30 \text{ days} \times 6 \text{ months} \checkmark \\
 &= 81\,000 \text{ kg} \checkmark
 \end{aligned}$$

$$\begin{aligned}
 &81\,000 \text{ kg} + 108\,000 \text{ kg} \\
 &= 189\,000 \text{ kg} \checkmark
 \end{aligned}$$

(3)

[35]



**QUESTION 3: ANIMAL PRODUCTION, PROTECTION AND CONTROL****3.1 Farming systems****3.1.1 Identification of production system**

Semi intensive system /Backyard/free-range ✓ (1)

**3.1.2 Two advantages of the production system**

- Less expensive/cheaper ✓
- Easy to manage ✓
- No special equipment needed ✓
- No specialised/expert knowledge needed ✓
- More environmentally friendly ✓/ Improved Animal welfare (Any 2) (2)

**3.1.3 Two problems associated with the production system**

- Easy to contract diseases ✓
- Poor feeding/feeding on less nutritious food ✓
- High risk for predators ✓
- More feed energy is utilised for non-production purpose/low production output/slow growth rate ✓
- High risk towards theft of animals ✓
- Exposed to extreme environmental conditions ✓ (Any 2) (2)

**3.2 Equipment used in handling farm animals**

3.2.1 B/E ✓ (1)

3.2.2 B/C ✓ (1)

3.2.3 B/D ✓ (1)

3.2.4 A ✓ (1)





## 3.3 Identification of the handling equipment

## 3.3.1 Diagram A

Rope ✓

## Diagram B

Nose holder ✓

(2)

## 3.3.2 One of the purposes of the equipment in Diagram A

Handled to make it to lie down/bring the animal down ✓

(1)


## 3.3.3 Part of the head that is used to grip the animal

- Nostril ✓
- Ear ✓

(Any 1) (1)

## 3.4 The temperature ranges in farm animals and their expected growth rates, expressed as averages for their population.

## 3.4.1 Graph on Temperature range and growth rates of cows and pigs



Rubric for marking of the graph

Criteria	Yes: (1 mark)	No: (0 marks)
1 Line graph	1 ✓	
2 X-axis correctly labelled	1 ✓	
3 Y-axis correctly labelled	1 ✓	
4 Plotting growth rate for pigs	1 ✓	
5 Plotting growth rate for cows	1 ✓	
6 Correct heading	1 ✓	

(6)





**3.4.2 Farm animals that require more environmental control**

Pigs ✓

**Justification**

They are most affected by lower temperatures (lower growth rate at lower temperatures) ✓

(2)

**3.5 Diseases affecting farm animals****A** – Bacteria ✓**B** – Blisters on the tongue, nose, lips, in the mouth and between the toes ✓**C** – Protozoan ✓**D** – Bont tick ✓**E** – Ringworm ✓

(5)

**3.6 The diagram indicating various stages of the life cycle of a parasite****3.6.1 Name of parasite**

Bont tick ✓

**Type of parasite**

External parasite/ecto-parasite ✓

(2)

**3.6.2 Classification of parasites**

Three-host tick ✓

(1)

**3.6.3 Two economic implications**

- Banning of exports/decrease in trade ✓
- High treatment/medication costs to control/prevent diseases ✓
- Decreased production ✓
- Loss of income/profit ✓
- Loss of livestock/death ✓

(Any 2) (2)

**3.7 Salt poisoning****3.7.1 Two symptoms of salt poisoning**

- Excessive salivation ✓
- Increased thirst ✓
- Vomiting and regurgitation ✓
- Constipation ✓
- Aggressiveness ✓
- Hypersensitivity ✓
- Red and dry mucous membranes of mouth ✓
- Increased urination and defecation ✓
- Abnormal pain and diarrhoea ✓

(Any 2) (2)

**3.7.2 Two measures to treat animals with salt poisoning**

- Provision of fresh, clean water in small amounts ✓
- Provision of isotonic saline solution/hypertonic dextrose ✓
- Remove source of salt poisoning ✓

(Any 2) (2)

**[35]**

**QUESTION 4: ANIMAL REPRODUCTION****4.1 Female reproductive organs****4.1.1 Identification of the parts of the reproductive system****A** – uterine horn ✓**C** – fallopian tube/oviduct ✓**E** – cervix ✓**F** – vagina ✓

(4)

**4.1.2 Parts linking with the following functions:**

(a) Serves as the birth canal – F ✓

(1)

(b) Site of fertilisation – C ✓

(1)

(c) Organ not directly involved in reproduction – I ✓

(1)

(d) Deposition of semen during artificial insemination – J ✓

(1)

**4.2 Infertility and sterility in bulls****4.2.1 Differentiation between infertility and sterility**• **Sterility** is the total loss of fertility ✓• **Infertility** is the temporary loss of fertility ✓

(2)

**4.2.2 TWO congenital defects that lead to sterility**

• Hypoplasia ✓

• Cryptorchidism ✓

• Hermaphroditism ✓

• Sperm defects ✓

(Any 2) (2)

**4.2.3 TWO conditions that may cause a bull to be unable to copulate**

• Injuries to the penis ✓

• Defective penis/corkscrew/too short ✓

• Poorly developed hind legs ✓

• Diseases causing inflammation of the joints ✓

(Any 2) (2)





#### 4.3 The diagram that illustrates a technique used in animal reproduction

##### 4.3.1 Process illustrated by the diagram

Nuclear transfer/cloning ✓

(1)

##### 4.3.2 Identification of cells

**A** – Recipient cell with nucleus/egg cell/ovum ✓

**B** – The nucleus of the donor cell ✓

**D** – The fused cell ✓

(3)

##### 4.3.3 Two types of the process

• Reproductive cloning ✓

• Therapeutic cloning ✓

(2)

#### 4.4 Difficulty in giving birth

##### 4.4.1 Scientific term for difficult births

Dystocia ✓

(1)

##### 4.4.2 Three reasons for difficult births in heifers

- Heifers are physically smaller ✓ and less developed (younger)/age ✓
- Incorrect presentation/position/posture/ Posterior presentation ✓
- Too large foetus/hydrocephalus ✓
- Deformities of the foetus ✓
- Torsion/twisting of the foetus ✓
- Prolapsed uterus ✓
- Multiple births/twins ✓
- Size of pelvic area ✓
- Weak/ ineffective labour ✓
- Cervix failing to dilate ✓
- Prolonged gestation/pregnancy period ✓
- Malnutrition ✓
- Diseases ✓

(Any 3) (3)

##### 4.4.3 TWO managerial measures to reduce difficult births

- Use bulls renowned for small calves/low birth weight ✓
- Mate heifers at the ideal age/mass/not too early ✓
- Use a controlled/well-planned breeding season ✓
- Well planned feeding programme/avoid overfeeding ✓
- Planned health programme ✓

(Any 2) (2)

##### 4.4.4 Definition of placenta retention

- The failure to expel the placenta/membranes ✓
- within 12 hours after parturition/birth ✓
- with negative effects/complications ✓

(Any 2) (2)





#### 4.5 Normal lactation

##### 4.5.1 Identification of the week when the cow reached her maximum production

Week 8 ✓

(1)

##### 4.5.2 TWO benefits of the milk produced within the first 3 days of parturition for the calf

- It provides antibodies to increase the calf's resistance to diseases ✓
- Supplies nutrients to the calf ✓
- Necessary for the normal growth, functioning and maturation of the alimentary canal ✓
- Higher butterfat content ✓

(Any 2) (2)

##### 4.5.3 Explanation of the relationship between the percentage of butter fat content and the quantity of milk produced

When milk production reached its maximum during week 8 and decreased until week 42 ✓ butterfat content decreased at week 8 and increased until week 42 ✓

(2)

##### 4.5.4 Period in lactating cow after 42 weeks

Dry period ✓

(1)

##### 4.5.5 Importance of dry period

It gives the glandular tissue of the udder time to recover ✓

(1)

[35]

**TOTAL SECTION B: 105**

**TOTAL: 150**

