

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great
Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



SA EXAM PAPERS

SA EXAM PAPERS

Proudly South African



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

**SENIOR CERTIFICATE EXAMINATIONS/
NATIONAL SENIOR CERTIFICATE EXAMINATIONS
SENIORSERTIFIKAAT-EKSAMEN/
NASIONALE SENIORSERTIFIKAAT-EKSAMEN**

**PHYSICAL SCIENCES: CHEMISTRY (P2)
FISIESE WETENSKAPPE: CHEMIE (V2)**

MAY/JUNE 2025/MEI/JUNIE 2025

MARKING GUIDELINES/NASIENRIGLYNE

DEPARTMENT OF BASIC EDUCATION
PRIVATE BAG X895, PRETORIA 0001
01 -06- 2025
APPROVED MARKING GUIDELINE
PUBLIC EXAMINATION

MARKS/PUNTE: 150

**These marking guidelines consist of 22 pages.
Hierdie nasienriglyne bestaan uit 22 bladsye.**

Approved
Alexander (IM)
DBE
01/06/2025


Approved
Dusane
DBE IM
01-06-2025

Approved
Umalusi
Umalusi Ext Mt.
01/06/2025

Approved!
Vubaluv.
UMALUUP
EXTERNAL
MODERATOR
1/6/2025

**QUESTION 1/VRAAG 1**

- 1.1 B ✓✓ ACCEPT/AANVAAR 2 (2)
- 1.2 D ✓✓ (2)
- 1.3 D ✓✓ (2)
- 1.4 A ✓✓ (2)
- 1.5 B ✓✓ (2)
- 1.6 C ✓✓ (2)
- 1.7 D ✓✓ (2)
- 1.8 B ✓✓ (2)
- 1.9 C ✓✓ (2)
- 1.10 C ✓✓ (2)
- [20]**

**QUESTION 2/VRAAG 2**

- 2.1 Compounds with one or more multiple bonds between C atoms in the hydrocarbon chain. ✓✓ (2 or 0)
Verbindings met een of meer meervoudige bindings tussen C-atome in die koolwaterstofkettings. (2 of 0)

OR/OF

A hydrocarbon with two or more bonds between the C-atoms.
'n Koolwaterstof met twee of meer bindings tussen die C-atome.

OR/OF

Hydrocarbons containing not only single bonds between C atoms.
Koolwaterstowwe wat nie slegs enkelbindings tussen die C-atome bevat nie.

ACCEPT/AANVAAR:

Compounds with one or more double/triple bonds between C atoms in the hydrocarbon chain.
Verbindings met een of meer dubbel/trippelbindings tussen C-atome in die koolwaterstofkettings. (2)

- 2.2
- 2.2.1 E ✓ (1)
- 2.2.2 F ✓ (1)
- 2.3 Ketones/Ketone ✓
Aldehydes/Aldehiede ✓ (2)



Handwritten signatures and initials




2.4 Tertiary/Tersiêre ✓

The hydroxyl group/functional group (-OH) is bonded to a C atom that is bonded to three other C atoms. ✓


Die hidroksiel/funksionele groep (-OH) is gebind aan 'n C-atoom wat aan drie ander C-atome gebind is.

OR/OF

The functional group (—C—) is bonded to three other C atoms.



Die funksionele groep (—C—) is gebind aan drie ander C-atome.



(2)

2.5

2.5.1

Marking criteria:

- Correct stem, i.e. hexane. ✓
- Both substituent (ethyl and iodo) correctly identified. ✓
- IUPAC name completely correct including numbering, sequence, hyphens and commas. ✓

Nasienkriteria:

- Korrekte stam, d.i. heksaan. ✓
- Beide substituent (etiel en jodo) korrek geïdentifiseer. ✓
- IUPAC-naam heeltemal korrek insluitende nommering, volgorde, koppeltekeens en kommas. ✓

3-ethyl-4-iodohexane/3-etiel-4-jodoheksaan ✓✓✓

(3)

2.5.2

Marking criteria/Nasienkriteria:

- Correct stem and substituents: methyl and propanol ✓
Korrekte stam en substituent: metiel en propanol
- IUPAC name completely correct including numbering, sequence, hyphens and commas. ✓
IUPAC-naam heeltemal korrek insluitende nommering, volgorde, koppeltekeens en kommas.


2-methylpropan-1-ol/ 2-methyl-1-propanol/ methylpropan-1-ol/

methyl-1-propanol ✓✓

2-metielpropan-1-ol/ 2-metiel-1-propanol / metielpropan-1-ol/

metiel-1-propanol

(2)



SA EXAM PAPERS

Handwritten signatures and initials




2.6

2.6.1 Esterification/Condensation/Verestering/Esterifikasie/Kondensasie ✓

(1)

2.6.2

**Marking criteria/Nasienkriteria:**

- Functional group correct. ✓
Funksionele groep korrek.
- Whole structure correct. ✓
Hele struktuur korrek.

IF/INDIEN

- More than one functional group/wrong functional group:
Meer as een funksionele groep/foutiewe funksionele groep: 0/2
- If condensed structural formulae used/Indien gekondenseerde struktuurformules gebruik: Max./Maks. 1/2


(2)

2.7

2.7.1 C₂H₄O ✓

(1)

2.7.2

**Marking criteria/Nasienkriteria:**


- Correct functional group. ✓
Korrekte funksionele groep.
- Whole structure correct. ✓
Hele struktuur korrek.

IF/INDIEN

- More than one functional group/wrong functional group:
Meer as een funksionele groep/foutiewe funksionele groep: 0/2
- If condensed structural formulae used/Indien gekondenseerde struktuurformules gebruik: Max./Maks. 1/2

(2)

[19]



SA EXAM PAPERS

**QUESTION 3/VRAAG 3**

- 3.1.1 (A series of organic) compounds that can be described by the same general formula. ✓✓ (2 or 0)

OR

(A series of organic) compounds in which one member differs from the next by a CH_2 group.

(’n Reeks organiese) verbindings wat deur dieselfde algemene formule beskryf kan word. ✓✓ (2 of 0)

OF

(’n Reeks organiese) verbindings waarin die een lid van die volgende verskil met ’n CH_2 -groep (2)

3.1.2

- (a) Formyl (group)/Formiel(groep) ✓ (1)

(b)

Marking criteria:

- Correct chain length, i.e. Meth. ✓
- Everything else correct. ✓

Nasienkriteria:

- Korrekte kettinglengte d.i. Met. ✓
- Alles verder reg ✓

Methanal/Metanaal ✓✓ (2)

3.1.3

- (a) Homologous series/Functional group/Type of intermolecular forces/Straight chain/Atmospheric pressure ✓
Homoloë reeks/Funksionele groep/Tipe intermolekulêre kragte/ Reguitketting/ Atmosferiese druk (1)

- (b) The boiling points of the carboxylic acids increase with an increase in the chain length/the number of carbon atoms/surface area/molecular mass./


OR

The boiling points of the carboxylic acids decrease with a decrease in the chain length/number of carbon atoms/surface area/molecular mass. ✓

Die kookpunte van die karboksielsure neem toe met ’n toename in die kettinglengte/aantal koolstofatome/reaksieoppervlak/molekulêre massa./

OF

Die kookpunte van die karboksielsure neem af met ’n afname in die kettinglengte/aantal koolstofatome/reaksieoppervlak/molekulêre massa. (1)

**SA EXAM PAPERS**

Handwritten signatures and initials: J.S., V.M.P., A., and a large stylized signature.



(c)

Marking criteria:

For increasing or decreasing number of C atoms

- Compare the strength of intermolecular forces. ✓
- Compare the energy required to overcome intermolecular forces. ✓

Nasienkriteria:

Vir toename of afname in aantal C-atome

- Vergelyk die sterkte van intermolekulêre kragte. ✓
- Vergelyk die energie benodig om intermolekulêre kragte te oorkom. ✓

As the number of C atoms/chain length/surface area/contact area/molecular mass increases

- The strength of intermolecular/London/dispersion forces increases. ✓
- More energy is needed to overcome intermolecular forces/London/dispersion forces. ✓

OR

As the number of C atoms/chain length/surface area/contact area/molecular mass decreases

- The strength of intermolecular/London/dispersion forces decreases. ✓
- Less energy is needed to overcome intermolecular forces/London/dispersion forces. ✓

Met toename in aantal C-atome/kettinglengte/reaksieoppervlak/kontakarea/molekulêre massa.

- Die sterkte van die die intermolekulêre kragte/Londonkragte/dispersiekragte neem toe.
- Meer energie word benodig om die intermolekulêre kragte/Londonkragte/dispersiekragte te oorkom/breek.

OF


Met afname in aantal C-atome/kettinglengte/reaksieoppervlak/kontakarea/molekulêre massa.

- Die sterkte van die die intermolekulêre kragte/Londonkragte/dispersiekragte neem af.
- Minder energie word benodig om die intermolekulêre kragte/Londonkragte/dispersiekragte te oorkom/breek

(2)

3.1.4 75 °C ✓

(1)



Sub. VHF
M
DL



3.2

Marking criteria:

- Higher than ✓
- State that carboxylic acids have more than one (two) site for hydrogen bonding and alcohols have one site for hydrogen bonding. ✓
- Comparing the strength of IMFs. ✓
- Comparing the number of molecules in a vapour phase at a given temperature/energy needed to overcome IMFs. ✓

Nasienkriteria:

- Hoër as ✓
- Stel dat karboksiesure het meer as een (twee) plekke vir waterstofbindings en dat alkohole een plek het vir waterstofbinding. ✓
- Vergelyk die sterkte van die IMK's/energie benodig om IMK's te oorkom. ✓
- Vergelyk die hoeveelheid molekules in die dampfase by 'n gegewe temperatuur/energie nodig om die IMK te oorkom. ✓

- Higher than ✓
- Compound B/CH₃CH₂CH₂COOH/Carboxylic acid/Butanoic acid has (in addition to London forces and dipole-dipole forces), more than one site (two) for hydrogen bonding between molecules and compound A/CH₃CH₂CH₂CH₂CH₂OH/Alcohol/Pentan-1-ol has (in addition to London forces and dipole-dipole forces) one site for hydrogen bonding between molecules. ✓
- Intermolecular forces in compound B/CH₃CH₂CH₂COOH/Carboxylic acids/Butanoic acid are stronger. ✓
- More energy needed to overcome/break intermolecular forces in compound B/CH₃CH₂CH₂COOH/Carboxylic acid/Butanoic acid.


OR

- At a given temperature there will be fewer molecules of compound B/CH₃CH₂CH₂COOH/Carboxylic acids/Butanoic acid in the vapour phase. ✓

- Higher than ✓
- Compound A/CH₃CH₂CH₂CH₂CH₂OH/Alcohol/Pentan-1-ol has (in addition to London forces and dipole-dipole forces) one site for hydrogen bonding between molecules and compound B/CH₃CH₂CH₂COOH/Carboxylic acid/Butanoic acid has, (in addition to London forces and dipole-dipole forces), more than one site (two) for hydrogen bonding between molecules. ✓
- Intermolecular forces in compound A/CH₃CH₂CH₂CH₂CH₂OH/Alcohol/Pentan-1-ol are weaker. ✓
- Less energy needed to overcome/break intermolecular forces in compound A/CH₃CH₂CH₂CH₂CH₂OH/Pentan-1-ol/Alcohol.

OR

- At a given temperature there will be more molecules of compound A/CH₃CH₂CH₂CH₂CH₂OH/Alcohol/ Pentan-1-ol in the vapour phase. ✓




Handwritten signatures and initials.



- Hoër as
 - Verbinding B/CH₃CH₂CH₂COOH/Karboksielsure/Butanoësuur het, (in toevoeging tot Londonkragte en dipool-dipoolkragte), meer as een posisie (twee) vir waterstofbinding tussen molekule en verbinding A/CH₃CH₂CH₂CH₂CH₂OH/Alkohol/Pentan-1-ol het, (in toevoeging tot Londonkragte en dipool-dipoolkragte), een posisie vir waterstofbinding tussen molekule.
 - Intermolekulêre kragte in verbinding B/CH₃CH₂CH₂COOH/Karboksielsure/Butanoësuur is sterker.
 - Meer energie word benodig om intermolekulêre kragte in verbinding B/CH₃CH₂CH₂COOH/Karboksielsure/Butanoësuur te oorkom/breek
- OF**
- By 'n gegewe temperatuur sal daar minder molekules van verbinding B/CH₃CH₂CH₂COOH/Karboksielsure/Butanoësuur in die dampfase wees.
-
- Hoër as
 - Verbinding A/CH₃CH₂CH₂CH₂CH₂OH/Alkohol/Pentan-1-ol het, (in toevoeging tot Londonkragte en dipool-dipoolkragte), een posisie vir waterstofbinding tussen molekule en verbinding B/CH₃CH₂CH₂COOH/Karboksielsure/Butanoësuur het, (in toevoeging tot Londonkragte en dipool-dipoolkragte), meer as een posisie (twee) vir waterstofbinding tussen molekule.
 - Intermolekulêre kragte in verbinding A/CH₃CH₂CH₂CH₂CH₂OH/Alkohol/Pentan-1-ol is swakker.
 - Minder energie word benodig om intermolekulêre kragte in A/CH₃CH₂CH₂CH₂CH₂OH/Alkohol/Pentan-1-ol te oorkom/breek.
- OF**
- By 'n gegewe temperatuur sal daar meer molekules van verbinding A/CH₃CH₂CH₂CH₂CH₂OH/Alkohol/Pentan-1-ol in die dampfase wees.

(4)
[14]



SA EXAM PAPERS

Copyright reserved/Kopiereg voorbehou

Proudly South African

Please turn over/Blaai om asseblief

mk. van
13

**QUESTION 4/VRAAG 4**

4.1

4.1.1 Hydrogenation/Hidrogenering/Hidrogenasie ✓ (1)

4.1.2 Dehydration/Dehidrasie/Dehidratering ✓ (1)

4.2

Marking criteria:

- Correct chain length, i.e. But. ✓
- Everything else correct: IUPAC name completely correct including numbering. ✓

Nasienkriteria:

- Korrekte kettinglengte d.i. But. ✓
- Alles verder reg: IUPAC-naam heeltemal korrek insluitende nommering. ✓

Butan-1-ol/1-butanol ✓✓ (2)

4.3

4.3.1

Marking criteria/Nasienkriteria:

- Whole condensed structural formula of alkene correct. ✓
Hele gekondenseerde struktuurformule van die alkeen korrek.
- H₂O. ✓
- Hydroxyl group/OH. ✓
Hidroksielgroep/OH.
- Whole condensed structural formula of alcohol correct (OH on second C-atom). ✓
Hele gekondenseerde struktuurformule van alkohol korrek (OH op tweede C-atoom)

IF/INDIEN


- Any additional reactants or products /Enige addisionele reaktanse of produkte:
Deduct 1 mark/Trek 1 punt af.

- Structural formulae used/Struktuurformule gebruik. Max./Maks. 3/4


- Molecular formulae used/Molekulêre formule gebruik. Max./Maks. 1/4

- Only reactants without arrow/Slegs reaktanse sonder pyl Max./Maks. 2/4

Marking rule 6.3.10/Nasienreël 6.3.10

**OR/OF**

(4)

4.3.2 Sulphuric acid/H₂SO₄/Phosphoric acid/H₃PO₄/Swawelsuur/Fosforsuur ✓ (1)**SA EXAM PAPERS**

Proudly South African

Handwritten signatures and initials.





4.4

Marking criteria/Nasienkriteria:

- $X_2 = Br_2/Cl_2$. ✓
- Whole structural formula of haloalkane correct. ✓
Hele struktuurformule van haloalkaan korrek.
- $H_2O/NaOH/KOH/LiOH$. ✓
- Whole structural formula of alcohol correct. ✓
Hele struktuurformule van alkohol korrek.
- $HX/NaX/KX/LiX$ where/waar $X = Br/Cl$ ✓

IF/INDIEN

- Any additional reactants or products /Enige addisionele reaktanse of produkte:
Max./Maks. $\frac{4}{5}$
- Condensed structural formulae used/Gekondenseerde struktuurformule gebruik:
deduct 1 mark/trek 1 punt af.
- If inorganic product does not correspond with inorganic reactant: no mark for inorganic product./Indien anorganiese produk nie met die anorganiese reaktans ooreenstem nie, geen punt vir anorganiese produk.
- Molecular formulae used:/Molekulêre formule gebruik: Max./Maks. $\frac{3}{5}$
- Marking rule 6.3.10/Nasienreël 6.3.10



Ignore phases./Ignoreer fases.

Marking criteria/Nasienkriteria:

- C_4H_{10} ✓ O_2 , CO_2 and/en H_2O ✓ Balancing/Balansering ✓
- Ignore double arrows./Ignoreer dubbelpyle.
- Marking rule 6.3.10/Nasienreël 6.3.10.

IF/INDIEN:

- Structural formulae C_4H_{10} used:/Struktuurformule C_4H_{10} gebruik: Max./Maks. $\frac{2}{3}$
- Balancing mark only if everything else is correct/
Balanseringspunt slegs indien alles korrek.

(3)
[17]

**QUESTION 5/VRAAG 5**

5.1

NOTE/NOTAGive the mark for per unit time only if in context of reaction rate.Gee die punt vir per eenheidtyd slegs indien in konteks van reaksietempo.**ANY ONE:**

- Change in concentration ✓ of products/reactants per (unit) time. ✓
- Change in amount/number of moles/volume/mass of products or reactants per (unit) time.
- Amount/number of moles/volume/mass of products formed/reactants used per (unit) time.
- Rate of change in concentration/amount/number of moles/volume/mass. ✓✓ (2 or 0)

ENIGE EEN:

- Verandering in konsentrasie ✓ van produkte/reaktanses per (eenheid)tyd. ✓
- Verandering in hoeveelheid/getal mol/volume/massa van produkte of reaktanses per (eenheid)tyd.
- Hoeveelheid/getal mol/volume/massa van produkte gevorm/reaktanses gebruik per (eenheid)tyd.
- Tempo van verandering in konsentrasie/ hoeveelheid/getal mol/ volume/ massa. ✓✓ (2 of 0) (2)

5.2

ANY ONE:

Temperature ✓/

(Initial) amount/Mass of magnesium carbonate/Surface area


ENIGE EEN:

Temperatuur ✓/

(Aanvanklike) hoeveelheid/Massa van magnesiumkarbonaat/

Reaksieoppervlak (1)

5.3

CO₂/gas escapes from the reaction flask. ✓CO₂/gas ontsnap uit die reaksiefles. ✓ (1)

Handwritten signatures and initials: J. van der Merwe, V. P., and others.



5.4

Marking criteria:

- (a) Mass subtraction ✓
 (b) Formula: $n = \frac{m}{M}$ or $V = nV_m$ ✓
 (c) Substitute $M = 44 \text{ g} \cdot \text{mol}^{-1}$ in
 $n(\text{CO}_2) = \frac{m}{M}$ with $m(\text{CO}_2)$ from (a) ✓
 (d) Substitute $24,5 \text{ dm}^3$ in $V = nV_m$ with
 $n(\text{CO}_2)$ ✓
 (e) Substitute V_{CO_2} and 120 in rate formula ✓
 (f) Final correct answer:
 $2,92 \times 10^{-3} (\text{dm}^3 \cdot \text{s}^{-1})$ ✓
 Range: $2,08 \times 10^{-3}$ to 3×10^{-3}

Nasienkriteria:

- (a) Afrek van massas. ✓
 (b) Formula: $n = \frac{m}{M}$ or $V = nV_m$ ✓
 (c) Vervang $M = 44 \text{ g} \cdot \text{mol}^{-1}$ in
 $n(\text{CO}_2) = \frac{m}{M}$ met $m(\text{CO}_2)$ van (a) ✓
 (d) Vervang $24,5 \text{ dm}^3$ in $V = nV_m$ met
 $n(\text{CO}_2)$ ✓
 (e) Vervang V_{CO_2} en 120 in tempoformule ✓
 (f) Finale korrekte antwoord:
 $2,92 \times 10^{-3} (\text{dm}^3 \cdot \text{s}^{-1})$ ✓
 Gebied: $2,08 \times 10^{-3}$ tot 3×10^{-3}

$$m(\text{CO}_2) = 144,5 - 143,87 \checkmark \text{ (a)}$$

$$= 0,63 \text{ g}$$

$$n(\text{CO}_2) = \frac{m}{M} \checkmark \text{ (b)}$$

$$= \frac{0,63}{44} \checkmark \text{ (c)}$$

$$= 1,43 \times 10^{-2} \text{ mol}$$

$$V(\text{CO}_2) = nV_m$$

$$= (1,43 \times 10^{-2})(24,5) \checkmark \text{ (d)}$$

$$= 0,35 \text{ dm}^3$$

$$\text{Ave rate/}$$

$$\text{gem tempo} = \frac{\Delta V(\text{CO}_2)}{\Delta t}$$

$$= \frac{0,35 - (0)}{120 - (0)} \checkmark \text{ (e)}$$

$$= 2,92 \times 10^{-3} (\text{dm}^3 \cdot \text{s}^{-1}) \checkmark \text{ (f)}$$

(6)

5.5

Marking criteria:

- A ✓
- Comparison of the curves of the graph ✓
- Comparison of concentration of HCl (from table) ✓
- Explanation of collision theory for LOWER concentration ✓✓

Nasienkriteria:

- A ✓
- Vergelyk die kurwes van grafiek ✓
- Vergelyk die konsentrasie van HCl (vanaf tabel) ✓
- Verduidelik botsingsteorie vir LAER konsentrasie ✓✓

A ✓

- Gradient is least steep/lowest reaction rate/least amount of gas produced in 120 s. ✓
- Lowest concentration of $\text{HCl}(\text{aq})$. ✓
- Least/Less particles per unit volume. ✓
- Least/Less effective collisions per unit time/second. ✓ **OR**
 Lowest/Lower frequency of effective collisions.
- *Gradient is die laagste/laagste reaksietempo/minste hoeveelheid gas geproduseer in 120 s.* ✓
- Laagste konsentrasie van $\text{HCl}(\text{aq})$. ✓
- Minste/Minder deeltjies per eenheidsvolume. ✓
- Minste/Minder effektiewe botsings per eenheidstyd/sekonde. ✓ **OF**
 Laagste/Laer frekwensie van effektiewe botsings.

(5)

5.6

The same/ Dieselfde ✓

The same amount MgCO_3 is used in each experiment. ✓Dieselfde hoeveelheid MgCO_3 is gebruik in elke eksperiment.(2)
[17]

Sub. vol

[Signature]

[Signature]

**QUESTION 6/VRAAG 6**

6.1

6.1.1 Remains the same/Bly dieselfde ✓ (1)

6.1.2 Decreases/Neem af ✓ (1)

6.1.3 Remains the same/Bly dieselfde ✓ (1)

- 6.2
- Decrease in pressure favours the reaction that produces a greater number of moles/amount of gas. ✓
'n Verlaging in druk bevoordeel die reaksie wat 'n groter aantal mol/hoeveelheid gas produseer.
 - Forward reaction is favoured. ✓ / [CO] increases AND [CO₂] decreases
Voorwaartse reaksie word bevoordeel./ [CO] neem toe EN [CO₂] neem af (2)

6.3

Marking criteria:(a) Substitute 44 in $n = \frac{m}{M}$ ✓(b) Change in mass of carbon:
 $m(C_i) - m(C_f)/n(C_i) - n(C_f)$ ✓(c) Substitute 12 in $n = \frac{m}{M}$ ✓

(d) Use mole ratio 1:1 ✓

(e) $n(\text{CO}_2)_{\text{eq}} = n(\text{CO}_2)_{\text{initial}} - n(\text{CO}_2)_{\text{used}}$ OR
 $m(\text{CO}_2)_{\text{eq}} = m(\text{CO}_2)_{\text{initial}} - m(\text{CO}_2)_{\text{used}}$ ✓(f) Final answer: 6,16 g ✓
RANGE: 6 to 6,16 g**NOTE:**If (b) $\Delta m(C)$ or $\Delta n(C)$ is not calculated
max $\frac{2}{6}$ **Nasienkriteria:**(a) Vervang 44 in $n = \frac{m}{M}$ ✓(b) Verandering in massa:
 $m(C_i) - m(C_f)/n(C_i) - n(C_f)$ ✓(c) Vervang 12 in $n = \frac{m}{M}$ ✓

(d) Gebruik molverhouding 1:1 ✓

(e) $n(\text{CO}_2)_{\text{ewe}} = n(\text{CO}_2)_{\text{begin}} - n(\text{CO}_2)_{\text{gebruik}}$ OR
 $m(\text{CO}_2)_{\text{ewe}} = m(\text{CO}_2)_{\text{begin}} - m(\text{CO}_2)_{\text{gebruik}}$ ✓(f) Finale antwoord: 6,16 g ✓
GEBIED: 6 tot 6,16 g**NOTA:**Indien (b) $\Delta m(C)$ of $\Delta n(C)$ nie bereken
maks $\frac{2}{6}$ **OPTION 1/OPSIE 1:**

$$\Delta m(C) = 14 - 4,44 \quad \checkmark \text{ (b)}$$

$$= 9,56 \text{ g}$$

$$n(\text{CO}_2)_{\text{initially}} = \frac{m}{M}$$

$$= \frac{41,2}{44} \quad \checkmark \text{ (a)}$$

$$= 0,94 \text{ mol (0,936)}$$

$$n(C)_{\text{used}} = \frac{m}{M}$$

$$n(C)_{\text{used}} = \frac{9,56}{12} \quad \checkmark \text{ (c)}$$

$$= 0,80 \text{ mol (0,797)}$$

$$n(\text{CO}_2)_{\text{used}} = n(C)$$

$$= 0,80 \text{ mol (0,797)} \quad \checkmark \text{ (d)}$$

$$n(\text{CO}_2)_{\text{eq}} = n(\text{CO}_2)_{\text{initially}} - n(\text{CO}_2)_{\text{used}}$$

$$= 0,94 - 0,80 \quad \checkmark \text{ (e)}$$

$$= 0,14 \text{ mol}$$

$$n(\text{CO}_2) = \frac{m}{M}$$

$$0,14 = \frac{m}{44}$$

$$X = m(\text{CO}_2) = 6,16 \text{ (g)} \quad \checkmark \text{ (f)}$$

OPTION 2/OPSIE 2:

$$\Delta m(C) = 14 - 4,44 \quad \checkmark \text{ (b)}$$

$$= 9,56 \text{ g}$$

$$n(C)_{\text{used}} = \frac{m}{M}$$

$$n(C)_{\text{used}} = \frac{9,56}{12} \quad \checkmark \text{ (c)}$$

$$= 0,80 \text{ mol (0,797)}$$

$$n(\text{CO}_2)_{\text{used}} = n(C)$$

$$= 0,80 \text{ mol (0,797)} \quad \checkmark \text{ (d)}$$

$$n(\text{CO}_2) = \frac{m}{M}$$


$$0,80 = \frac{m}{44} \quad \checkmark \text{ (a)}$$

$$m(\text{CO}_2) = 35,05 \text{ g}$$

$$m(\text{CO}_2)_{\text{eq}} = m(\text{CO}_2)_{\text{initially}} - m(\text{CO}_2)_{\text{used}}$$

$$= 41,2 - 35,05 \quad \checkmark \text{ (e)}$$

$$X = 6,15 \text{ (g)} \quad \checkmark \text{ (f)}$$



**OPTION 3/OPSIE 3:**


$$\begin{aligned}
 n(\text{CO}_2)_{\text{initially}} &= \frac{m}{M} \\
 &= \frac{41,2}{44} \checkmark \text{ (a)} \\
 &= 0,94 \text{ mol (0,936)}
 \end{aligned}$$

$$\begin{aligned}
 n(\text{C})_{\text{used}} &= \frac{9,56}{12} \checkmark \text{ (c)} \\
 &= 0,80 \text{ mol (0,797)}
 \end{aligned}$$

	C	CO ₂
Ratio/Verhouding	1	1
Initial quantity (mol) Aanvangshoeveelheid (mol)	1,17	0,936
Change (mol) Verandering (mol)	0,8	0,8 $\checkmark$ (d)
Quantity at equilibrium (mol)/ Hoeveelheid by ewewig (mol)	0,37 $\checkmark$ (b)	0,14 $\checkmark$ (e)

$$\begin{aligned}
 n(\text{CO}_2) &= 0,139 \text{ mol} \\
 m(\text{CO}_2) &= 0,139 (44) \\
 X &= 6,16 \text{ (g)} \checkmark \text{ (f)}
 \end{aligned}$$

(6)



SA EXAM PAPERS

Copyright reserved/Kopiereg voorbehou

Proudly South African

Please turn over/Blaai om asseblief



6.4

POSITIVE MARKING FROM QUESTION 6.3:**POSITIEWE NASIEN VANAF VRAAG 6.3:****Marking criteria**

- (a) Use of ratio $n(\text{CO}_2) : n(\text{CO}) = 1:2$. ✓
 (b) Divide by 3 dm^3 . ✓
 (c) Correct K_c expression (formulae in square brackets). ✓
 (d) Substitute of concentration into K_c expression. ✓
 (e) Final answer: 5,98 ✓
 RANGE: 5,98 – 7,29

Nasienkriteria:

- (a) Gebruik verhouding $n(\text{CO}_2) : n(\text{CO}) = 1:2$. ✓
 (b) Deel deur 3 dm^3 . ✓
 (c) Korrekte K_c uitdrukking (formules in vierkantige hakies). ✓
 (d) Vervang konsentrasies in korrekte K_c uitdrukking. ✓
 (e) Finale antwoord: 5,98 ✓
 GEBIED: 5,98 – 7,29

NOTE/NOTA:

Mark calculations of this question that may be done in QUESTION 6.3.

Merk berekeninge van hierdie vraag wat in VRAAG 6.3 gedoen is.

CALCULATIONS USING NUMBER OF MOLES**BEREKENINGE WAT AANTAL MOL GEBRUIK****OPTION 1/OPSIE 1:**

$$n(\text{CO}_2)_{\text{initial}} = \frac{m}{M}$$

$$= \frac{41,2}{44}$$

$$= 0,936 \text{ mol}$$

	CO_2	CO
Ratio/Verhouding	1	2
Initial quantity (mol) Aanvangshoeveelheid (mol)	0,936	0
Change (mol) Verandering (mol)	0,8	1,6
Quantity at equilibrium (mol)/ Hoeveelheid by ewewig (mol)	0,14	1,6
Equilibrium concentration ($\text{mol} \cdot \text{dm}^{-3}$) Ewewigskonsentrasie ($\text{mol} \cdot \text{dm}^{-3}$)	0,047	0,53


✓ (a)

Divide by/deel deur 3 ✓ (b)

$$K_c = \frac{[\text{CO}]^2}{[\text{CO}_2]} \quad \checkmark \text{ (c)}$$

$$= \frac{(0,53)^2}{0,047} \quad \checkmark \text{ (d)}$$

$$= 5,98 \quad \checkmark \text{ (e)}$$

Wrong K_c expressionVerkeerde K_c -uitdrukking: Max./Maks. $\frac{2}{5}$ No K_c expression/Geen K_c - uitdrukking: $\frac{4}{5}$ 

Sub. ✓
 ✓
 A


CALCULATIONS USING CONCENTRATION
BEREKENINGE WAT KONSENTRASIE GEBRUIK
OPTION 2/OPSIE 2:

$$c(\text{CO}_2) = \frac{m}{MV} = \frac{41,2}{(44)(3)} = 0,31 \text{ mol}\cdot\text{dm}^{-3}$$

Divide by 3 ✓ (b)

$$c = \frac{n}{V} = \frac{0,8}{3} = 0,27 \text{ mol}\cdot\text{dm}^{-3} (0,267)$$


	CO ₂	CO
Ratio/Verhouding	1	2
Initial concentration (mol·dm ⁻³) Aanvangskonsentrasie (mol·dm ⁻³)	0,31	0
Change in concentration (mol·dm ⁻³) Verandering in konsentrasie (mol·dm ⁻³)	0,27	0,54
Equilibrium concentration (mol·dm ⁻³) Ewewigskonsentrasie (mol·dm ⁻³)	0,04	0,54

✓ (a)

$$K_c = \frac{[\text{CO}]^2}{[\text{CO}_2]} \quad \checkmark (c)$$

$$= \frac{(0,54)^2}{0,04} \quad \checkmark (d)$$

$$= 7,29 \quad \checkmark (e)$$

Wrong K_c expressionVerkeerde K_c-uitdrukking: Max./Maks. 2/5No K_c expression/Geen K_c- uitdrukking: 4/5

Sub. Vch

**OPTION 3/OPSIE 3:**

$$n(\text{CO}_2)_{\text{initial}} = \frac{m}{M}$$

$$= \frac{41,2}{44}$$

$$= 0,936 \text{ mol}$$

$$\Delta n(\text{CO}_2) = 0,8 \text{ mol}$$

$$n(\text{CO}_2)_{\text{eqm}} = n(\text{CO}_2)_{\text{initial}} - \Delta n(\text{CO}_2)$$

$$= 0,936 - 0,8$$

$$= 0,136 \text{ mol}$$

$$n(\text{CO})_{\text{formed}} = 2\Delta n(\text{CO}_2)_{\text{used}} = 1,6 \text{ mol} \quad \checkmark \text{ (a)}$$

$$n(\text{CO})_{\text{eqm}} = \Delta n(\text{CO})_{\text{formed}} = 1,6 \text{ mol}$$

$$[\text{CO}_2]_{\text{eqm}} = \frac{0,136}{3} = 4,53 \times 10^{-2} \text{ mol} \cdot \text{dm}^{-3}$$

$$[\text{CO}]_{\text{eqm}} = \frac{1,6}{3} = 0,53 \text{ mol} \cdot \text{dm}^{-3}$$

 $\checkmark \text{ (b)}$

$$K_c = \frac{[\text{CO}]^2}{[\text{CO}_2]} \quad \checkmark \text{ (c)}$$

$$= \frac{(0,53)^2}{4,53 \times 10^{-2}} \quad \checkmark \text{ (d)}$$

$$= 6,2 \quad \checkmark \text{ (e)}$$

Wrong K_c expressionVerkeerde K_c -uitdrukking: Max./Maks. $\frac{2}{5}$ No K_c expression/Geen K_c -uitdrukking: $\frac{4}{5}$

(5)

6.5 Y ✓✓

(2)

6.6 Remains the same/Bly dieselfde ✓

(1)

[19]

DEPARTMENT OF BASIC EDUCATION
PRIVATE BAG X895, PRETORIA 0001
0.1 -06- 2025
APPROVED MARKING GUIDELINE
PUBLIC EXAMINATION



Vnl

**QUESTION 7/VRAAG 7**

7.1 An acid produces hydrogen ions /H⁺/hydronium ions/ H₃O⁺ in aqueous solution/water. ✓✓ (2 or 0)
'n Suur is 'n stof wat waterstofione/H⁺/hidroniumione/H₃O⁺ vorm in waterige oplossing/water. (2 of 0) (2)

7.2 (COOH)₂ ✓ (1)

7.2.1 NaCl ✓ (1)


7.2.2 HCO₃⁻ ✓
OR/OF NH₃ (1)

7.2.3 NaOH ✓✓
OR/OF Mg(OH)₂ (2)

7.3

Marking criteria: (a) Calculate n(H_xY) ✓ (b) Calculate n(NaOH) ✓ (c) Final answer: x = 2 ✓ (d) Reactants ✓ Products ✓ Balancing ✓ NOTE: Ignore ⇌ and phases Marking rule 6.3.10	Nasienkriteria: (a) Bereken n(H_xY) ✓ (b) Bereken n(NaOH) ✓ (c) Finale antwoord: x = 2 ✓ (d) Reaktanse ✓ Produkte ✓ Balansering ✓ NOTA: Ignoreer ⇌ en fases Nasienreël 6.3.10
OPTION 1/OPSIE 1: n = cV n_{acid} = (0,11)(0,02364) ✓ (a) = 2,6 x 10⁻³ n_{base} = (0,26)(0,02) ✓ (b) = 5,2 x 10⁻³ (0,0052) $\frac{n(\text{H}_x\text{Y})}{n(\text{NaOH})} = \frac{n_a}{n_b}$ $\frac{2,6 \times 10^{-3}}{5,2 \times 10^{-3}} = \frac{1}{n_b}$ n_b = 2 ∴ x = 2 ✓ (c)	OPTION 2/OPSIE 2: $\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$ (a) ✓ $\frac{(23,64)(0,11)}{(20)(0,26)} = \frac{1}{n_b}$ (b) ✓ $n_b = 2$ ∴ x = 2 ✓ (c) H_xY : NaOH 2,6 x 10⁻³ : 5,2 x 10⁻³ 1 : 2
H₂Y(aq) + 2NaOH(aq) ✓ → Na₂Y (aq) + 2H₂O(l) ✓ Bal ✓ (d)	

(6)



Handwritten signatures and initials.



7.4


Marking criteria:

- (a) Any formula: $pH = -\log[H_3O^+]/pH = -\log[H^+]/[H_3O^+] = 10^{-pH}$ ✓
 (b) Substitute 1,61 in $pH = -\log[H_3O^+]$ ✓
 (c) Calculate $n(HCl)_{\text{unused}}$ using $c = \frac{n}{V}$ ✓
 (d) Calculate $n(HCl)_{\text{initial}}$ using $c = \frac{n}{V}$ ✓
 (e) Calculate $n(HCl)_{\text{used}} = n(HCl)_{\text{initial}} - n(HCl)_{\text{unused}}$ ✓
 (f) Using ratio 1:2 with **USED** HCl from of (e) to calculate $n(CaCO_3)$ ✓
 (g) Substitute 100 AND $n(CaCO_3)$ from (f) in $n = \frac{m}{M}$ ✓
 (h) Mass of impurity = $m_{\text{sample}} - m(CaCO_3)$ ✓
 (i) Final answer: 0,25 g ✓ (Range: 0,2 g to 0,3 g)

Nasienkriteria:

- (a) Enige formule: $pH = -\log[H_3O^+]/pH = -\log[H^+]/[H_3O^+] = 10^{-pH}$ ✓
 (b) Vervang 1,61 in $pH = -\log[H_3O^+]$ ✓
 (c) Bereken $n(HCl)_{\text{ongebruik}}$ using $c = \frac{n}{V}$ ✓
 (d) Bereken $n(HCl)_{\text{begin}}$ using $c = \frac{n}{V}$ ✓
 (e) Bereken $n(HCl)_{\text{gebruik}} = n(HCl)_{\text{begin}} - n(HCl)_{\text{ongebruik}}$ ✓
 (f) Gebruik ratio 1:2 van HCl **GEBRUIK** van (e) om $n(CaCO_3)$ te bereken ✓
 (g) Vervang 100 EN $n(CaCO_3)$ van (f) in $n = \frac{m}{M}$ ✓
 (h) Massa of onsuiverheid = $m_{\text{monster}} - m(CaCO_3)$ ✓
 (i) Finale antwoord: 0,25 g ✓ (Gebied: 0,2 g tot 0,3 g)

$pH = -\log[H_3O^+] \quad \checkmark (a)$
 (b) $\checkmark 1,61 = -\log[H_3O^+]$
 $[H_3O^+] = 10^{-1,61}$
 $= 2,45 \times 10^{-2} \text{ mol} \cdot \text{dm}^{-3} (0,0245)$
 $n(HCl)_{\text{unused}} = n(H_3O^+) = cV$
 $= (2,45 \times 10^{-2})(0,2) \quad \checkmark (c)$
 $= 4,9 \times 10^{-3} \text{ mol} (0,0049)$
 $n(HCl)_{\text{initial}} = cV$
 $= (0,15)(0,2) \quad \checkmark (d)$
 $= 3 \times 10^{-2} \text{ mol} (0,03)$
 $n(HCl)_{\text{used}} = 3 \times 10^{-2} - 4,9 \times 10^{-3} \quad \checkmark (e)$
 $= 2,51 \times 10^{-2} \text{ mol} (0,0251)$
 Reaction ratio $nCaCO_3 : nHCl = 1:2$
 $n(CaCO_3) = \frac{1}{2}(2,51 \times 10^{-2}) = 1,25 \times 10^{-2} \text{ mol} \quad \checkmark (f)$
 $n(CaCO_3) = \frac{m}{M}$
 $1,25 \times 10^{-2} = \frac{m}{100} \quad \checkmark (g)$
 $m(CaCO_3) = 1,25 \text{ g}$
 $m \text{ of impurity in the sample} = 1,5 - 1,25 \quad \checkmark (h)$
 $= 0,25 \text{ g} \quad \checkmark (i)$

(9)
[22]

Sub.

Vwh

A

**QUESTION 8/VRAAG 8**

8.1 H^+/H_3O^+ ions/hydrogen ions/hydronium ions/oxonium ions ✓
Waterstofione/hidroniumione/oksoniumione (1)

8.2 0,77 V ✓ (1)

8.3 A ✓ (1)

8.4 H_2 is a stronger reducing agent ✓ than $Fe^{2+}/Fe(II)$ ions ✓ and will reduce $Fe^{3+}/Fe(III)$ ions ✓ (to $Fe^{2+}/Fe(II)$ ions).

H_2 is 'n sterker reduseermiddel as $Fe^{2+}/Fe(II)$ -ione en sal $Fe^{3+}/Fe(III)$ -ione reduseer (na $Fe^{2+}/Fe(II)$ -ione).

OR/OF





Fe^{2+} -ion is a weaker reducing agent ✓ than H_2 ✓ and therefore $Fe^{3+}/Fe(III)$ ions (to $Fe^{2+}/Fe(II)$ ions) will be reduced. ✓

Fe^{2+} -ioon is 'n sterker reduseermiddel as H_2 en sal $Fe^{3+}/Fe(III)$ -ione reduseer (na $Fe^{2+}/Fe(II)$ -ione). (3)

8.5

8.5.1 Pt/Platinum ✓ (1)

8.5.2 $H_2 \rightarrow 2H^+ + 2e^-$ ✓✓

NOTE/NOTA:


• Ignore if charge omitted on electron./Ignoreer indien lading weggelaat op elektron.

• If charge (+) omitted on H^+ /Indien lading (+) weggelaat op H^+ :


Example/Voorbeeld: $H_2 \rightarrow 2H + 2e^-$ Max/Maks: $\frac{1}{2}$

(2)

8.5.3



OR/OF



ACCEPT/AANVAAR:



(3)

8.6

The reaction reaches equilibrium/no charges/electrons flow. ✓
Die reaksie bereik ewewig/geen ladings/elektrone vloei.

(1)
[13]

DEPARTMENT OF BASIC
EDUCATION
PRIVATE BAG X8395, PRETORIA 0001
11-06-2025



SA EXAM PAPERS

Please turn over/Blaai om asseblief

Proudly South African

**QUESTION 9/VRAAG 9****9.1 ANY ONE:**


- The (chemical) process in which electrical energy is converted to chemical energy. ✓✓ **(2 or 0)**
- The use of electrical energy to produce a chemical change.
- Decomposition of an ionic compound by means of electrical energy.
- The process during which an electric current passes through a solution/ionic liquid/molten ionic compound.

ENIGE EEN:

- Die (chemiese) proses waarin elektriese energie omgeskakel word na chemiese energie. ✓✓ **(2 of 0)**
- Die gebruik van elektriese energie om 'n chemiese verandering teweeg te bring.
- Ontbinding van 'n ioniese verbinding met behulp van elektriese energie.
- Die proses waardeur 'n elektriese stroom deur 'n oplossing/ioniese vloeistof/gesmelte ioniese verbinding beweeg.

(2)**9.2** $2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$ ✓✓**NOTE/NOTA:**

- $\text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq}) \leftarrow 2\text{H}_2\text{O}(\text{l}) + 2\text{e}^-$ ($\frac{2}{2}$)
 - $2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$ ($\frac{1}{2}$)
 - $\text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq}) \rightleftharpoons 2\text{H}_2\text{O}(\text{l}) + 2\text{e}^-$ ($\frac{0}{2}$)
 - $2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \leftarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$ ($\frac{0}{2}$)
 - Ignore if charge omitted on electron. / Ignoreer indien lading weggelaat op elektron.
 - If charge (-) omitted on OH^- / Indien lading (-) weggelaat op OH^-
- Example/Voorbeeld: $2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}(\text{aq})$ ✓
- Max./Maks: $\frac{1}{2}$
- Ignore phases/Ignoreer fases

(2)**9.3** X to Y/X tot Y ✓**(1)**



9.4

Marking criteria:(a) Substitute 300×10^{-3} and 24 dm^3 into

$$n = \frac{V}{V_m} \checkmark$$

(b) Using ratio 1:2 to calculate $n(e^-)$ ✓(c) Substitute $6,02 \times 10^{23} \text{ mol}^{-1}$ in

$$n = \frac{N}{N_A} \checkmark$$

(d) Final correct answer:

$$1,505 \times 10^{22} \text{ electrons} \checkmark$$

Range: $1,505 \times 10^{22}$ to $2,41 \times 10^{22}$ electrons**Nasienkriteria:**(a) Vervang 300×10^{-3} en 24 dm^3 in

$$n = \frac{V}{V_m} \checkmark$$

(b) Gebruik verhouding 1:2 om $n(e^-)$ te bereken ✓(c) Vervang $6,02 \times 10^{23} \text{ mol}^{-1}$ in

$$n = \frac{N}{N_A} \checkmark$$

(d) Finale korrekte antwoord:

$$1,505 \times 10^{22} \text{ elektrone} \checkmark$$

Gebied: $1,505 \times 10^{22}$ tot $2,41 \times 10^{22}$ elektrone**OPTION 1/OPSIE 1:**

$$n(\text{Cl}_2) = \frac{V}{V_m}$$

$$= \frac{300 \times 10^{-3}}{24} \checkmark (a)$$

$$= 0,0125 \text{ mol (0,01)}$$

$$\begin{aligned} n(e^-) &= 2n(\text{Cl}_2) \downarrow \\ &= 2(0,0125) \checkmark (b) \\ &= 0,025 \text{ mol} \end{aligned}$$

$$\begin{aligned} n(e^-) &= \frac{N}{N_A} \downarrow \\ 0,025 &= \frac{N}{6,02 \times 10^{23}} \checkmark (c) \\ N &= 1,505 \times 10^{22} \text{ (electrons)} \checkmark (d) \end{aligned}$$

OPTION 2/OPSIE 2:

$$n(\text{Cl}_2) = \frac{V}{V_m}$$

$$= \frac{300 \times 10^{-3}}{24} \checkmark (a)$$

$$= 0,0125 \text{ mol (0,01)}$$

$$\begin{aligned} n(\text{Cl}_2) &= \frac{N}{N_A} \downarrow \\ 0,0125 &= \frac{N}{6,02 \times 10^{23}} \checkmark (c) \\ N &= 7,525 \times 10^{21} (\text{Cl}_2) \end{aligned}$$

$$\begin{aligned} N(e^-) &= 2n(\text{Cl}_2) \downarrow \\ &= 2(7,525 \times 10^{21}) \checkmark (b) \\ &= 1,505 \times 10^{22} \text{ (electrons)} \checkmark (d) \end{aligned}$$

TOTAL/TOTAAL:(4)
[9]
150

DEPARTMENT OF BASIC EDUCATION
PRIVATE BAG X895, PRETORIA 0001
01 -06- 2025
APPROVED MARKING GUIDELINE PUBLIC EXAMINATION



Sub. ✓
A. ✓




MARKING GUIDELINES

1. Calculations

- 1.1 **Marks will be awarded for:** correct formula, correct substitution, correct answer with unit.
- 1.2 **No marks** will be awarded if an **incorrect or inappropriate formula is used**, even though there may be relevant symbols and applicable substitutions.
- 1.3 When an error is made during **substitution into a correct formula**, a mark will be awarded for the correct formula and for the correct substitutions, but **no further marks** will be given.
- 1.4 If **no formula** is given, but **all substitutions are correct**, the candidate will **forfeit one mark**.
Example: No K_c expression, correct substitution:
- $$K_c = \frac{(2)^2}{(2)(1)^3} \checkmark = 2 \checkmark \quad \left(\frac{2}{3}\right)$$
- 1.5 Marks are only awarded for a formula if a **calculation has been attempted**, i.e. substitutions have been made or a numerical answer has been given.
- 1.6 Marks can only be allocated for substitutions when values are substituted into formulae and not when listed before a calculation starts.
- 1.7 The final answer to all calculations, when not specified in the question, must be rounded off to a minimum of TWO decimal places.
- 1.8 If a final answer to a calculation is correct, full marks will not automatically be awarded. Markers will always ensure that the correct/appropriate formula is used and that workings, including substitutions, are correct.
- 1.9 Mathematical manipulations and change of subject of appropriate formulae carry no marks, but if a candidate starts off with the correct formula and then changes the subject of the formula incorrectly, marks will be awarded for the formula and the correct substitutions. The mark for the incorrect numerical answer is forfeited.

Example:

CORRECT	ANSWER (1)	POSSIBLE	ANSWER (2)	POSSIBLE
$n = \frac{m}{M} \checkmark$ $0,01 \checkmark = \frac{m}{52} \checkmark$ $m = 0,52 \text{ g} \checkmark$ (4)	$n = \frac{m}{M} \checkmark$ $0,01 \checkmark = \frac{52}{m} \times$ $m = 5\,200 \text{ g} \times$ (2)	$m = \frac{n}{M} \times$ $= \frac{0,01}{52}$ $= 0,002 \text{ g}$ (0)	$n = \frac{m}{M} \checkmark$ $m = \frac{M}{n} \times$ $= \frac{52}{0,01} \checkmark$ $= 5\,200 \text{ g} \times$ (3)	$n = \frac{m}{M} \checkmark$ $= 0,52 \text{ g} \checkmark$ (2)



Sub. Vind
A.




2. Units

- 2.1 Candidates will only be penalised once for the repeated use of an incorrect unit **within a question**.
- 2.2 Units are only required in the final answer to a calculation.
- 2.3 Marks are only awarded for an answer and not for a unit per se. Candidates will therefore forfeit the mark allocated for the answer in each of the following situations:
- Correct answer + wrong unit
 - Wrong answer + correct unit
 - Correct answer + no unit
- 2.4 Separate compound units with a multiplication dot, not a full stop, for example $\text{mol}\cdot\text{dm}^{-3}$. Accept $\text{mol}\cdot\text{dm}^{-3}$ (or mol/dm^3) for marking purposes.





3. General

- 3.1 If one answer or calculation is required, but two are given by the candidate, only the first one will be marked, irrespective of which one is correct. If two answers are required, only the first two will be marked, etc.
- 3.2 When a chemical **FORMULA** is asked, and the **NAME** is given as answer, the candidate forfeits the marks. The same rule applies when the **NAME** is asked and the **FORMULA** is given.
- 3.3 When redox half-reactions are to be written, the correct arrow should be used.

If the equation



is the correct answer, the marks must be given as follows:



- 3.4 When candidates are required to give an explanation involving the relative strength of oxidising and reducing agents, do not accept the following:
- Stating the position of a substance on Table 4 only (e.g. Cu is above Mg).
 - Using relative reactivity only (e.g. Mg is more reactive than Cu).
 - The correct answer would be for instance: Mg is a stronger reducing agent than Cu, and therefore Mg will be able to reduce Cu^{2+} ions to Cu. The answer can also be given in terms of the relative strength as electron acceptors and donors.
- 3.5 One mark is forfeited when the charge of an ion is omitted per equation (not for the charge on an electron).
- 3.6 The error-carrying principle does not apply to chemical equations or half-reactions. For example, if a learner writes the wrong oxidation/reduction half-reaction in the sub-question and carries the answer over to another sub-question (balancing of equations or calculation of E_{cell}^0), then the learner must not be credited for this substitution.
- 3.7 In the structural formula of an organic molecule all hydrogen atoms must be shown. Marks must be deducted if hydrogen atoms are omitted.

DEPARTMENT OF BASIC EDUCATION
PRIVATE BAG X895, PRETORIA 0001
01 -06- 2025
APPROVED MARKING GUIDELINE
PUBLIC EXAMINATION




Handwritten signatures and initials:
 1. A signature starting with 'S'.
 2. Initials 'VF'.
 3. A signature starting with 'A'.



- 3.8 When a structural formula is asked, marks must be deducted if the learner writes the condensed formula.
- 3.9 When a IUPAC name is asked and the candidate omits the hyphen(s) (e.g. instead of pent-1-ene or 1-pentene the candidate writes pent 1 ene or 1 pentene), marks will be forfeited.
- 3.10 When a chemical reaction is asked, marks are awarded for correct reactants, correct products and correct balancing.
If only a reactant(s) followed by an arrow, or only a product(s) preceded by an arrow, is/are written, marks may be awarded for the reactant(s) or product(s). If only a reactant(s) or only a product(s) is/are written, without an arrow, no marks are awarded for the reactant(s) or product(s).

Examples:




4. Positive marking

Positive marking regarding calculations will be followed in the following cases:

- 4.1 **Sub-question to sub-question:** When a certain variable is calculated in one sub-question (e.g. QUESTION 3.1) and needs to be substituted in another (QUESTION 3.2 or QUESTION 3.3), e.g. if the answer for QUESTION 3.1 is incorrect and is substituted correctly in QUESTION 3.2 or QUESTION 3.3, **full marks** are to be awarded for the subsequent sub-questions.
- 4.2 **A multistep question in a sub-question:** If the candidate has to calculate, for example, current in the first step and gets it wrong due to a substitution error, the mark for the substitution and the final answer will be forfeited.

5. Negative marking

Normally an incorrect answer cannot be correctly motivated if based on a conceptual mistake. If the candidate is therefore required to motivate in QUESTION 3.2 the answer given to QUESTION 3.1, and QUESTION 3.1 is incorrect, no marks can be awarded for QUESTION 3.2. However, if the answer for, for example, QUESTION 3.1 is based on a calculation, the motivation for the incorrect answer in QUESTION 3.2 could be considered.



SA EXAM PAPERS

Proudly South African



86. ✓
A. ✓