

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



SA EXAM PAPERS

SA EXAM PAPERS
Proudly South African



FINAL



KWAZULU-NATAL PROVINCE

EDUCATION
REPUBLIC OF SOUTH AFRICA

GRADE 12

**NATIONAL
SENIOR CERTIFICATE**

PHYSICAL SCIENCES P2 (CHEMISTRY)

PREPARATORY EXAMINATION

SEPTEMBER 2025

MARKING GUIDELINES

MARKS: 150

This marking guideline consists of 11 pages.



QUESTION 1

- 1.1 B ✓✓ (2)
- 1.2 D ✓✓ (2)
- 1.3 C ✓✓ (2)
- 1.4 C ✓✓ (2)
- 1.5 C ✓✓ (2)
- 1.6 C ✓✓ (2)
- 1.7 D ✓✓ (2)
- 1.8 C ✓✓ (2)
- 1.9 D ✓✓ **DO NOT MARK** (2)
- 1.10 B ✓✓ (2)
- [20]**

QUESTION 2

- 2.1.1 4 – bromo – 5 – ethyl – 2 - methylheptane ✓✓✓

Marking criteria:

- correct stem i.e. heptane ✓
- substituents correctly identified i.e. bromo, ethyl, methyl ✓
- IUPAC name completely correct including numbering, sequence and hyphen ✓

(3)

- 2.1.2 3 – methylbut – 1 - ene ✓✓

Marking criteria:

- correct stem and substituents i.e. dibromo, methyl and heptane ✓
- IUPAC name completely correct including numbering, sequence and hyphen ✓

(2)

- 2.2 $C_nH_{2n}O_2$ ✓ (1)
- 2.3.1 E ✓ (1)
- 2.3.2 C ✓ (1)



- 2.4.1 Compounds with the same molecular formula, ✓ but different functional groups/homologous series. ✓

Marking criteria:

If any one of the underlined key words/phrases in the **correct context** is omitted, deduct 1 mark.

(2)

- 2.4.2 $C_5H_{10}O_2$ ✓ ✓

(2)

- 2.4.3 Pentanoic acid or methylbutanoic acid ✓ ✓

Marking criteria:

- correct homologous series: carboxylic acid. ✓
- IUPAC name completely correct. ✓

(2)

[14]

QUESTION 3

- 3.1 The temperature at which the vapour pressure of a substance equals atmospheric pressure. ✓ ✓

Marking criteria:

If any one of the underlined key words/phrases in the **correct context** is omitted, deduct 1 mark.

The underlined phrases must be in the correct context.

(2)

- 3.2.1 Ensure a fair test/comparison. ✓

(1)

- 3.2.2 London forces/Dispersion forces. ✓

(1)

- 3.2.3 Hydrogen bonding ✓

(1)

- 3.4 Y. ✓

Y has the lowest boiling point. ✓

Or Y has the lowest intermolecular forces compared to W and X.

(2)

- 3.5

Marking criteria:

- Relate boiling point with length of carbon chain/branching/number of side chains/surface area. ✓ ✓
- Compare the strength of the intermolecular forces. ✓
- Compare the energy required to overcome the intermolecular forces. ✓

X has the lowest boiling point ✓ and therefore has the shorter carbon chain/is branched/more compact/more spherical/smaller surface area over which the intermolecular forces act than W. ✓

Weaker intermolecular forces/Van der Waals forces/London forces than W. ✓

Less energy needed to overcome the intermolecular forces ✓ OR


(4)

W has the higher boiling point ✓ and therefore has the longer carbon chain/no branches/less compact/less spherical/largest surface area over which the intermolecular forces act than X. ✓

Stronger/more intermolecular forces/Van der Waals forces/London forces than X ✓

More energy needed to overcome the intermolecular forces ✓

3.6

**Marking criteria:**

- Functional group on first carbon ✓
- 3 carbons in the longest chain ✓
- 2 – methyl groups on the second carbon ✓

(3)

3.7

Marking criteria:

- LOWER THAN ✓
- Correctly identify intermolecular forces in both compounds. ✓
- Compare the strength of the intermolecular forces. ✓
- Compare the energy required to overcome the intermolecular forces. ✓

LOWER THAN ✓

2,2 – dimethylpropane has only London/Dispersion forces between molecules and Compound Y has hydrogen bonds ✓ and dipole-dipole forces in addition to London/Dispersion forces

Intermolecular forces between molecules of 2,2 – dimethylpropane are weaker ✓

Require less energy to overcome. ✓

(4)
[18]

**QUESTION 4**

4.1 addition✓ (1)

4.2 The C-atom bonded to the hydroxyl/-OH✓ is bonded to three other carbon atoms.✓

Marking criteria:


If any one of the underlined key words/phrases in the **correct context** is omitted, deduct 1 mark.

The underlined phrases must be in the correct context.

4.3.1 Water/H₂O✓ (2)

4.3.2 Sulphuric acid/H₂SO₄ or Phosphoric acid/H₃PO₄✓ (1)

4.4.1

**Marking criteria:**

- Correct functional group✓
- substituent correctly identified i.e. methyl✓
- IUPAC name completely correct including numbering, sequence and hyphen ✓

4.4.2 2 - methyl - 1 - bromopentane✓✓✓ (3)

Marking criteria:

- correct stem i.e. pentane ✓
- substituents correctly identified i.e. bromo, methyl✓
- IUPAC name completely correct including numbering, sequence and hyphen ✓

4.5.1 hydroxyl✓ (1)

4.5.2 substitution✓ (1)

4.5.3 Dilute KOH/potassium hydroxide✓ (1)





Marking criteria:

- Both reactants correct. ✓
- Both products correct ✓
- Balancing ✓

(3)
[17]

QUESTION 5

5.1 The CO_2 produced escapes from the container. ✓ (1)

- 5.2
- Change in concentration ✓ of products/reactants per (unit) time. ✓
 - Change in amount/number of moles/volume/mass ✓ of products/reactants per (unit) time. ✓
 - Amount/number of moles/volume/mass of products formed/reactants used per (unit) time. ✓✓

Marking criteria:

If any one of the underlined key words/phrases in the **correct context** is omitted, deduct 1 mark.

The underlined phrases must be in the correct context.

- Rate of change in concentration/amount/number of moles/volume/mass. ✓✓ (2 or 0) (2)
- 5.3 • Reaction is complete/no more CO_2 is produced/ CaCO_3 is used up ✓✓ (2)
- 5.4 • Crush the calcium carbonate chips to a powder. ✓
- Add a catalyst. ✓ (2)

5.5 **Marking criteria:**

- Correct substitution ($\frac{62,25}{100}$) in the formula $n = \frac{m}{M}$ to calculate $n(\text{CaCO}_3)$ ✓
- Ratio: $n(\text{CaCO}_3)$ used equals $n(\text{CO}_2)$ produced ✓
- Use $n = \frac{m}{M}$ to calculate $m(\text{CO}_2)$ ✓
- Substitute for both n and M correctly i.e.: $n = 0,6225$ and $M = 44$ ✓
- Rate formula: $\text{rate} = \frac{\Delta m}{\Delta t}$ ✓
- Correct substitution of 1,37 for rate ✓
- Correct substitution for Δm in the formula: $\text{rate} = \frac{\Delta m}{\Delta t}$ ✓
- Final answer $\Delta t = 19,99$ minutes ✓



$$\begin{aligned}
 n(\text{CO}_2) \text{ lost} &= n(\text{CaCO}_3) \text{ reacted} \quad \checkmark \\
 &= \frac{m}{M} \quad \checkmark \\
 &= \frac{62,25}{100} \quad \checkmark \\
 &= 0,6225 \text{ mols}
 \end{aligned}$$

$$\begin{aligned}
 m(\text{CO}_2) \text{ lost} &= nM \\
 &= (0,6225)(44) \quad \checkmark \\
 &= 27,39 \text{ g}
 \end{aligned}$$

$$\begin{aligned}
 \text{rate} &= \frac{\Delta m}{\Delta t} \quad \checkmark \\
 1,37 \checkmark &= \frac{27,39 \checkmark}{\Delta t} \\
 \Delta t &= 19,99 \text{ minutes} \checkmark
 \end{aligned}$$

(8)

5.6.1 INCREASES. ✓

(1)

5.6.2 REMAINS THE SAME ✓
CaCO₃ is the limiting reagent ✓

(2)

- 5.7
- More reacting molecules per unit volume. ✓
 - More molecules correctly orientated. ✓
 - More effective collisions per unit time/second. ✓
- OR
- Frequency of effective collisions increases.

(3)

[21]

QUESTION 66.1 Add more reactants/increase the amount/mols/concentration of the reactants. ✓
OR

Remove some of the products/ decrease the amount/mols/concentration of the products.

OR

Decrease in pressure

(1)



6.2

Marking criteria:

- Correctly calculate 70,59% of 5 = 3,53 ✓
- Using the correct mol ratio ✓
- Calculating the quantity(mol) at equilibrium of all three substances ✓
- Using the concentration of Q₂ at equilibrium and the number of mols of A₂ at equilibrium in the equation $c = n/V$ to calculate the volume of the container. ✓✓
- Calculating the equilibrium concentrations of the reactants ✓
- K_c expression ✓
- Correct substitution of equilibrium concentrations into K_c expression ✓
- K_c = 5,09 ✓

	A ₂ Q	A ₂	Q ₂	
Initial quantity (mol)	5	0	0	
Change (mol)	3,53 ✓	3,53	1,765	✓
Quantity at equilibrium (mol)	1,47	3,53	1,765	✓
Equilibrium concentration (mol.dm ⁻³)	0,735	1,765	0,8825	✓

$$c = \frac{n}{V} \quad \checkmark$$

$$0,8825 = \frac{1,765}{V} \quad \checkmark$$

$$V = 2 \text{ dm}^3$$

$$K_c = \frac{[A_2]^2[Q_2]}{[A_2Q]^2} \quad \checkmark$$

$$= \frac{(1,765)^2(0,8825)}{(0,735)^2} \quad \checkmark$$

$$= 5,09 \quad \checkmark$$

(9)

6.3 The rates of the forward and reverse reaction do not increase equally. ✓

(1)

6.4

Marking criteria:

If any one of the underlined key words/phrases in the correct context is omitted, deduct 1 mark.

The underlined phrase must be in the correct context.

When the equilibrium in a closed system is disturbed, the system will re-instate a new equilibrium by favouring the reaction that will oppose the disturbance. ✓✓

(2)



- 6.5 GREATER THAN ✓ (1)
- 6.6 The rate of the forward reaction decreases less than the rate of the reverse reaction. ✓
 The forward reaction is favoured. ✓
 Concentration of the products increases while the concentration of the reactants decreases. ✓ (3)
[17]

QUESTION 7

- 7.1.1 CH_3COO^- ✓ (1)
- 7.1.2 GREATER THAN 7 ✓ (1)
- 7.1.3 $\text{CH}_3\text{COO}^- + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{COOH} + \text{OH}^-$
 LHS ✓
 RHS ✓
 Balancing ✓ (3)
- 7.2.1 It is a substance that ionises completely ✓ in water to produce a high concentration of hydronium ions. ✓ (2)

7.2.2

Marking criteria:

- Calculate $n(\text{HCl})_{\text{excess}}$ ✓
- Substitute for c and V in $n = cV$ ✓
- Ratio $\text{HCl} : \text{Na}_2\text{CO}_3 = 2:1$ ✓
- Substitute for c and V ✓
- Calculate $n(\text{HCl})_{\text{total}}$ by adding above 2 values ✓
- Ratio of $c(\text{HCl}) = c(\text{H}_3\text{O}^+)$ ✓
- Substitute into $c = n/V$ to calculate concentration of H_3O^+ ✓
- Equation: $\text{pH} = -\log[\text{H}_3\text{O}^+]$ ✓
- Substitute into $\text{pH} = -\log[\text{H}_3\text{O}^+]$ ✓
- Final answer ✓






$$\begin{aligned}
 n(\text{HCl})_{\text{excess}} &= cV \quad \checkmark \\
 &= (0,013)(0,075) \quad \checkmark \\
 &= 0,000975 \text{ mols} \\
 n(\text{HCl})_{\text{reacted with Na}_2\text{CO}_3} &= 2cV \quad \checkmark \\
 &= (2)(0,075)(0,025) \quad \checkmark \\
 &= 0,00375 \text{ mols} \\
 n(\text{HCl})_{\text{total}} &= 0,000975 + 0,00375 \quad \checkmark \\
 &= 0,004725 \text{ mols} \\
 c(\text{H}_3\text{O}^+) &= c(\text{HCl}) \quad \checkmark \\
 &= \frac{n}{V} \\
 &= \frac{0,004725}{0,05} \quad \checkmark \\
 &= 0,0945 \text{ mol.dm}^{-3} \\
 \text{pH} &= -\log[\text{H}_3\text{O}^+] \quad \checkmark \\
 &= -\log 0,0945 \quad \checkmark \\
 &= 1,02 \quad \checkmark
 \end{aligned}$$

(10)

[17]

QUESTION 88.1 Pt $\checkmark$ (1)8.2 Remain the same. $\checkmark$ (1)8.3.1 $\text{H}_2 \rightarrow 2\text{H}^+ + 2\text{e}^-$ $\checkmark\checkmark$ **Notes**

- Ignore phases



Ignore if charge on electron omitted.

(2)

(1)

8.3.2 $25^\circ\text{C}/298 \text{ K}$ $\checkmark$ 8.3.3 1 mol.dm^{-3} $\checkmark$ (1)8.4 Cu^{2+} $\checkmark$ (1)

- 8.5 $\text{Pt(s)} / \text{H}_2(\text{g}) / \text{H}^+(\text{aq})(1 \text{ mol.dm}^{-3}) \checkmark // \checkmark \text{Cu}^{2+}(\text{aq})(1 \text{ mol.dm}^{-3}) / \text{Cu(s)} \checkmark$ (3)
Ignore the phases and concentrations

8.6

Notes

- Accept any other correct formula from the data sheet.

Any other formula using unconventional abbreviations, e.g. $E^\circ_{\text{cell}} = E^\circ_{\text{OA}} - E^\circ_{\text{RA}}$ followed by correct substitutions Max: $\frac{3}{4}$

$$E^\circ_{\text{cell}} = E^\circ_{\text{reduction}} - E^\circ_{\text{oxidation}} \checkmark$$

$$0,75 \checkmark = 0,34 \checkmark - E^\circ_{\text{oxidation}}$$

$$E^\circ_{\text{oxidation}} = -0,41 \text{ V} \checkmark$$

Yes $\checkmark$, Cr^{2+} is not a solid. $\checkmark$

(6)

[16]**QUESTION 9**

- 9.1 Electrolytic $\checkmark$
They both have a power supply $\checkmark$
Both cells convert electrical energy to chemical energy. (2)
- 9.2 An electrolyte is a substance of which the aqueous solution contains ions. $\checkmark\checkmark$
OR
A substance that dissolves in water to give a solution that conducts electricity. (2)
- 9.3 $2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2 + 2\text{OH}^- \checkmark\checkmark$
Ignore phases

Notes

- $2\text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{H}_2 + 2\text{OH}^-$ (1/2)
- $\text{H}_2 + 2\text{OH}^- \leftarrow 2\text{H}_2\text{O} + 2\text{e}^-$ (2/2)
- $\text{H}_2 + 2\text{OH}^- \rightleftharpoons 2\text{H}_2\text{O} + 2\text{e}^-$ (0/2)
- $2\text{H}_2\text{O} + 2\text{e}^- \leftarrow \text{H}_2 + 2\text{OH}^-$ (0/2)

(2)

- 9.4 Cu or Copper $\checkmark$ (1)

- 9.5 Decreases $\checkmark$

Cu^{2+} (or Copper(II) ions) are reduced $\checkmark$ to Cu $\checkmark$ (or Copper). (3)

TOTAL: **[10]**
150

NB: MARK SCRIPT OUT OF 148 AND THEREAFTER CONVERT TO TOTAL MARKS : 150

