

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great
Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



SA EXAM PAPERS

SA EXAM PAPERS

Proudly South African



LIMPOPO
PROVINCIAL GOVERNMENT
REPUBLIC OF SOUTH AFRICA

DEPARTMENT OF EDUCATION

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

PHYSICAL SCIENCES PAPER 2

MEMO

PREPARATORY 2025

MARKS: 150

TIME: 3 Hours

This Marking Guidelines consist of 10 pages including the cover page



SA EXAM PAPERS

Proudly South African

QUESTION 1

- | | | |
|------|-----|-----|
| 1.1 | C✓✓ | (2) |
| 1.2 | A✓✓ | (2) |
| 1.3 | B✓✓ | (2) |
| 1.4 | A✓✓ | (2) |
| 1.5 | B✓✓ | (2) |
| 1.6 | A✓✓ | (2) |
| 1.7 | A✓✓ | (2) |
| 1.8 | B✓✓ | (2) |
| 1.9 | B✓✓ | (2) |
| 1.10 | D✓✓ | (2) |

[20]**QUESTION 2**

- 2.1 Chain isomers are compounds with the same molecular formula but different types of chains. ✓✓ (2)
- 2.2.1 C✓ (1)
- 2.2.2 4 – bromo – 2,2 – dimethylpentane✓✓✓

Notes:

1. Bromo, dimethyl identified✓
2. Pentane✓
3. Whole structure correct✓

(3)

- 2.2.3 Pent – 1 – ene✓✓

Notes:


1. Correct functional group✓
2. Whole structure correct✓

(2)

- 2.2.4 C_nH_{2n} ✓

(1)

2.3

**Notes:**

1. Correct functional group✓
2. Whole structure correct✓

(2)

- 2.4 $C_5H_{12} + 8O_2 \rightarrow 5CO_2 + 6H_2O$ (✓balance) (3)

- 2.5 Aldehydes✓ (1)





2.6 $20\% \text{ C} - \text{atoms in comp. C} = \frac{20}{100} \times 5$
 $= 1$

Compound E has 6 C – atoms

$\therefore x = 6$ and $y = 14$ **OR** ($x = 6$ and $y = 14$)

(4)

[19]**QUESTION 3**

3.1 Melting point is the temperature at which the solid and the liquid phases of a substance are at equilibrium. ✓✓ (2)

3.2 As the chain length/molar mass/number of carbon atoms increases, the boiling/melting point increases. ✓✓ (2)

3.3 Stronger than. ✓ (1)

3.4 Fair. ✓ The compounds belong to the same homologous/intermolecular force/functional group/there is only one independent variable. ✓ (2)

3.5 Gas. ✓ (1)

- 3.6
- Propane has a longer chain length/larger/ surface area/ bigger molecular size/more carbon atoms than ethane. ✓
 - Both have London forces/dispersion/dipole-dipole which are stronger between propane molecules than in ethane. ✓/the intermolecular forces of propane are stronger
 - More energy is needed to overcome the intermolecular force in propane than in ethane. ✓

OR


- Ethane has shorter chain length/smaller surface area/smaller molecular size/less carbon atoms than propane. ✓
- Both have London forces/dispersion/dipole-dipole which are weaker between ethane molecules than in propane. ✓
- Less energy is needed to overcome the intermolecular forces in ethane than in propane. ✓

[11]

QUESTION 4

4.1.1 Substitution/hydrolysis✓ (1)

4.1.2 Butan – 2 – ol✓✓ Accept 2-butanol (1 mark for butanol, 1 mark for position) (2)


**Notes:**

1. Correct functional group✓
2. Whole structure correct✓

(2)

4.2.2 (Strong) Heat✓ (1)

4.3.1



(5)

4.3.2 **Positive marking form 4.3.1**

But-2-ene.✓✓ /Accept 2-Butene (1 mark for butene, 1 mark for position)

(2)

4.4.1 Addition/hydrogenation.✓ (1)

4.4.2 Ni/Pt/Pd/Nickel/Platinum/Palladium.✓ (1)

4.5.1 Cracking is the chemical process in which longer chain hydrocarbon molecules are broken down to shorter more useful molecules.✓✓ (2)

4.5.2 Catalyst (Pt, Pd and Ni).✓ (1)

[18]

QUESTION 5

5.1 Rate of reaction is the change in concentration of reactants or products per unit time. ✓✓ (2)

5.2 Neutralisation reaction. ✓/Acid-base reaction (1)

5.3 State of division/surface area. ✓ (1)

5.4 Rate of reaction. ✓ (1)

$$\begin{aligned} 5.5 \quad n(\text{CO}_2) &= \frac{v}{v_m} \checkmark \\ &= \frac{53}{22\,400} \checkmark \\ &= 0,002366 \text{ mol} \end{aligned}$$

$$\begin{aligned} n(\text{HCl}) &= \frac{2(0,002366)}{22\,400} \checkmark \\ &= 0,004732 \text{ mol} \end{aligned}$$

$$\begin{aligned} \text{Rate} &= \frac{\Delta n}{\Delta t} \checkmark \\ &= \frac{0,004732}{60} \checkmark \\ &= 0,0000789 \text{ mol} \cdot \text{s}^{-1} / (7,89 \times 10^{-5} \text{ mol} \cdot \text{s}^{-1}) \checkmark \end{aligned} \quad (5)$$

[10]

QUESTION 6

- 6.1.1 Chemical equilibrium is a dynamic equilibrium when the rate of the forward reaction equals to the rate of the reverse reaction. ✓✓ (2)

6.1.2

	SO ₂	O ₂	SO ₃	
Ratio	2	1	2	
Initial amount(mol)	x	0,5	0	
Change	-0,75	-0,375✓	0,75	✓Ratio
Equilibrium	x - 0,75	0,125✓	0,75	
Concentration	$\frac{x - 0,75}{2}$	0,0625	0,375	✓ ÷ 2

$$K_c = \frac{[SO_3]^2}{[SO_2]^2 [O_2]} \checkmark$$

$$10 \checkmark = \frac{(0,375)^2}{\left(\frac{x-0,75}{2}\right)^2 (0,0625)} \checkmark$$

$$x = 1,70 \text{ mol} \checkmark$$

(8)

- 6.1.3 **Positive marking from 6.1.2**

$$n = \frac{m}{M} \checkmark$$

$$1,70 = \frac{m}{64} \checkmark$$

$$\therefore m = 108,8g \checkmark$$

(3)

- 6.2.1 Decreases. ✓

(1)

- 6.2.2 Stays the same. ✓

(1)

- 6.2.3 Decreases. ✓ A decrease in pressure favours the reaction that produces more moles of gas/more volume. ✓ The reverse reaction is favoured. ✓

(3)

[18]

QUESTION 7

7.1 A weak base is a base that ionises/dissociates incompletely in water to form a low concentration of hydroxide ions.✓✓ (2)

7.2 **Option 1:**

$$[\text{H}_3\text{O}^+] = 2 \times 0,02 = 0,04 \text{ mol.dm}^{-3}$$

$$K_w = [\text{H}_3\text{O}^+][\text{OH}^-]$$

$$1 \times 10^{-14} = 0,04 [\text{OH}^-] \quad [\text{OH}^-] = 2,5 \times 10^{-13} \text{ mol.dm}^{-3}$$

Option 2:

$$\text{pH} = -\log[\text{H}_3\text{O}^+]$$

$$= -\log(0,04)$$

$$= 1,398$$

$$\text{pH} + \text{pOH} = 14$$

$$1,398 + \text{pOH} = 14$$

$$\text{pOH} = 12,602$$

$$\text{pOH} = -\log[\text{OH}^-]$$

$$12,602 = -\log[\text{OH}^-]$$

$$[\text{OH}^-] = 2,50 \times 10^{-13} \text{ mol.dm}^{-3}$$

(4)

7.3.1 $n(\text{H}_2\text{SO}_4) = cv$ ✓

$$= (0,02)(0,025)$$
✓

$$= 0,0005 \text{ mol}$$

$$n(\text{NH}_3) = 2(0,0005)$$
✓

$$= 0,001 \text{ mol}$$

$$[\text{NH}_3] \text{ in diluted solution} = \frac{c}{V}$$

$$= \frac{0,001}{0,0312}$$
✓

$$= 0,032051 \text{ mol}$$

$$n(\text{NH}_3) \text{ in } 250 \text{ cm}^3 = cv$$

$$= 0,032051 \times 0,25$$
✓

$$= 0,0080128 \text{ mol}$$

$$n(\text{NH}_3) \text{ in } 250 \text{ cm}^3 = n(\text{NH}_3) \text{ in } 20 \text{ cm}^3 \text{ of undiluted solution}$$

$$[\text{NH}_3] = \frac{n}{V}$$

$$= \frac{0,0080128}{0,02}$$
✓

$$= 0,4 \text{ mol} \cdot \text{dm}^{-3}$$
✓

(7)

7.3.2 Below 7✓



(4)

7.4.1 An Ampholyte is a substance that can act as either an acid or a base✓✓

(2)

7.4.2 HC_2O_4^- ✓

(1)

7.4.3 $\text{H}_2\text{C}_2\text{O}_4$ ✓ and H_3O^+ ✓

(2)

[22]

QUESTION 8

- 8.1 A galvanic cell is a cell in which chemical energy is converted to electrical energy.✓✓ (2)
- 8.2 Concentration of $1\text{ mol}\cdot\text{dm}^{-3}$ ✓
Temperature of $25^{\circ}\text{C}/298\text{K}$ ✓ (2)
- 8.3.1 Pt✓ (1)
- 8.3.2 Cr✓ (1)
- 8.4 $2\text{ mol}\cdot\text{dm}^{-3}$ ✓ (2)
- 8.5 $3\text{Sn}^{4+} + 2\text{Cr} \longrightarrow 3\text{Sn}^{2+} + 2\text{Cr}^{3+}$ ✓ (✓balancing) (3)
- 8.6 $E^{\theta}_{\text{cell}} = E^{\theta}_{\text{cathode}} - E^{\theta}_{\text{anode}}$ ✓
 $= +0,15 - (-0,74)$ ✓
 $= +0,89\text{V}$ ✓ (4)
- 8.7 Decreases.✓ (1)

[16]





QUESTION 9

9.1 Electrolysis is the chemical process in which electrical energy is converted to chemical energy.✓✓ (2)

9.2 $\text{Cu} \longrightarrow \text{Cu}^{2+} + 2\text{e}^-$ ✓✓

Notes/Notas:

Ignore phases/ignoreer fases



(2)

9.3 Impure copper.✓ (1)

9.4.1 Zinc/Nickel/Zn/Ni.✓ (1)

9.4.2 Zinc/Nickel is a stronger reducing agent✓ than copper.✓ (2)

9.4.3 They will fall to the bottom.✓ (1)

9.5 $\text{mass of copper} + \text{impurities} = 600 - 390$ ✓
 $= 210\text{g}$

$\text{mass of pure copper} = 320 - 150$ ✓
 $= 170\text{g}$

$$\% \text{ purity} = \frac{170}{210} \times 100$$
✓

$$= 80,95\%$$
✓

(4)

9.6 It remains constant.✓ The rate of oxidation of copper is equal to the rate of reduction of copper ions.✓✓ (3)

[16]**TOTAL: 150**