

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



SA EXAM PAPERS

SA EXAM PAPERS
Proudly South African



This Paper was downloaded from SAEXAMPAPERS



Province of the
EASTERN CAPE
EDUCATION

Iphondo leMpuma Kapa: Isebe leMfundo
Provinsie van die Oos Kaap: Departement van Onderwys
Porafensio Ya Kapa Botjhabela: Lefapha la Thuto

NATIONAL SENIOR CERTIFICATE

GRADE 12

SEPTEMBER 2025

PHYSICAL SCIENCES P2: (CHEMISTRY)

MARKS: 150

TIME: 3 hours

This question paper consists of 20 pages, including 4 data sheets.



SA EXAM PAPERS

Proudly South African

INSTRUCTIONS AND INFORMATION

1. Write your name and surname in the appropriate space on the ANSWER BOOK.
2. This question paper consists of NINE questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two sub questions, e.g. between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. You may use appropriate mathematical instruments.
8. Show ALL formulae and substitutions in ALL calculations.
9. Round off your FINAL numerical answers to a minimum of TWO decimal places.
10. Give brief motivations, discussions, etc. where required.
11. You are advised to use the attached DATA SHEETS.
12. Write neatly and legibly.

QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Choose the answer and write only the letter (A–D) next to the question numbers (1.1 to 1.10) in the ANSWER BOOK, for example 1.11 D.

- 1.1 Which ONE of the following homologous series can be described by the general formula $C_nH_{2n}O$?
- A Alcohols
- B Aldehydes
- C Esters
- D Carboxylic acids (2)
- 1.2 The empirical formula of methyl propanoate is ...
- A $C_4H_8O_2$
- B C_4H_8O
- C C_2H_4O
- D CH_2O (2)
- 1.3 Consider the following compounds from the SAME HOMOLOGOUS SERIES with different boiling points.


Compounds	Boiling point (°C)
C_4H_8O	63
C_4H_8O	75

Which ONE of the following combinations is CORRECT regarding the homologous series to which compounds belong and type of isomers they form?

	Homologous series	Type of isomers
A	Ketones	Positional isomers
B	Ketones	Chain isomers
C	Aldehydes	Positional isomers
D	Aldehydes	Chain isomers


(2)

- 1.4 The organic molecule shown below undergoes thermal cracking.



Which ONE of the following are possible products from the reaction?


- A Hexane and butane
 - B Octane, ethane and hydrogen gas
 - C Pentane, propene and ethene
 - D Propane, butane and ethene (2)
- 1.5 Which ONE of the following is TRUE regarding an EXOTHERMIC REACTION?
- A No energy is absorbed or released.
 - B The amount of energy released is greater than the amount of energy absorbed.
 - C The amount of energy absorbed is greater than the amount of energy released.
 - D The amount of energy absorbed is equal to the amount of energy released. (2)
- 1.6 Consider the reaction below that reaches chemical equilibrium in a closed container.



Which ONE of the following changes will increase the yield of NH_3 ?

- A Addition of a catalyst
- B Increase temperature
- C Decrease in the volume of the container
- D Decrease in the concentration of N_2 (2)

1.7 The diagrams shows a titration set-up and titration curve obtained.



Consider the statements regarding substance **X**.

- I** The K_a value for substance **X** is greater than 1.
- II** The amount of **X** is equal the amount of NaOH at the equivalence point.
- III** Substance **X** ionises in water to produce low concentration of H_3O^+

Which of the above statement(s) is/are TRUE?

- A **I** and **II** only
- B **II** only
- C **II** and **III** only
- D **I** and **III** only (2)

1.8 The reaction represented by the balanced equation below reaches equilibrium in a closed container.



Which ONE of the following changes will favour the reverse reaction?

- A Addition of hydrogen (H_2)
- B Addition of potassium hydroxide (KOH)
- C Decrease the pH of the equilibrium mixture
- D Cool the equilibrium mixture (2)


- 1.9 The standard hydrogen half-cell ($\text{H}^+ | \text{H}_2 | \text{Pt}$) is connected to $\text{Mn} | \text{Mn}^{2+}$ half-cell to form a galvanic cell.

Which ONE of the following combinations CORRECTLY identifies the oxidising agent and direction of electron flow?

	Oxidising agent	Direction of flow of electrons	
		From half-cell	To half-cell
A	Mn	$\text{H}^+ \text{H}_2 \text{Pt}$	$\text{Mn} \text{Mn}^{2+}$
B	Mn^{2+}	$\text{Mn} \text{Mn}^{2+}$	$\text{H}^+ \text{H}_2 \text{Pt}$
C	H^+	$\text{H}^+ \text{H}_2 \text{Pt}$	$\text{Mn} \text{Mn}^{2+}$
D	H^+	$\text{Mn} \text{Mn}^{2+}$	$\text{H}^+ \text{H}_2 \text{Pt}$

(2)

- 1.10 The electrolytic cell below is used for the electrolysis of concentrated sodium chloride (NaCl).



Consider the statements regarding the electrolysis of concentrated sodium chloride (NaCl).

- I** The pH of the electrolyte increases over time.
- II** H_2O is oxidised at electrode **P**.
- III** Chlorine gas (Cl_2) is produced at electrode **Q**.

Which of the above statement(s) is/are TRUE?

- A **I** and **II** only
- B **III** only
- C **II** and **III** only
- D **I** and **III** only

(2)

[20]


QUESTION 2 (Start on a NEW page.)

The table below shows organic molecules (**A–F**) from different homologous series.

A $ \begin{array}{ccccccc} & & & & \text{CH}_3 & & \\ & & & & & & \\ \text{CH}_2 & - & \text{CH} & - & \text{CH}_2 & - & \text{CH} \\ & & & & & & \\ \text{CH}_3 & & \text{Cl} & & & & \text{CH}_3 \end{array} $	B 3,4-dimethylpentan-2-ol
C $\text{CH}_3\text{CH}(\text{CH}_3)\text{CCCH}_3$	D $ \begin{array}{ccccccc} & \text{H} & & \text{H} & & \text{H} & & \text{O} \\ & & & & & & & \\ \text{H} & - \text{C} & - & \text{C} & - & \text{C} & - & \text{C} - \text{H} \\ & & & & & & & \\ & \text{H} & & & & \text{H} & & \\ & & & & & & & \\ & & & \text{H} - \text{C} - \text{H} & & & & \\ & & & & & & & \\ & & & \text{H} & & & & \end{array} $
E 3-methylbutanone	F C_xH_y

- 2.1 Define *homologous series*. (2)
- 2.2 Write down the LETTER that represents EACH of the following:
- 2.2.1 Unsaturated hydrocarbon (1)
- 2.2.2 A compound with a carbonyl group bonded to two saturated carbon atoms (1)
- 2.2.3 Compounds that are functional isomers (2)
- 2.3 Write down the IUPAC name of:
- 2.3.1 Compound **A** (3)
- 2.3.2 Compound **C** (2)
- 2.4 Draw the STRUCTURAL FORMULA of compound **B**. (2)
- 2.5 Compound **D** has a positional isomer.
- 2.5.1 Define the term *positional isomer*. (2)
- 2.5.2 Draw the positional isomer of compound **D**. (2)

- 2.6 Compound **F** (C_xH_y) is an alkane that undergoes complete combustion with excess oxygen as shown below.




- 2.6.1 Determine the value **P**. (2)

- 2.6.2 Write down the CONDENSED STRUCTURAL FORMULA of compound **F**. (2)

[21]

QUESTION 3 (Start on a NEW page.)

The vapour pressure against temperature curves are drawn for THREE STRAIGHT CHAIN ALKANES. The vapour pressure are measured in mmHg. The atmospheric pressure is 760 mmHg.

GRAPH OF VAPOUR PRESSURE VERSUS TEMPERATURE

- 3.1 Define *vapour pressure*. (2)
- 3.2 At what phase are the alkanes at room temperature?
Write down only GAS, LIQUID or SOLID.
Give a reason for the answer. (2)
- 3.3 How will the boiling points change if the atmospheric pressure decreases?
Choose from INCREASES, DECREASES or REMAIN THE SAME. (1)
- 3.4 What can be concluded from the curve regarding the relationship between intermolecular forces and vapour pressure? (2)
- 3.5 The table below shows the boiling point of propan-1-ol and propanoic acid.


	COMPOUND	BOILING POINT (°C)
A	Propan-1-ol	97
B	Propanoic acid	141

- 3.5.1 Define the term *boiling point*. (2)
- 3.5.2 Fully explain the difference in the boiling points. (4)

[13]

QUESTION 4 (Start on a NEW page.)

Consider the flow diagram below:

4.1 Consider reaction **I**.

Write down the:

4.1.1 Name of reaction (1)

4.1.2 STRUCTURAL FORMULA of the carboxylic acid need (2)

4.2 Consider reaction **II**.

Write down the:

4.2.1 NAME or FORMULA of the inorganic reagent needed (1)

4.2.2 IUPAC name of unsaturated hydrocarbon **P** (2)

4.3 Consider reaction **III**.

4.3.1 Write down the name of the type of addition reaction. (1)

4.3.2 Is compound **Q** a PRIMARY, SECONDARY or TERTIARY HALOALKANE? Give a reason for the answer. (2)

4.4 Compound **Q** is mixed with concentrated sodium hydroxide (NaOH) and strongly heated.

Write down the:

4.4.1 Type of reaction that occurred (1)

4.4.2 Balanced equation by using STRUCTURAL FORMULAE for the organic compounds and write the structural formula for the major product only in the reaction. (6)

[16]


QUESTION 5 (Start on a NEW page.)

5.1 Define the term *reaction rate*. (2)

A group of learners want to determine the percentage purity of a sample that contains calcium carbonate (CaCO_3). They react 2,5 g sample of impure calcium carbonate (CaCO_3) with EXCESS dilute hydrochloric acid (HCl) according to the balanced equation.



They measure the volume of carbon dioxide gas produced over time and obtained the graph below.



5.2 How will the learners know that the reaction has reached completion? (1)

5.3 Calculate the:

5.3.1 Average rate at which CO_2 is produced in $\text{cm}^3 \cdot \text{s}^{-1}$ (3)


5.3.2 Percentage purity of CaCO_3 if the molar volume of CO_2 is $24\,000 \text{ cm}^3 \cdot \text{mol}^{-1}$ at 25°C (5)

5.4 The experiment is repeated by using powdered sample of calcium carbonate instead of chunked sample of calcium carbonate.

5.4.1 Will the reaction rate INCREASE, DECREASE or REMAIN THE SAME? (1)

5.4.2 Explain the answer to QUESTION 5.4.1 by referring to the collision theory. (3)

- 5.5 The graph below represents the Maxwell-Boltzmann distribution curve for CO_2 (g) at 25°C .



5.5.1 What does the shaded area to the right of line **P** represents? (1)

5.5.2 How will an increase in temperature affect the shaded area?


Choose from INCREASES, DECREASES or NO EFFECT.

Give a reason for the answer.

(2)
[18]

QUESTION 6 (Start on a NEW page.)

When copper(II)chloride is dissolved in water the following chemical equilibrium equation is obtained. This can be used to demonstrate Le Chatelier's principle.



6.1 State *Le Chatelier's principle*. (2)

6.2 State whether the solution will become **BLUE** or **GREEN** when the following changes are made.


The colour of the equilibrium mixture is **BLUE**.

6.2.1 The mixture is cooled. (1)

6.2.2 Addition of saturated sodium chloride solution. (1)

6.3 Fully explain the answer to QUESTION 6.2.2 by referring to Le Chatelier's principle. (3)

6.4 Calcium carbonate is allowed to decompose at 1 000 °C in a sealed container according to the balanced chemical equation below.



Initially 0,24 mol of calcium carbonate is placed in a 2 dm³ container and heated. An equilibrium is established at 1 000 °C. At this temperature the equilibrium constant is 0,0385.

6.4.1 Will the K_c value INCREASE, DECREASE or REMAIN THE SAME if the temperature at which calcium carbonate decomposes is INCREASED?

Fully explain the answer. (3)

6.4.2 Calculate the mass of unreacted calcium carbonate that will remain in the container when the system reaches equilibrium at 1 000 °C. (7)

[17]

QUESTION 7 (Start on a NEW page.)

7.1 Consider the reactions below and answer questions that follow.

REACTION	CHEMICAL EQUATIONS
A	$\text{HSO}_4^- + \text{H}_2\text{O} \rightarrow \text{SO}_4^{2-} + \text{H}_3\text{O}^+$
B	$\text{KOH} + \text{CH}_3\text{COOH} \rightarrow \text{CH}_3\text{COOK} + \text{H}_2\text{O}$

7.1.1 Define a base in term of the *Arrhenius theory*. (2)

From the reactions above:

7.1.2 Which ONE of the reactions (**A** or **B**) represents the Arrhenius model? (1)

7.1.3 Write down a conjugate base of HSO_4^- (1)


7.2 Consider reaction **B**.

7.2.1 Give a reason why CH_3COOH can be regarded as a weak acid. (2)

7.2.2 Will CH_3COOK be ACIDIC, NEUTRAL or ALKALINE solution in water? (1)

7.2.3 Explain the answer in QUESTION 7.2.2 by using the relevant equation. (3)

7.3 During a titration, $12,5 \text{ cm}^3$ of sodium hydroxide (NaOH) solution with a concentration of $0,2 \text{ mol} \cdot \text{dm}^{-3}$ neutralises 25 cm^3 of sulphuric acid (H_2SO_4) solution, according to the following balanced chemical equation:




7.3.1 Write down the name of a suitable indicator for this neutralisation reaction. (1)

7.3.2 Calculate the pH of the H_2SO_4 solution. (7)

[18]

QUESTION 8 (Start on a NEW page.)

- 8.1 A galvanic cell is set up under standard conditions using unknown metal **M** and silver as an electrode. The standard electrode potential for the unknown metal **M** is less than 0 V.



- 8.1.1 Write down the energy conversion in this cell. (2)
- 8.1.2 Which electrode is the anode in the cell (Ag or Metal **M**)?
Explain the answer. (3)
- 8.2 The initial potential difference for this cell is 1,54 V.
- 8.2.1 Identify metal **M** by calculation. (5)
- 8.2.2 Write down the cell notation for the cell. (3)
- 8.3 How will the initial reading on the voltmeter change when the following changes are made to the cell above?


Write down only INCREASES, DECREASES or REMAINS THE SAME

- 8.3.1 Surface area of the Ag electrode is increased. (1)
- 8.3.2 The concentration of AgNO_3 solution is increased to $2 \text{ mol} \cdot \text{dm}^{-3}$. (1)

[15]

QUESTION 9 (Start on a NEW page.)

The electrolytic cell, represented in the diagram below, is set up to purify copper which contains zinc and silver impurities.



- 9.1 Define the term *electrolyte*. (2)
- 9.2 Is this reaction ENDOTHERMIC or EXOTHERMIC? (1)
- 9.3 Which of the electrodes (**X** or **Z**) is connected to the positive terminal of the battery? (1)
- 9.4 Write down the half reaction that takes place at electrode **Z**. (2)
- 9.5 During the electrolysis it is observed that zinc is also oxidised but not silver.
Give a reason for this observation by referring to the relative strengths of the reducing agents. (1)
- 9.6 An unknown solution (MCl_2) was electrolysed using copper electrodes. After some time $1,806 \times 10^{22}$ electrons were gained at the cathode while the cathode gained 0,6 g.

Determine, by calculations, the formula of an unknown solution. (5)
[12]

TOTAL: 150

**NATIONAL SENIOR CERTIFICATE
NASIONALE SENIOR SERTIFIKAAT**

**DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)**

**GEGEWENS VIR FISIESE WETENSKAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)**

TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESE KONSTANTES

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure <i>Standaarddruk</i>	p^θ	$1,013 \times 10^5 \text{ Pa}$
Molar gas volume at STP <i>Molêre gasvolume teen STD</i>	V_m	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard temperature <i>Standaardtemperatuur</i>	T^θ	273 K
Charge on electron <i>Lading op elektron</i>	e	$-1,6 \times 10^{-19} \text{ C}$
Avogadro's constant <i>Avogadro se konstante</i>	N_A	$6,02 \times 10^{23} \text{ mol}^{-1}$

TABLE 2: FORMULAE/TABEL 2: FORMULES

$n = \frac{m}{M}$ or/of $n = \frac{N}{N_A}$ or/of $n = \frac{V}{V_m}$	$c = \frac{n}{V}$ or/of $c = \frac{m}{MV}$ $\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$	$\text{pH} = -\log[\text{H}_3\text{O}^+]$ $K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14}$ at /by 298K
$E^\theta_{\text{cell}} = E^\theta_{\text{cathode}} - E^\theta_{\text{anode}} / E^\theta_{\text{sel}} = E^\theta_{\text{katode}} - E^\theta_{\text{anode}}$ $E^\theta_{\text{cell}} = E^\theta_{\text{reduction}} - E^\theta_{\text{oxidation}} / E^\theta_{\text{sel}} = E^\theta_{\text{reduksie}} - E^\theta_{\text{oksidasie}}$ $E^\theta_{\text{cell}} = E^\theta_{\text{oxidising agent}} - E^\theta_{\text{reducing agent}} / E^\theta_{\text{sel}} = E^\theta_{\text{oksideermiddel}} - E^\theta_{\text{reduseermiddel}}$		
$q = I\Delta t$ $n = \frac{Q}{e}$ or/of $n = \frac{Q}{q_e}$		



TABLE 3: THE PERIODIC TABLE OF ELEMENTS/TABEL 3: DIE PERIODIEKE TABEL VAN ELEMENTE

Atoomgetal Atomic number																															
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18														
(I)	(II)											(III)	(IV)	(V)	(VI)	(VII)	(VIII)														
KEY/ SLEUTEL																															
							Elektronegatiwiteit Electronegativity	29 σ↓ Cu		Simbool Symbol																					
										Approximate relative atomic mass/ Benaderde relatiewe atoommassa																					
1 2,1 H	4 9 Be	12 24 Mg	19 39 K	20 40 Ca	23 47 V	24 52 Cr	25 55 Mn	26 56 Fe	27 59 Co	28 59 Ni	29 63,5 Cu	30 65 Zn	31 70 Ga	32 73 Ge	33 75 As	34 79 Se	35 84 Br	36 84 Kr	54 131 Xe	86 222 Rn											
3 7 Li	11 23 Na	12 24 Mg	13 27 Al	14 28 Si	15 31 P	16 32 S	17 35,5 Cl	18 40 Ar	19 39 K	20 40 Ca	21 45 Sc	22 48 Ti	23 51 V	24 52 Cr	25 55 Mn	26 56 Fe	27 59 Co	28 59 Ni	29 63,5 Cu	30 65 Zn	31 70 Ga	32 73 Ge	33 75 As	34 79 Se	35 84 Br	36 84 Kr	54 131 Xe	86 222 Rn			
55 133 Cs	56 137 Ba	57 139 La	58 140 Ce	59 141 Pr	60 144 Nd	61 147 Pm	62 150 Sm	63 152 Eu	64 157 Gd	65 159 Tb	66 163 Dy	67 165 Ho	68 167 Er	69 169 Tm	70 173 Yb	71 175 Lu	72 177 Hf	73 181 Ta	74 184 W	75 186 Re	76 190 Os	77 192 Ir	78 195 Pt	79 197 Au	80 201 Hg	81 204 Tl	82 207 Pb	83 209 Bi	84 210 Po	85 210 At	86 222 Rn
87 226 Fr	88 226 Ra	89 227 Ac	90 232 Th	91 232 Pa	92 238 U	93 238 Np	94 238 Pu	95 238 Am	96 238 Cm	97 238 Bk	98 238 Cf	99 238 Es	100 238 Fm	101 238 Md	102 238 No	103 238 Lr	104 238 Rf	105 238 Db	106 238 Sg	107 238 Bh	108 238 Hs	109 238 Mt	110 238 Ds	111 238 Rg	112 238 Cn	113 238 Nh	114 238 Fl	115 238 Mc	116 238 Lv	117 238 Ts	118 238 Og

TABLE 4A: STANDARD REDUCTION POTENTIALS
TABEL 4A: STANDAARD REDUKSIEPOTENSIALE

Half-reactions/Halfreaksies		E^{θ} (V)
$F_2(g) + 2e^-$	$\rightleftharpoons 2F^-$	+ 2,87
$Co^{3+} + e^-$	$\rightleftharpoons Co^{2+}$	+ 1,81
$H_2O_2 + 2H^+ + 2e^-$	$\rightleftharpoons 2H_2O$	+ 1,77
$MnO_4^- + 8H^+ + 5e^-$	$\rightleftharpoons Mn^{2+} + 4H_2O$	+ 1,51
$Cl_2(g) + 2e^-$	$\rightleftharpoons 2Cl^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^-$	$\rightleftharpoons 2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^-$	$\rightleftharpoons 2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^-$	$\rightleftharpoons Mn^{2+} + 2H_2O$	+ 1,23
$Pt^{2+} + 2e^-$	$\rightleftharpoons Pt$	+ 1,20
$Br_2(l) + 2e^-$	$\rightleftharpoons 2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^-$	$\rightleftharpoons NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^-$	$\rightleftharpoons Hg(l)$	+ 0,85
$Ag^+ + e^-$	$\rightleftharpoons Ag$	+ 0,80
$NO_3^- + 2H^+ + e^-$	$\rightleftharpoons NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^-$	$\rightleftharpoons Fe^{2+}$	+ 0,77
$O_2(g) + 2H^+ + 2e^-$	$\rightleftharpoons H_2O_2$	+ 0,68
$I_2 + 2e^-$	$\rightleftharpoons 2I^-$	+ 0,54
$Cu^+ + e^-$	$\rightleftharpoons Cu$	+ 0,52
$SO_2 + 4H^+ + 4e^-$	$\rightleftharpoons S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^-$	$\rightleftharpoons 4OH^-$	+ 0,40
$Cu^{2+} + 2e^-$	$\rightleftharpoons Cu$	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^-$	$\rightleftharpoons SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^-$	$\rightleftharpoons Cu^+$	+ 0,16
$Sn^{4+} + 2e^-$	$\rightleftharpoons Sn^{2+}$	+ 0,15
$S + 2H^+ + 2e^-$	$\rightleftharpoons H_2S(g)$	+ 0,14
$2H^+ + 2e^-$	$\rightleftharpoons H_2(g)$	0,00
$Fe^{3+} + 3e^-$	$\rightleftharpoons Fe$	- 0,06
$Pb^{2+} + 2e^-$	$\rightleftharpoons Pb$	- 0,13
$Sn^{2+} + 2e^-$	$\rightleftharpoons Sn$	- 0,14
$Ni^{2+} + 2e^-$	$\rightleftharpoons Ni$	- 0,27
$Co^{2+} + 2e^-$	$\rightleftharpoons Co$	- 0,28
$Cd^{2+} + 2e^-$	$\rightleftharpoons Cd$	- 0,40
$Cr^{3+} + e^-$	$\rightleftharpoons Cr^{2+}$	- 0,41
$Fe^{2+} + 2e^-$	$\rightleftharpoons Fe$	- 0,44
$Cr^{3+} + 3e^-$	$\rightleftharpoons Cr$	- 0,74
$Zn^{2+} + 2e^-$	$\rightleftharpoons Zn$	- 0,76
$2H_2O + 2e^-$	$\rightleftharpoons H_2(g) + 2OH^-$	- 0,83
$Cr^{2+} + 2e^-$	$\rightleftharpoons Cr$	- 0,91
$Mn^{2+} + 2e^-$	$\rightleftharpoons Mn$	- 1,18
$Al^{3+} + 3e^-$	$\rightleftharpoons Al$	- 1,66
$Mg^{2+} + 2e^-$	$\rightleftharpoons Mg$	- 2,36
$Na^+ + e^-$	$\rightleftharpoons Na$	- 2,71
$Ca^{2+} + 2e^-$	$\rightleftharpoons Ca$	- 2,87
$Sr^{2+} + 2e^-$	$\rightleftharpoons Sr$	- 2,89
$Ba^{2+} + 2e^-$	$\rightleftharpoons Ba$	- 2,90
$Cs^+ + e^-$	$\rightleftharpoons Cs$	- 2,92
$K^+ + e^-$	$\rightleftharpoons K$	- 2,93
$Li^+ + e^-$	$\rightleftharpoons Li$	- 3,05

Increasing oxidising ability/Toenemende oksiderende vermoë

Increasing reducing ability/Toenemende reduserende vermoë




TABLE 4B: STANDARD REDUCTION POTENTIALS
TABEL 4B: STANDAARD REDUKSIEPOTENSIALE

Half-reactions/Halfreaksies			$E^{\ominus}$ (V)
$\text{Li}^+ + \text{e}^-$	$\rightleftharpoons$	Li	-3,05
$\text{K}^+ + \text{e}^-$	$\rightleftharpoons$	K	-2,93
$\text{Cs}^+ + \text{e}^-$	$\rightleftharpoons$	Cs	-2,92
$\text{Ba}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Ba	-2,90
$\text{Sr}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Sr	-2,89
$\text{Ca}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Ca	-2,87
$\text{Na}^+ + \text{e}^-$	$\rightleftharpoons$	Na	-2,71
$\text{Mg}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Mg	-2,36
$\text{Al}^{3+} + 3\text{e}^-$	$\rightleftharpoons$	Al	-1,66
$\text{Mn}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Mn	-1,18
$\text{Cr}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Cr	-0,91
$2\text{H}_2\text{O} + 2\text{e}^-$	$\rightleftharpoons$	$\text{H}_2(\text{g}) + 2\text{OH}^-$	-0,83
$\text{Zn}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Zn	-0,76
$\text{Cr}^{3+} + 3\text{e}^-$	$\rightleftharpoons$	Cr	-0,74
$\text{Fe}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Fe	-0,44
$\text{Cr}^{3+} + \text{e}^-$	$\rightleftharpoons$	Cr^{2+}	-0,41
$\text{Cd}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Cd	-0,40
$\text{Co}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Co	-0,28
$\text{Ni}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Ni	-0,27
$\text{Sn}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Sn	-0,14
$\text{Pb}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Pb	-0,13
$\text{Fe}^{3+} + 3\text{e}^-$	$\rightleftharpoons$	Fe	-0,06
$2\text{H}^+ + 2\text{e}^-$	$\rightleftharpoons$	$\text{H}_2(\text{g})$	0,00
$\text{S} + 2\text{H}^+ + 2\text{e}^-$	$\rightleftharpoons$	$\text{H}_2\text{S}(\text{g})$	+0,14
$\text{Sn}^{4+} + 2\text{e}^-$	$\rightleftharpoons$	Sn^{2+}	+0,15
$\text{Cu}^{2+} + \text{e}^-$	$\rightleftharpoons$	Cu^+	+0,16
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^-$	$\rightleftharpoons$	$\text{SO}_2(\text{g}) + 2\text{H}_2\text{O}$	+0,17
$\text{Cu}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Cu	+0,34
$2\text{H}_2\text{O} + \text{O}_2 + 4\text{e}^-$	$\rightleftharpoons$	4OH^-	+0,40
$\text{SO}_2 + 4\text{H}^+ + 4\text{e}^-$	$\rightleftharpoons$	$\text{S} + 2\text{H}_2\text{O}$	+0,45
$\text{Cu}^+ + \text{e}^-$	$\rightleftharpoons$	Cu	+0,52
$\text{I}_2 + 2\text{e}^-$	$\rightleftharpoons$	2I^-	+0,54
$\text{O}_2(\text{g}) + 2\text{H}^+ + 2\text{e}^-$	$\rightleftharpoons$	H_2O_2	+0,68
$\text{Fe}^{3+} + \text{e}^-$	$\rightleftharpoons$	Fe^{2+}	+0,77
$\text{NO}_3^- + 2\text{H}^+ + \text{e}^-$	$\rightleftharpoons$	$\text{NO}_2(\text{g}) + \text{H}_2\text{O}$	+0,80
$\text{Ag}^+ + \text{e}^-$	$\rightleftharpoons$	Ag	+0,80
$\text{Hg}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	$\text{Hg}(\ell)$	+0,85
$\text{NO}_3^- + 4\text{H}^+ + 3\text{e}^-$	$\rightleftharpoons$	$\text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+0,96
$\text{Br}_2(\ell) + 2\text{e}^-$	$\rightleftharpoons$	2Br^-	+1,07
$\text{Pt}^{2+} + 2\text{e}^-$	$\rightleftharpoons$	Pt	+1,20
$\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^-$	$\rightleftharpoons$	$\text{Mn}^{2+} + 2\text{H}_2\text{O}$	+1,23
$\text{O}_2(\text{g}) + 4\text{H}^+ + 4\text{e}^-$	$\rightleftharpoons$	$2\text{H}_2\text{O}$	+1,23
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^-$	$\rightleftharpoons$	$2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+1,33
$\text{Cl}_2(\text{g}) + 2\text{e}^-$	$\rightleftharpoons$	2Cl^-	+1,36
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^-$	$\rightleftharpoons$	$\text{Mn}^{2+} + 4\text{H}_2\text{O}$	+1,51
$\text{H}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^-$	$\rightleftharpoons$	$2\text{H}_2\text{O}$	+1,77
$\text{Co}^{3+} + \text{e}^-$	$\rightleftharpoons$	Co^{2+}	+1,81
$\text{F}_2(\text{g}) + 2\text{e}^-$	$\rightleftharpoons$	2F^-	+2,87

Increasing oxidising ability/Toenemende oksiderende vermoë

Increasing reducing ability/Toenemende reduserende vermoë

