

SA EXAM PAPERS This Paper was downloaded from SAEXAMPAPERS
SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



**SA EXAM
PAPERS**

SA EXAM PAPERS

Proudly South African



DEPARTMENT OF EDUCATION
DEPARTEMENT VAN ONDERWYS
LEFAPHA LA THUTO
ISEBE LEZEMFUNDO

**PROVINSIALE VOORBEREIDENDE EKSAMEN/
PROVINCIAL PREPARATORY EXAMINATION**

GRAAD/GRADE 12

FISIESE WETENSKAPPE/PHYSICAL SCIENCES

VRAESTEL/PAPER 2

CHEMIE/CHEMISTRY

SEPTEMBER 2025

PUNTE/MARKS: 150

TYD/TIME: 3 uur/hours

**Hierdie vraestel bestaan uit 15 bladsye, 'n grafiekblad en 4 gegewensblaaie./
This question paper consists of 15 pages, a graph sheet and 4 data sheets.**



INSTRUCTIONS AND INFORMATION


1. Write your name in the appropriate space on the ANSWER BOOK.
2. This question paper consists of NINE questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two subquestions, e.g. between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. Show ALL formulae and substitutions in ALL calculations.
8. Round off your FINAL numerical answers to a minimum of TWO decimal places.
9. Give brief motivations, discussions, etc. where required.
10. You are advised to use the attached DATA SHEETS.
11. Write neatly and legibly.



QUESTION 1: MULTIPLE-CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Each question has only ONE correct answer. Choose the correct answer and write only the letter (A–D) next to the question numbers (1.1 to 1.10) in the ANSWER BOOK, e.g. 1.11 E.

- 1.1 Which of the following reactants is used to prepare ethyl propanoate?
- A Ethane and propanoic acid
- B Propanol and ethanoic acid
- C Ethanol and propanoic acid
- D Ethene and propanol (2)
- 1.2 The structural formula of an organic compound is given below.



The IUPAC name of this compound is ...


- A 4-bromo-3-ethyl-5,6-dimethylheptane.
- B 4-bromo-5-ethyl-2,6-dimethylheptane.
- C 3-ethyl-4-bromo-2,6-dimethylheptane.
- D 4-bromo-3-ethyl-2,6-dimethylheptane. (2)
- 1.3 Which ONE of the following reaction types may be used to prepare ethene from pentane?
- A Addition
- B Cracking
- C Substitution
- D Hydrolysis (2)

- 1.4 Which ONE of the following combinations of values of activation energy (E_A) and heat of reaction (ΔH) CANNOT be obtained for any reaction?

	E_A (kJ·mol ⁻¹)	ΔH (kJ·mol ⁻¹)
A	60	-120
B	60	+120
C	120	+60
D	120	-60

(2)

- 1.5 Consider the following reaction that reaches equilibrium in a beaker:



A few drops of concentrated $\text{HCl}(\text{aq})$ are now added to the beaker.

Which ONE of the following combinations correctly identifies the DISTURBANCE ON THE SYSTEM and the SYSTEM'S RESPONSE to the disturbance?

	DISTURBANCE IN THE SYSTEM	SYSTEM'S RESPONSE
A	$[\text{H}^+]$ increases	Forward reaction favoured
B	$[\text{H}^+]$ decreases	Reverse reaction favoured
C	$[\text{CrO}_4^{2-}]$ increases	Forward reaction favoured
D	$[\text{CrO}_4^{2-}]$ decreases	Reverse reaction favoured





(2)

- 1.6 Which ONE of the descriptions below is TRUE for a chemical reaction in equilibrium?

	CONCENTRATIONS OF REACTANTS AND PRODUCTS	RATES OF FORWARD AND REVERSE REACTIONS
A	Remain constant	Equal
B	Remain constant	Not equal
C	Equal	Equal
D	Not equal	Not equal

(2)

1.7 The pH of a solution of H_3PO_4 is 4,5. Which ONE of the following relations is CORRECT for the H_3PO_4 solution?







(2)

1.8 When sulphuric acid is titrated with ammonia, ammonium sulphate ($(\text{NH}_4)_2\text{SO}_4$) is produced. Which ONE of the combinations is correct for the indicator used in this titration and the pH of $(\text{NH}_4)_2\text{SO}_4$?

	INDICATOR	pH
A	Methyl orange	pH < 7
B	Phenolphthalein	pH > 7
C	Phenolphthalein	pH < 7
D	Bromothymol blue	pH = 7

(2)

1.9 Which ONE of the following is a NON-SPONTANEOUS redox reaction? Refer to the Table of Standard Reduction Potentials (Table 4A or 4B).



(2)



1.10 The following characteristics may be used to describe an electrochemical cell:

- I The chemical reaction is spontaneous
- II The reaction requires energy from an electrical source.
- III The anode is the positive electrode of the cell.

Which ONE of the following combinations are specific to an electrolytic cell?

- A Only I
- B Only II
- C I and III
- D II and III

(2)
[20]



QUESTION 2 (Start on a new page.)

The letters **A** to **F** in the table below represent six organic compounds.

A	$\text{CH}_3(\text{CH}_2)_4\text{CO}_2\text{H}$	D	4-methylpentan-1-ol
B	$ \begin{array}{c} \text{H} \qquad \text{O} \qquad \text{H} \\ \qquad // \qquad \\ \text{H}-\text{C}-\text{O}-\text{C}-\text{C}-\text{H} \\ \qquad \qquad \\ \text{H}-\text{C}-\text{H} \qquad \text{H} \\ \\ \text{H}-\text{C}-\text{H} \\ \\ \text{H} \end{array} $	E	$ \begin{array}{c} \text{H} \qquad \qquad \text{H} \\ \qquad \qquad \\ \text{H}-\text{C}-\text{C}\equiv\text{C}-\text{C}-\text{H} \\ \qquad \qquad \\ \text{H} \qquad \qquad \text{CH}_3 \end{array} $
C	$\text{C}_3\text{H}_6\text{O}$	F	$(\text{CH}_3)_2\text{CC}(\text{C}_2\text{H}_5)\text{CH}_2\text{CH}_3$

Write down the:

- 2.1 LETTER of the compound that represents an aldehyde (1)
- 2.2 STRUCTURAL FORMULA of the functional group of compound **A** (1)
- 2.3 LETTERS of the compounds that represent unsaturated hydrocarbons (2)
- 2.4 STRUCTURAL FORMULA of compound **D** (3)
- 2.5 General formula for compound **E** (1)
- 2.6 IUPAC NAMES of the TWO compounds represented by **C** (2)
- 2.7 LETTER of the compound that represents an ester (1)
- 2.8 LETTER of the compound that has a hydroxyl group as a functional group (1)
- 2.9 NAME of the functional group of the ketones (1)
- 2.10 IUPAC NAME of compound **B** (2)

[15]

QUESTION 3 (Start on a new page.)

An investigation was carried out to determine the various factors affecting the boiling point of different alcohols **A** to **C**, shown in the table below:

A	$\text{CH}_3\text{CH}(\text{OH})\text{CH}_3$
B	$\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$
C	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$

3.1 Which ONE of the compounds (**A**, **B** or **C**) has the highest boiling point?

Explain the answer. (3)

3.2 Write down the type of alcohol represented by compound **A**.
Choose from PRIMARY, SECONDARY or TERTIARY.

Give a reason for the answer. (2)

Consider compounds **D** to **G**, shown in the table below.

	COMPOUND	MOLECULAR MASS ($\text{g}\cdot\text{mol}^{-1}$)
D	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$	72
E	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$	74
F	$\text{CH}_3\text{CH}_2\text{COOH}$	74
G	$\text{CH}_3\text{CH}_2\text{OOCH}$	74

3.3 Which ONE of compounds **E**, **F** or **G** has the lowest vapour pressure?

Fully explain the answer. (4)

3.4 Compounds **F** and **G** are isomers.
Identify the type of isomerism they represent.


Give a reason for the answer. (2)

3.5 Write down a balanced equation, using MOLECULAR FORMULAE, for the combustion of compound **D** in excess oxygen. (3)

[14]

QUESTION 4 (Start on a new page.)

Study the flow diagram of various organic reactions and answer the questions that follow:



- 4.1 For Reaction I, write down the:
- 4.1.1 NAME of the **substitution** reaction (1)
- 4.1.2 STRUCTURAL FORMULA for compound **A** (2)
- 4.2 Compound **B** is formed in the presence of water. Write down the:
- 4.2.1 NAME of the substitution reaction (1)
- 4.2.2 IUPAC NAME of organic compound **B** (2)
- 4.3 Reaction III is an elimination reaction. Write down the:
- 4.3.1 NAME of the elimination reaction (1)
- 4.3.2 ONE reaction condition (1)
- 4.3.3 IUPAC NAME of the organic product formed (2)
- 4.3.4 Balanced equation for the above reaction using CONDENSED STRUCTURAL FORMULAE. (3)
- 4.4 Reaction IV takes place in the presence of a catalyst. Write down the:
- 4.4.1 Name of the catalyst (1)
- 4.4.2 Type of reaction (1)



- 4.5 In another reaction, the position isomer of compound **B** reacts with methanoic acid in the presence of a catalyst.

Write down the:


4.5.1 Name of this reaction (1)

4.5.2 Balanced equation using STRUCTURAL FORMULAE (5)
[21]



QUESTION 5 (Start on a new page.)

10 g of aluminium powder reacts with a limiting amount of sulphuric acid according to the following balanced equation:



Aluminium is in excess.

The rate of the reaction between aluminium powder and sulphuric acid is investigated at 20 °C. Hydrogen gas is collected in a syringe, and the volume of hydrogen gas is measured at different time intervals.

The following results were obtained:

TIME (minutes)	H ₂ VOLUME (dm ³)
0	0
2	1,2
4	2,4
6	3,6
8	4,2
10	4,5
12	4,5

- 5.1 Define the term *reaction rate*. (2)
- 5.2 Calculate the average reaction rate, in dm³·min⁻¹, during the first 10 minutes. (2)
- 5.3 Use the data in the table to draw a graph of the volume of hydrogen gas collected versus time on the attached GRAPH SHEET. (3)
- 5.4 Use the collision theory to explain the change in the reaction rate over time. Assume that the temperature remains constant. (3)
- 5.5 Calculate the mass of aluminium powder that was left at the end of the reaction. The molar gas volume at this temperature is 24 dm³·mol⁻¹ (6)
- 5.6 A change in reaction conditions will affect the sketch drawn on the GRAPH SHEET in QUESTION 5.3.


On the same GRAPH SHEET sketch graphs for the following changes:

- 5.6.1 The temperature is increased to 40 °C. Label this graph **B** (2)
- 5.6.2 10 g aluminium granules are used. Label this graph **C** (2)

[20]


QUESTION 6 (Start on a new page.)

Consider the following balanced equation for a reversible reaction in a closed system below:



6.1 Define the term *dynamic equilibrium*. (2)

The sketch graph below shows the relationship between the value of the equilibrium constant (K_c) and the temperature for this reaction.



6.2 Write down the sign of the heat of reaction (ΔH) for the above reaction. Choose from POSITIVE or NEGATIVE.

Explain the answer. (3)

6.3 The pressure on the system is decreased at constant temperature.

Use Le Chatelier's principle to explain how this change affects the yield of $\text{NO}(\text{g})$. (3)

6.4 1,4 mol of ammonia gas (NH_3) and 0,8 mol of oxygen gas (O_2) are placed in a 250 cm^3 container at 1 000 kPa and allowed to reach equilibrium at $800 \text{ }^\circ\text{C}$. At this temperature and pressure, the amount of water vapour present is 12,6 g.

6.4.1 Calculate the value of the equilibrium constant (K_c) (7)


6.4.2 More ammonia gas is now added to the equilibrium mixture. How will this change affect the yield of $\text{NO}(\text{g})$?

Choose from INCREASES, DECREASES or REMAINS THE SAME. (1)

[16]


QUESTION 7 (Start on a new page.)

- 7.1 Hydrogen iodide (HI) reacts with water according to the following balanced equation:




The K_a value for HI is $3,2 \times 10^9$ at 25°C .

- 7.1.1 Is hydrogen iodide a weak or a strong acid? Give a reason for the answer. (2)
- 7.1.2 Write down the formulae for the TWO bases in the reaction above. (2)
- 7.2 60 cm^3 of a hydrogen iodide (HI) solution reacts with an unknown quantity of magnesium hydroxide (Mg(OH)_2) according to the following balanced equation:



The EXCESS hydrogen iodide is then neutralised with 18 cm^3 NaOH with a concentration of $0,3\text{ mol.dm}^{-3}$ according to the following balanced equation:




- 7.2.1 Calculate the pH of the excess hydrogen iodide. (5)
- 7.2.2 Calculate the initial mass of magnesium hydroxide used in Reaction 1 if the initial concentration of hydrogen iodide was $0,4\text{ mol.dm}^{-3}$. (6)
- 7.2.3 Bromothymol blue is used as an indicator. Explain why it is the most suitable indicator by referring to the pH at the equivalence point. (2)

[17]

QUESTION 8 (Start on a new page.)

The diagram below represents an electrochemical cell operating under standard conditions, featuring tin (Sn) and an unknown metal, **Z**.



8.1 It is found that the mass of electrode **Z** increases whilst the cell is functioning.

For this cell, write down:

8.1.1 The electrode that acts as the ANODE. Choose from **Sn** or **Z**.

Give a reason for the answer. (2)

8.1.2 How will the concentration of Sn^{2+} change while the cell is in operation?

Choose from INCREASES, DECREASES or REMAINS THE SAME. (1)

8.1.3 The reduction half-reaction (2)

8.1.4 In which direction (**A** or **B**) will CATIONS move in the salt bridge? (1)

8.1.5 The cell notation (3)

8.2 The initial emf of this cell is 0,94 V. Identify metal **Z** by means of a calculation. (5)

A stronger reducing agent is now used with the same oxidising agent under the same conditions.


8.3 How will this affect the initial emf of the cell?

Choose from INCREASES, DECREASES or NO EFFECT. (1)

[15]

QUESTION 9 (Start on a new page.)

The diagram below represents the cell used for electroplating an iron spoon with copper.




- 9.1 Define the term *electrolyte*. (2)
- 9.2 What type of power **source** is used to drive the reaction in this cell? Choose from AC or DC. (2)
- Give a reason for the answer. (2)
- 9.3 Define the term *reduction* in terms of electron transfer. (2)
- 9.4 Write down a half-reaction that takes place at the ANODE. (2)
- 9.5 Write down ONE observation that can be made after the cell has been operating for some time. (1)
- 9.6 Name the metal that is labelled 'Electrode'. (1)
- 9.7 How will the concentration of the copper (II) nitrate solution change while the cell is in operation? Choose from INCREASES, DECREASES or REMAINS THE SAME. (2)
- Give a reason for the answer. (2)

[12]**TOTAL: 150**

QUESTION 5.3

Hand in this **GRAPH SHEET** with your **ANSWER BOOK**.



**INFORMATION FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)**

**GEGEWENS VIR FISIESTE WETENSAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)**

TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESTE KONSTANTES

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure <i>Standaarddruk</i>	p^θ	$1,013 \times 10^5 \text{ Pa}$
Molar gas volume at STP <i>Molêre gas volume by STD</i>	V_m	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard temperature <i>Standaardtemperatuur</i>	T^θ	273 K
Charge on electron <i>Lading op elektron</i>	e	$-1,6 \times 10^{-19} \text{ C}$
Avogadro's constant <i>Avogadro-konstante</i>	N_A	$6,02 \times 10^{23} \text{ mol}^{-1}$


TABLE 2: FORMULAE/TABEL 2: FORMULES

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ or/of $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$
$\frac{c_a v_a}{c_b v_b} = \frac{n_a}{n_b}$	$\text{pH} = -\log[\text{H}_3\text{O}^+]$
$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14}$ at/by 298 K	
$E_{\text{cell}}^\theta = E_{\text{cathode}}^\theta - E_{\text{anode}}^\theta / E_{\text{sel}}^\theta = E_{\text{katode}}^\theta - E_{\text{anode}}^\theta$	
or/of $E_{\text{cell}}^\theta = E_{\text{reduction}}^\theta - E_{\text{oxidation}}^\theta / E_{\text{sel}}^\theta = E_{\text{reduksie}}^\theta - E_{\text{oksidasie}}^\theta$	
or/of $E_{\text{cell}}^\theta = E_{\text{oxidising agent}}^\theta - E_{\text{reducing agent}}^\theta / E_{\text{sel}}^\theta = E_{\text{oksideermiddel}}^\theta - E_{\text{reduseermiddel}}^\theta$	



TABLE 3: THE PERIODIC TABLE OF ELEMENTS
TABEL 3: DIE PERIODIEKE TABEL VAN ELEMENTE

1 (I)	2 (II)	3	4	5	6	7	8	9	10	11	12	13 (III)	14 (IV)	15 (V)	16 (VI)	17 (VII)	
1 H 1,01	4 Be 9,01	12 Mg 24,31	20 Ca 40,08	38 Sr 87,62	56 Ba 137,33	88 Ra 226,03	21 Sc 44,96	39 Y 88,91	57 La 138,91	89 Ac 227,03	22 Ti 47,88	40 Zr 91,22	58 Ce 140,12	90 Th 232,04	59 Pr 140,91	91 Pa 231,04	87 Fr 223,02
3 Li 6,94	9 Na 22,99	19 K 39,10	37 Rb 85,47	55 Cs 132,91	85 Fr 223,02	23 V 50,94	41 Nb 92,91	73 Ta 180,95	91 Hf 178,49	59 Ce 140,12	90 Th 232,04	24 Cr 51,99	42 Mo 95,94	74 W 183,84	92 U 238,03	60 Nd 144,24	92 U 238,03
4 Be 9,01	12 Mg 24,31	20 Ca 40,08	38 Sr 87,62	56 Ba 137,33	88 Ra 226,03	25 Mn 54,94	43 Tc 98,91	75 Re 186,21	93 Nb 92,91	61 Pm 144,91	93 Np 237,05	26 Fe 55,85	44 Ru 101,07	76 Os 190,23	94 Pu 242,02	62 Sm 150,36	94 Pu 242,02
5 B 10,81	13 Al 26,98	27 Ga 69,72	49 In 114,82	81 Tl 204,38	113 Nh 286,10	27 Co 58,93	45 Rh 101,07	77 Ir 192,22	95 Pd 106,42	63 Eu 151,96	95 Am 243,06	28 Si 28,09	50 Sn 118,71	82 Pb 207,20	126 Lv 293,10	64 Gd 157,25	96 Cm 247,07
6 C 12,01	14 Si 28,09	32 Ge 72,64	60 Zn 65,38	112 Cn 283,80	114 Fl 289,10	30 Zn 65,38	46 Pd 106,42	78 Pt 195,08	108 Ag 107,87	65 Tb 158,93	97 Bk 247,07	15 P 30,97	73 As 74,92	159 Ho 162,50	209 Og 294,11	66 Dy 162,50	98 Cf 251,08
7 N 14,01	15 P 30,97	33 As 74,92	62 Ni 58,69	114 Po 209,00	116 Lv 293,10	31 Ga 69,72	47 Cu 63,55	79 Au 196,97	110 Dn 286,10	67 Er 167,26	99 Es 254,09	16 S 32,06	75 Sb 121,76	173 Yb 173,05	210 Nh 286,10	68 Er 167,26	100 Fm 254,09
8 O 16,00	16 S 32,06	34 Se 78,96	64 Cu 63,55	116 Og 294,11	118 Og 294,11	35 Br 79,90	48 Cd 112,41	80 Hg 200,59	118 Og 294,11	69 Tm 168,93	101 Md 258,11	17 Cl 35,45	83 Bi 208,98	175 Lu 174,97	211 Nh 286,10	69 Tm 168,93	102 No 259,10
9 F 18,99	17 Cl 35,45	35 Br 79,90	66 Zn 65,38	118 Og 294,11	120 Og 294,11	36 Kr 83,80	59 Ni 58,69	81 Au 196,97	119 Og 294,11	70 Yb 173,05	102 No 259,10	18 Ar 39,95	85 At 210,09	177 Lu 176,93	212 Nh 286,10	70 Yb 173,05	103 Lr 260,11



Approximate relative atomic mass
Benaderde relatiewe atoommassa



TABLE 4A: STANDARD REDUCTION POTENTIALS
TABEL 4A: STANDAARD-REDUKSIEPOTENSIALE

Half-reactions/Halfreaksies	E^{θ} (V)
$F_2(g) + 2e^- \rightleftharpoons 2F^-$	+ 2,87
$Co^{3+} + e^- \rightleftharpoons Co^{2+}$	+ 1,81
$H_2O_2 + 2H^+ + 2e^- \rightleftharpoons 2H_2O$	+1,77
$MnO_4^- + 8H^+ + 5e^- \rightleftharpoons Mn^{2+} + 4H_2O$	+ 1,51
$Cl_2(g) + 2e^- \rightleftharpoons 2Cl^-$	+ 1,36
$Cr_2O_7^{2-} + 14H^+ + 6e^- \rightleftharpoons 2Cr^{3+} + 7H_2O$	+ 1,33
$O_2(g) + 4H^+ + 4e^- \rightleftharpoons 2H_2O$	+ 1,23
$MnO_2 + 4H^+ + 2e^- \rightleftharpoons Mn^{2+} + 2H_2O$	+ 1,23
$Pt^{2+} + 2e^- \rightleftharpoons Pt$	+ 1,20
$Br_2(l) + 2e^- \rightleftharpoons 2Br^-$	+ 1,07
$NO_3^- + 4H^+ + 3e^- \rightleftharpoons NO(g) + 2H_2O$	+ 0,96
$Hg^{2+} + 2e^- \rightleftharpoons Hg(l)$	+ 0,85
$Ag^+ + e^- \rightleftharpoons Ag$	+ 0,80
$NO_3^- + 2H^+ + e^- \rightleftharpoons NO_2(g) + H_2O$	+ 0,80
$Fe^{3+} + e^- \rightleftharpoons Fe^{2+}$	+ 0,77
$O_2(g) + 2H^+ + 2e^- \rightleftharpoons H_2O_2$	+ 0,68
$I_2 + 2e^- \rightleftharpoons 2I^-$	+ 0,54
$Cu^+ + e^- \rightleftharpoons Cu$	+ 0,52
$SO_2 + 4H^+ + 4e^- \rightleftharpoons S + 2H_2O$	+ 0,45
$2H_2O + O_2 + 4e^- \rightleftharpoons 4OH^-$	+ 0,40
$Cu^{2+} + 2e^- \rightleftharpoons Cu$	+ 0,34
$SO_4^{2-} + 4H^+ + 2e^- \rightleftharpoons SO_2(g) + 2H_2O$	+ 0,17
$Cu^{2+} + e^- \rightleftharpoons Cu^+$	+ 0,16
$Sn^{4+} + 2e^- \rightleftharpoons Sn^{2+}$	+ 0,15
$S + 2H^+ + 2e^- \rightleftharpoons H_2S(g)$	+ 0,14
$2H^+ + 2e^- \rightleftharpoons H_2(g)$	0,00
$Fe^{3+} + 3e^- \rightleftharpoons Fe$	- 0,06
$Pb^{2+} + 2e^- \rightleftharpoons Pb$	- 0,13
$Sn^{2+} + 2e^- \rightleftharpoons Sn$	- 0,14
$Ni^{2+} + 2e^- \rightleftharpoons Ni$	- 0,27
$Co^{2+} + 2e^- \rightleftharpoons Co$	- 0,28
$Cd^{2+} + 2e^- \rightleftharpoons Cd$	- 0,40
$Cr^{3+} + e^- \rightleftharpoons Cr^{2+}$	- 0,41
$Fe^{2+} + 2e^- \rightleftharpoons Fe$	- 0,44
$Cr^{3+} + 3e^- \rightleftharpoons Cr$	- 0,74
$Zn^{2+} + 2e^- \rightleftharpoons Zn$	- 0,76
$2H_2O + 2e^- \rightleftharpoons H_2(g) + 2OH^-$	- 0,83
$Cr^{2+} + 2e^- \rightleftharpoons Cr$	- 0,91
$Mn^{2+} + 2e^- \rightleftharpoons Mn$	- 1,18
$Al^{3+} + 3e^- \rightleftharpoons Al$	- 1,66
$Mg^{2+} + 2e^- \rightleftharpoons Mg$	- 2,36
$Na^+ + e^- \rightleftharpoons Na$	- 2,71
$Ca^{2+} + 2e^- \rightleftharpoons Ca$	- 2,87
$Sr^{2+} + 2e^- \rightleftharpoons Sr$	- 2,89
$Ba^{2+} + 2e^- \rightleftharpoons Ba$	- 2,90
$Cs^+ + e^- \rightleftharpoons Cs$	- 2,92
$K^+ + e^- \rightleftharpoons K$	- 2,93
$Li^+ + e^- \rightleftharpoons Li$	- 3,05

Increasing strength of oxidising agents/Toenemende sterkte van oksidermiddels

Increasing strength of reducing agents/Toenemende sterkte van reduseermiddels



TABLE 4B: STANDARD REDUCTION POTENTIALS
TABEL 4B: STANDAARD-REDUKSIEPOTENSIALE

Increasing strength of oxidising agents/Toenemende sterkte van oksidermiddels

Half-reactions/ <i>Halfreaksies</i>	E^{θ} (V)
$\text{Li}^+ + e^- \rightleftharpoons \text{Li}$	-3,05
$\text{K}^+ + e^- \rightleftharpoons \text{K}$	-2,93
$\text{Cs}^+ + e^- \rightleftharpoons \text{Cs}$	-2,92
$\text{Ba}^{2+} + 2e^- \rightleftharpoons \text{Ba}$	-2,90
$\text{Sr}^{2+} + 2e^- \rightleftharpoons \text{Sr}$	-2,89
$\text{Ca}^{2+} + 2e^- \rightleftharpoons \text{Ca}$	-2,87
$\text{Na}^+ + e^- \rightleftharpoons \text{Na}$	-2,71
$\text{Mg}^{2+} + 2e^- \rightleftharpoons \text{Mg}$	-2,36
$\text{Al}^{3+} + 3e^- \rightleftharpoons \text{Al}$	-1,66
$\text{Mn}^{2+} + 2e^- \rightleftharpoons \text{Mn}$	-1,18
$\text{Cr}^{2+} + 2e^- \rightleftharpoons \text{Cr}$	-0,91
$2\text{H}_2\text{O} + 2e^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-$	-0,83
$\text{Zn}^{2+} + 2e^- \rightleftharpoons \text{Zn}$	-0,76
$\text{Cr}^{3+} + 3e^- \rightleftharpoons \text{Cr}$	-0,74
$\text{Fe}^{2+} + 2e^- \rightleftharpoons \text{Fe}$	-0,44
$\text{Cr}^{3+} + e^- \rightleftharpoons \text{Cr}^{2+}$	-0,41
$\text{Cd}^{2+} + 2e^- \rightleftharpoons \text{Cd}$	-0,40
$\text{Co}^{2+} + 2e^- \rightleftharpoons \text{Co}$	-0,28
$\text{Ni}^{2+} + 2e^- \rightleftharpoons \text{Ni}$	-0,27
$\text{Sn}^{2+} + 2e^- \rightleftharpoons \text{Sn}$	-0,14
$\text{Pb}^{2+} + 2e^- \rightleftharpoons \text{Pb}$	-0,13
$\text{Fe}^{3+} + 3e^- \rightleftharpoons \text{Fe}$	-0,06
$2\text{H}^+ + 2e^- \rightleftharpoons \text{H}_2(\text{g})$	0,00
$\text{S} + 2\text{H}^+ + 2e^- \rightleftharpoons \text{H}_2\text{S}(\text{g})$	+0,14
$\text{Sn}^{4+} + 2e^- \rightleftharpoons \text{Sn}^{2+}$	+0,15
$\text{Cu}^{2+} + e^- \rightleftharpoons \text{Cu}^+$	+0,16
$\text{SO}_4^{2-} + 4\text{H}^+ + 2e^- \rightleftharpoons \text{SO}_2(\text{g}) + 2\text{H}_2\text{O}$	+0,17
$\text{Cu}^{2+} + 2e^- \rightleftharpoons \text{Cu}$	+0,34
$2\text{H}_2\text{O} + \text{O}_2 + 4e^- \rightleftharpoons 4\text{OH}^-$	+0,40
$\text{SO}_2 + 4\text{H}^+ + 4e^- \rightleftharpoons \text{S} + 2\text{H}_2\text{O}$	+0,45
$\text{Cu}^+ + e^- \rightleftharpoons \text{Cu}$	+0,52
$\text{I}_2 + 2e^- \rightleftharpoons 2\text{I}^-$	+0,54
$\text{O}_2(\text{g}) + 2\text{H}^+ + 2e^- \rightleftharpoons \text{H}_2\text{O}_2$	+0,68
$\text{Fe}^{3+} + e^- \rightleftharpoons \text{Fe}^{2+}$	+0,77
$\text{NO}_3^- + 2\text{H}^+ + e^- \rightleftharpoons \text{NO}_2(\text{g}) + \text{H}_2\text{O}$	+0,80
$\text{Ag}^+ + e^- \rightleftharpoons \text{Ag}$	+0,80
$\text{Hg}^{2+} + 2e^- \rightleftharpoons \text{Hg}(\ell)$	+0,85
$\text{NO}_3^- + 4\text{H}^+ + 3e^- \rightleftharpoons \text{NO}(\text{g}) + 2\text{H}_2\text{O}$	+0,96
$\text{Br}_2(\ell) + 2e^- \rightleftharpoons 2\text{Br}^-$	+1,07
$\text{Pt}^{2+} + 2e^- \rightleftharpoons \text{Pt}$	+1,20
$\text{MnO}_2 + 4\text{H}^+ + 2e^- \rightleftharpoons \text{Mn}^{2+} + 2\text{H}_2\text{O}$	+1,23
$\text{O}_2(\text{g}) + 4\text{H}^+ + 4e^- \rightleftharpoons 2\text{H}_2\text{O}$	+1,23
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6e^- \rightleftharpoons 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+1,33
$\text{Cl}_2(\text{g}) + 2e^- \rightleftharpoons 2\text{Cl}^-$	+1,36
$\text{MnO}_4^- + 8\text{H}^+ + 5e^- \rightleftharpoons \text{Mn}^{2+} + 4\text{H}_2\text{O}$	+1,51
$\text{HClO}_2 + 2\text{H}^+ + 2e^- \rightleftharpoons 2\text{H}_2\text{O}$	+1,77
$\text{Co}^{3+} + e^- \rightleftharpoons \text{Co}^{2+}$	+1,81
$\text{F}_2(\text{g}) + 2e^- \rightleftharpoons 2\text{F}^-$	+2,87

Increasing strength of reducing agents/Toenemende sterkte van reduseermiddels

