

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za





education

Department:
Education
North West Provincial Government
REPUBLIC OF SOUTH AFRICA

PROVINCIAL ASSESSMENT

GRADE 12

MATHEMATICS P1

JUNE 2026

MARKS: 150

TIME: 3 hours

This question paper consists of 10 pages, 1 information sheet
and an answer book of 18 pages.

Proudly South African



INSTRUCTIONS AND INFORMATION

Read the following instructions carefully before answering the questions.

1. This question paper consists of 11 questions.
2. Answer ALL the questions in the SPECIAL ANSWER BOOK provided.
3. Clearly show ALL calculations, diagrams, graphs, etc. that you have used in determining your answers.
4. Answers only will NOT necessarily be awarded full marks.
5. You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.
6. If necessary, round off answers to TWO decimal places, unless stated otherwise.
7. Diagrams are NOT necessarily drawn to scale.
8. An information sheet with formulae is included at the end of the question paper.
9. Write neatly and legibly.



SA EXAM PAPERS

Proudly South African

**QUESTION 1**1.1 Solve for x :

1.1.1 $x^2 + x - 30 = 0$ (3)

1.1.2 $2x^2 - 8 = 5x$ (correct to TWO decimal places) (4)

1.1.3 $-7x^2 < -14x$ (4)

1.1.4 $\sqrt{x+6} - x = 4$ (5)

1.1.5 $4^x - 16 = 6.2^x$ (4)

1.2 Solve for x and y simultaneously:

$x + 2y + 5 = 0$ and $y^2 + x^2 + 10x + 5 = 0$ (6)

1.3 Given the irrational number $\frac{16}{\sqrt{2}}$:

1.3.1 Show that $\frac{16}{\sqrt{2}}$ can be written as $8\sqrt{2}$. (1)

1.3.2 Hence, show that $\frac{16}{\sqrt{2}} = 8a + 8b$ if $(a+b) > 0$ and $(a+b)^2 = 2$. (3)

[30]**QUESTION 2**

The sequence of the first differences of a quadratic pattern is:

107; 113; 119; ...

The third term of the quadratic number pattern is 320.

2.1 Write down the first four terms of the quadratic number pattern. (3)

2.2 Determine the general term (T_n) of the quadratic number pattern. (4)

2.3 Between which two consecutive terms of the quadratic number pattern will the first difference be 2 021? (4)

[11]**SA EXAM PAPERS**

Proudly South African

**QUESTION 3**

3.1 Given the arithmetic series: $6 + 13 + 20 + 27 + \dots$

3.1.1 Determine the value of the 24th term. (3)

3.1.2 Calculate the sum of the first 50 terms of the series. (2)

3.2 Find the value of m :

$$\sum_{k=1}^m 12 \left(\frac{1}{3} \right)^{k-1} = \frac{1456}{81} \quad (5)$$

3.3 The first two terms of a geometric series are $x + 2$ and $x^2 - 2x - 8$.

3.3.1 For which values of x will the series converge? (2)

3.3.2 If the sum to infinity of the series is $\frac{11}{3}$, determine the value of x . (4)

[16]

QUESTION 4

Prove, WITHOUT USING A FORMULA, that:


$$p + \frac{p}{2} + \frac{p}{4} + \frac{p}{8} + \dots + \frac{p}{2^n} = p \left(2 - \frac{1}{2^n} \right) \quad [5]$$



**QUESTION 5**

In the diagram below, the graphs of $f(x) = ax^2 + bx + c$ and $g(x) = -4x - 12$, are drawn.

- B and C are the x -intercepts of f .
- C is the x -intercept of g .
- f and g intersect at A and C.
- A is the turning point of f .
- The line $x = -1$ is the axis of symmetry of f .



- 5.1 Determine the coordinates of A, B and C. (4)
- 5.2 Show that $f(x) = 2x^2 + 4x - 6$. (3)
- 5.3 Write down the coordinates of the turning point of $f(x - 3) - 2$. (2)
- 5.4 Write down the value of k if $f(x) = k$ has equal roots. (1)
- 5.5 Write down the range for $-f(x)$. (2)
- 5.6 If $f(x) = 2x^2 + 4x - 6$, determine the length of OT if PRT is perpendicular to the x -axis with P a point on f , R is a point on g and $PRT = 16$ units. (6)

SA EXAM PAPERS

Proudly South African


[18]

**QUESTION 6**

Sketched below are the graphs of $f(x) = 3 \cdot 2^{x-p} + q$ and $g(x) = \frac{a}{x+k} + c$.

The asymptotes of g are $x = 2$ and $y = 1$.

The asymptote of f is $y = 1$.



6.1 Determine the equation of g . (4)

6.2 Determine the values of p and q . (4)

6.3 For which values of x is $f(x) \leq g(x)$? (2)


6.4 Determine the equation of the axis of symmetry of h for $m < 0$ if $h(x) = g(x) + 2$. (3)

[13]**SA EXAM PAPERS**

Proudly South African

**QUESTION 7**

Given: $f(x) = \log_b x$. A is the x -intercept of f and point $(125; -3)$ is on f .



7.1 Determine the value of b . (3)

7.2 Write down the coordinates of A. (2)

7.3 For which value(s) of x is $f(x) \geq -3$? (2)

7.4 Write down the equation of f^{-1} , the inverse of f , in the form $y = \dots$ (2)

7.5 Sketch the graph of f^{-1} on the set of axes provided in the ANSWER BOOK. Clearly indicate the y -intercept with the axes, at least the coordinates of ONE other point and the asymptote on your graph. (3)

7.6 $h(x)$ is the reflection of f^{-1} about the x -axis. Write down the equation of $h(x)$. (1)

[13]**QUESTION 8**

8.1 Determine $f'(x)$ from first principles if $f(x) = 3 - 2x^2$. (5)

8.2 Determine:

8.2.1 $f'(x)$ if it is given that $f(x) = -x^5 + 3x^4$ (2)

8.2.2 $\frac{dy}{dx}$ if $yx = 7x^3 - 5x^{\frac{2}{3}}$ (4)

**SA EXAM PAPERS**

Proudly South African



8.3 The tangent to the curve of $g(x) = 2x^3 + px^2 + qx - 7$ at $x = 1$ has the equation $y = 5x - 8$.


8.3.1 Show that $(1; -3)$ is the point of contact of the tangent to the graph of g . (1)

8.3.2 Hence, calculate the values of p and q . (6)
[18]

QUESTION 9

Sketched below is the graph of $f(x) = -x^3 + ax^2 + bx + c$.

- P and Q are the turning points of f .
- $x = -2$; $x = 3$ and $x = 6$ are the x -intercepts of f .



9.1 Show that $a = 7$; $b = 0$ and $c = -36$. (2)

9.2 Calculate the coordinates of the turning points P and Q. (5)

9.3 Determine the x -coordinate of the point of inflection. (3)

9.4 For which values of x will $x \cdot f'(x) > 0$? (2)

9.5 For which value of k will $f(x) - 4 = k$ have one root? (2)


[14]



**QUESTION 10**

The graph of $f'(x)$ is drawn below.

- 6 is the y -intercept of f' .
- 1 and 2 are the x -intercepts and $\left(\frac{3}{2}; -\frac{3}{4}\right)$ is the turning point of f' respectively.



- 10.1 Write down the x -coordinate of the minimum turning point of $f(x)$. (1)
- 10.2 For which values of x will $f(x)$ be decreasing? (2)
- 10.3 Write down the gradient of the tangent to f if $x = 0$. (1)
- 10.4 At which other x -value will a tangent to f be parallel to the answer in QUESTION 10.3? (1)
- 10.5 Is $f(x)$ concave up or concave down for $x > \frac{3}{2}$? (1)

[6]**SA EXAM PAPERS**

Proudly South African

**QUESTION 11**

A body moves according to the formulae $S(t) = 100t^2 - 200t + 300$, where S is the distance in meters, from a fixed point P , after t seconds.

11.1 Determine the distance from P after 10 seconds. (2)

11.2 Determine after how many seconds the body will reach a speed of 600 m/s. (2)

11.3 After how many seconds will the body reach its minimum distance from P ? (2)

[6]**TOTAL: 150****SA EXAM PAPERS**

Proudly South African



INFORMATION SHEET

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$A = P(1+i)n$$

$$A = P(1-in)$$

$$A = P(1-i)^n$$

$$A = P(1+i)^n$$

$$T_n = a + (n-1)d$$

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(r^n - 1)}{r - 1}; r \neq 1$$

$$S_\infty = \frac{a}{1-r}; -1 < r < 1$$

$$F = \frac{x[(1+i)^n - 1]}{i}$$

$$P = \frac{x[1 - (1+i)^{-n}]}{i}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$$

$$y = mx + c$$

$$y - y_1 = m(x - x_1)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \tan \theta$$

$$(x-a)^2 + (y-b)^2 = r^2$$

$$\text{In } \triangle ABC: \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\text{area } \triangle ABC = \frac{1}{2} ab \sin C$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\cos 2\alpha = \begin{cases} \cos^2 \alpha - \sin^2 \alpha \\ 1 - 2\sin^2 \alpha \\ 2\cos^2 \alpha - 1 \end{cases}$$

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha$$

$$\bar{x} = \frac{\sum fx}{n}$$

$$\sigma^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}$$

$$P(A) = \frac{n(A)}{n(S)}$$

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

$$\hat{y} = a + bx$$

$$b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$$



SA EXAM PAPERS

Proudly South African