

SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za





This Paper was downloaded from SAEXAMPAPERS

GAUTENG PROVINCE

EDUCATION
REPUBLIC OF SOUTH AFRICA

**JUNE EXAMINATION/
JUNIE EKSAMEN
GRADE/*GRAAD* 12**

2026

**MARKING GUIDELINES/
*NASIENRIGLYNE***

**MATHEMATICS/
*WISKUNDE***

(PAPER/*VRAESTEL* 2)

27 pages/*bladsye*



SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE****MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)****GR12 0626****NOTE:**

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- If a candidate has crossed out an attempt of a question and not redone the question, mark the crossed-out version.
- Consistent accuracy applies in ALL aspects of the marking guidelines. Stop marking at the second calculation error.
- Assuming answers/values to solve a problem is NOT acceptable.

LET WEL:

- *As 'n kandidaat 'n vraag TWEE KEER beantwoord het, sien slegs die EERSTE poging na.*
- *As 'n kandidaat 'n antwoord van 'n vraag doodtrek en dit nie oordoen nie, sien die doodgetrekte poging na.*
- *Volgehoue akkuraatheid word in ALLE aspekte van die nasienriglyne toegepas. Hou op nasien by die tweede berekeningsfout.*
- *Aannames van antwoorde/waardes om 'n probleem op te los, word NIE toegelaat NIE.*

GEOMETRY/MEETKUNDE	
S	A mark for a correct statement (A statement mark is independent of a reason.)
	<i>'n Punt vir 'n korrekte bewering (n Punt vir 'n bewering is onafhanklik van 'n rede.)</i>
R	A mark for a correct reason (A reason mark may only be awarded if the statement is correct.)
	<i>'n Punt vir 'n korrekte rede (n Punt word slegs vir die rede toegeken as die bewering korrek is.)</i>
S/R	Award a mark if the statement AND reason are both correct.
	<i>Ken 'n punt toe indien beide die bewering EN rede korrek is.</i>

**SA EXAM PAPERS**

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**

**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**

GR12 0626

AMENDMENT TO MARKING GUIDELINES

JUNE 2026 PROVINCIAL COMMON EXAMINATION

FOR ATTENTION: THE CHIEF INVIGILATOR

SUBJECT / VAK	MATHEMATICS/WISKUNDE
PAPER / VRAESTEL	2
DATE OF EXAMINATION	8 JUNE/JUNIE 2026

The errata for the Marking Guidelines of **MATHEMATICS P2/WISKUNDE V2** has reference.

There was an error in **QUESTION 2.4** which affected the possible candidate responses on BOTH the English AND Afrikaans versions of the question paper. This matter was addressed at the Marking Standardisation Meeting.

To ensure that candidates are not disadvantaged nor prejudiced in any way, you are advised to ask your Mathematics Educator to please **ignore QUESTION 2.4** when marking. This question carries 5 marks.

In other words, the paper must be marked out of a total of 145 instead of 150 and then the learners' marks must be converted to a mark out of 150. E.g. Should a learner attain $\frac{29}{145}$ then that mark is recalculated as $\frac{30}{150}$.

Use the formula: $\frac{a}{145} \times 100 = b$. Then, $\frac{b}{100} \times 150 = c$

C is the mark that is entered into SASAMS out of 150.

MR JONATHAN WILLIAMS


DIRECTOR: EXAMINATIONS MANAGEMENT

9 JUNE 2026



SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626
QUESTION/VRAAG 1


1.1	$m_{MT} = \frac{-1-1}{-1-2}$ $= \frac{2}{3}$	✓ substitution into gradient formula/ <i>substitusie in die gradiënt formule</i> ✓ answer/antwoord (2)
1.2	$y = \frac{2}{3}x + c$ $\frac{11}{2} = \frac{2}{3}(-1) + c$ $c = \frac{37}{6}$ $\therefore y = \frac{2}{3}x + \frac{37}{6}$ OR/OF $y - \frac{11}{2} = \frac{2}{3}(x - (-1))$ $\therefore y = \frac{2}{3}x + \frac{37}{6}$	✓ $m_{\text{line/lyn}} = \frac{2}{3}$ ✓ substitution into equation formula/ <i>substitusie in die vergelykings- formule</i> ✓ answer/antwoord (3)
1.3	$\frac{-1+k}{2} = 2$ $\therefore k = 5$ $\frac{-1+h}{2} = 1$ $\therefore h = 3$ Answer only: Full marks <i>Slegs antwoord: Volpunte</i>	✓ substitution into midpoint formula/ <i>substitusie in die middelpuntsformule</i> ✓ $k = 5$ ✓ $h = 3$ (3)



SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**

**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**

GR12 0626



1.4	$AM = \sqrt{(-1 - 5)^2 + (-1 - 3)^2}$ $= 2\sqrt{13} \text{ units/eenhede (Accept } \sqrt{52})$ OR/OF $MT = \sqrt{(-1 - 2)^2 + (-1 - 1)^2} = \sqrt{13}$ $\therefore AM = 2MT = 2\sqrt{13} \text{ units/eenhede}$	✓ substitution into distance formula/ substitusie in afstandformule ✓ answer in surd form/ antwoord in wortelvorm (2) ✓ substitution into distance formula/ substitusie in afstandformule ✓ answer in surd form/ antwoord in wortelvorm (2)
1.5	ARMP is a parallelogram/is 'n parallelogram (2 pairs of opp sides / 2 pare oorst sye) $\frac{x_R + (-1)}{2} = 2$ $x_R = 5$ $\frac{y_R + \frac{11}{2}}{2} = 1$ $y_R = -\frac{7}{2}$ Answer only: Full marks Slegs antwoord: Volpunte Translation implied Translasie geïmpliseer $\therefore R(5; -\frac{7}{2})$	✓ method/metode ✓ $x_R = 5$ ✓ $y_R = -\frac{7}{2}$ (3)



SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626

1.6

$$m_{AM} = \frac{2}{3}$$

$$\alpha = \tan^{-1}\left(\frac{2}{3}\right)$$

$$\alpha = 33,69^\circ$$

$$m_{PR} = \frac{\frac{11}{2} - 1}{-1 - 2} = -\frac{3}{2}$$

$$\theta = \tan^{-1}\left(-\frac{3}{2}\right) + 180$$

$$\theta = 123,69^\circ$$

$$M\hat{T}R = 123,69^\circ - 33,69^\circ = 90^\circ$$

$$\therefore P\hat{T}A = 90^\circ \text{ (vert opp } \angle\text{s)/(regoorst } \angle\text{e)}$$

OR/OF

$$m_{AM} = \frac{2}{3}$$

$$m_{PR} = \frac{\frac{11}{2} - 1}{-1 - 2} = -\frac{3}{2}$$

$$m_{AM} \times m_{PR} = \frac{2}{3} \times -\frac{3}{2} = -1$$

$$\therefore AM \perp PR$$

$$\therefore P\hat{T}A = 90^\circ$$

OR/OF

$$PT = \sqrt{(1-2)^2 + \left(\frac{11}{2} - 1\right)^2} = \frac{3\sqrt{13}}{2} = 5,41$$

$$AT = \sqrt{(5-2)^2 + (3-1)^2} = \sqrt{13} = 3,61$$

$$AP = \sqrt{(5 - (-1))^2 + \left(3 - \frac{11}{2}\right)^2} = 6,5$$

$$P\hat{T}A = \cos^{-1}\left(\frac{\left(\frac{3\sqrt{13}}{2}\right)^2 + (\sqrt{13})^2 - \left(\frac{13}{2}\right)^2}{2\left(\frac{3\sqrt{13}}{2}\right)(\sqrt{13})}\right)$$

$$\therefore P\hat{T}A = 90^\circ$$

$$\checkmark m_{AM} = \frac{2}{3}$$

$$\checkmark \alpha = 33,69^\circ$$

$$\checkmark m_{PR} = -\frac{3}{2}$$

$$\checkmark \theta = 123,69^\circ$$

$$\checkmark \text{answer/antwoord} \quad (5)$$

OR/OF

$$\checkmark m_{AM} = \frac{2}{3}$$

$$\checkmark m_{PR} = -\frac{3}{2}$$

$$\checkmark m_{AM} \times m_{PR} = -1$$

$$\checkmark AM \perp PR$$

$$\checkmark \text{answer/antwoord} \quad (5)$$

OR/OF

$$\checkmark PT = \frac{3\sqrt{13}}{2}$$

$$\checkmark AT = \sqrt{13}$$

$$\checkmark AP = \frac{13}{2}$$

$$\checkmark \text{substitution into cosine formula/}$$

$$\text{substitution in cos-formule}$$

$$\checkmark \text{answer/antwoord} \quad (5)$$


SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626

1.7	$m_{AP} = \frac{\frac{11}{2}-3}{-1-5} = -\frac{5}{12}$ $m_{AM} = \frac{2}{3}$ $m_{AP} \times m_{AM} = -\frac{5}{12} \times \frac{2}{3} \neq -1$ $\therefore \widehat{P\hat{A}M} \neq 90^{\circ}$	$\checkmark m_{AP} = -\frac{5}{12}$ $\checkmark m_{AM} = \frac{2}{3}$ $\checkmark m_{AP} \times m_{AM} \neq -1$	(3)	
1.8		$h = 3 \text{ units/eenhede}$ $MP = \frac{11}{2} - (-1) = \frac{13}{2}$ $= 6,5 \text{ units/eenhede}$ $\text{Area of /opp } \Delta MPT = \frac{1}{2} \times \frac{13}{2} \times 3 \text{ units}^2/\text{eenhede}^2$ $= \frac{39}{4} \text{ units}^2/\text{eenhede}^2$ $= 9,75 \text{ units}^2/\text{eenhede}^2$	$\checkmark h = 3 \text{ units}/h = 3 \text{ eenhede}$ $\checkmark \text{length of MP/}$ lengte van MP $\checkmark \text{substitution into area formula}$ $\text{/substitusie in oppervlakformule}$ $\checkmark \text{answer/antwoord}$	(4)


SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626

OR/OF $\widehat{PTM} = 90^\circ$ (<s on a str line) / (< e op 'n reguitlyn) $PT = \sqrt{(-1 - 2)^2 + \left(\frac{11}{2} - 1\right)^2} = \frac{3\sqrt{13}}{2} = 5,41$ $MT = \sqrt{(-1 - 2)^2 + (-1 - 1)^2} = \sqrt{13} = 3,61$ Area of / (opp van) ΔMPT $= \frac{1}{2} \times \frac{3\sqrt{13}}{2} \times \sqrt{13} \text{ units}^2/\text{eenhede}^2$ $= \frac{39}{4} \text{ units}^2/\text{eenhede}^2$ $= 9,75 \text{ units}^2/\text{eenhede}^2$ OR/OF $\widehat{PTM} = 90^\circ$ (<s on a str line) / (< e op 'n reguitlyn) $PT = \sqrt{(1 - 2)^2 + \left(\frac{11}{2} - 1\right)^2} = \frac{3\sqrt{13}}{2} = 5,41$ $MT = \sqrt{(-1 - 2)^2 + (-1 - 1)^2} = \sqrt{13} = 3,61$ Area of/Opp van ΔMPT $= \frac{1}{2} \times \frac{3\sqrt{13}}{2} \times \sqrt{13} \sin(90^\circ) \text{ units}^2/\text{eenhede}^2$ $= \frac{39}{4} \text{ units}^2/\text{eenhede}^2$ $= 9,75 \text{ units}^2/\text{eenhede}^2$	OR/OF $\checkmark PT = \frac{3\sqrt{13}}{2}$ $\checkmark MT = \sqrt{13}$ $\checkmark \text{ substitution into area formula}$ $\text{substitusie in opp. formule}$ $\checkmark \text{ answer/antwoord} \quad (4)$ OR/OF $\checkmark PT = \frac{3\sqrt{13}}{2}$ $\checkmark MT = \sqrt{13}$ $\checkmark \text{ substitution into area formula}$ $\text{substitusie in opp. formule}$ $\checkmark \text{ answer/antwoord} \quad (4)$
[25]	


SA EXAM PAPERS


Proudly South African

MARKING GUIDELINES/
NASIENRIGLYNE

MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)

GR12 0626

QUESTION/VRAAG 2



2.1.1	$ON = \sqrt{(6 - 0)^2 + (-8 - 0)^2}$ $ON = \sqrt{100}$ $\therefore ON = 10 \text{ units/eenhede}$	✓ substitution/substitusie ✓ answer/antwoord (2)
2.1.2	$ON = OS$ tangents from the same point/ <i>raaklyne vanuit dieselfde punt</i> $\therefore p = 10$	✓ S ✓ answer/antwoord (2)
2.1.3	$ON \perp NM$ tangent $\perp$ radius/ <i>raaklyn $\perp$ radius</i> $m_{ON} = \frac{-8-0}{6-0}$ $\therefore m_{ON} = -\frac{4}{3}$ $\therefore m_{NM} = \frac{3}{4}$	✓ $-\frac{4}{3}$ ✓ $\frac{3}{4}$ (2)



SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626

2.1.4	$m_{NM} = \frac{3}{4}$ $\frac{q - (-8)}{10 - 6} = \frac{3}{4}$ $\frac{q + 8}{4} = \frac{3}{4}$ $q = -5$	$y = \frac{3}{4}x + c$ $-8 = \frac{3}{4}(6) + c$ $c = -\frac{25}{2}$ $y = \frac{3}{4}x - \frac{25}{2}$ $q = \frac{3}{4}(10) - \frac{25}{2}$ $q = -5$	✓ equating and simplifying/ gelykstelling en vereenvoudiging ✓ answer/antwoord OR/OF ✓ substitution/vervanging ✓ answer/antwoord (2)
2.2	MS = r = 5 units/eenhede $(x - 10)^2 + (y + 5)^2 = 25$		✓ radius ✓ subs centre coordinates/ subst middelpuntkoördinate ✓ equation/vergelyking (3)
2.3	$(x - 10)^2 + (-5 + 5)^2 = 25$ $(x - 10)^2 = 25$ $\therefore k = 5$ or/of $k = 15$ $k = 5$ or/of $k = 15$		✓ substitution/vervanging ✓ both answer/beide antwoord (2)
2.4	Coordinates of the point directly opposite N is C. / Koördinate van die punt direk teenoor N is C. $\frac{x+6}{2} = 10$ $\frac{y-8}{2} = -5$ $x + 6 = 10 \times 2$ $y - 8 = -5 \times 2$ $\therefore x = 14$ $\therefore y = -2$ $\therefore C(14; -2)$ equation of the tangent at C/ vergelyking van raaklyn by C $y + 2 = -\frac{4}{3}(x - 14)$ $\therefore y = -\frac{4}{3}x + \frac{50}{3}$ $\therefore 0 < t < \frac{50}{3}$	DO NOT MARK THIS QUESTION	✓ substitution into correct midpoint formula/subst in korrekte middelpuntsformule ✓ x – value/waarde ✓ y – value/waarde ✓ equation of tangent at C/ vergelyking van raaklyn by C ✓ value of t/waarde van t (5)


SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626

2.5	They will not touch./<i>Hulle sal nie raak nie</i> The new circle is the old circle shifted up by 11 units/ <i>Die nuwe sirkel is die ou sirkel wat met 11 eenhede opwaarts gekuif het.</i> OR/OF sum of radii/<i>som van radiusse</i> = 10 distance between centres/<i>afstand tussen middelpunte</i> $= \sqrt{(10 - 10)^2 + (6 + 5)^2}$ $= 11$ $\therefore$ the circles do not touch or cut/<i>die sirkels raak of sny nie</i>	✓ answer/<i>antwoord</i> ✓ any valid reason/ <i>enige geldige rede</i> (2) ✓ distance between centres > sum of radii /<i>afstand tussen middelpunte > som van radiusse</i> ✓ the circles do not touch or cut/<i>die sirkels raak of sny nie.</i> (2)
[15]		


SA EXAM PAPERS


Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**

**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**

GR12 0626

QUESTION/VRAAG 3

3.1			
3.1.1	$\tan A = \frac{2q}{q}$ $\therefore \tan A = 2$ OR/OF $\frac{\sin A}{\cos A} = \frac{2q}{q} = 2 = \tan A$	✓ correct diagram/ korrekte diagram ✓ answer/antwoord (2) OR/OF ✓ quotient identity/ kwosiënt identiteit ✓ answer/antwoord (2)	
3.1.2	$(2q)^2 + q^2 = 1$ $5q^2 = 1$ $q^2 = \frac{1}{5}$ $q = \frac{1}{\sqrt{5}}$ $q = \frac{\sqrt{5}}{5}$	✓ Pythagoras ✓ simplification/ vereenvoudiging ✓ answer in any form/ antwoord in enige vorm (3)	



SA EXAM PAPERS

Proudly South African

MARKING GUIDELINES/ NASIENRIGLYNE	MATHEMATICS/WISKUNDE (PAPER/VRAESTEL 2)	GR12 0626
--	--	------------------

3.2	$x^2 + y^2 = r^2$ $\therefore x = \sqrt{1 - (p)^2}$	
3.2.1	$\sin 594^\circ$ $= \sin 234^\circ$ $= -\sin 54^\circ$ $= -p$	$\checkmark -\sin 54^\circ$ $\checkmark -p$ (2)
3.2.2	$\cos 36^\circ = p$ or/of $\cos 36^\circ$ $= \cos (90^\circ - 54^\circ)$ $= \sin 54^\circ$ $= p$	$\checkmark \checkmark$ answer/antwoord (2)
3.2.3	$\cos 18^\circ \sin 2(36^\circ)$ $= \sin 72^\circ$ $= \sin 2(36^\circ)$ $= 2 \sin 36^\circ \cdot \cos 36^\circ$ $= 2 \left(\frac{\sqrt{1-(p)^2}}{1} \right) \left(\frac{p}{1} \right)$ $= 2p\sqrt{1-(p)^2}$ OR/OR $\cos 18^\circ = \cos (54^\circ - 36^\circ)$ $= \cos 54^\circ \cos 36^\circ + \sin 54^\circ \sin 36^\circ$ $= \sqrt{1-p^2} \times p + p \times \sqrt{1-p^2}$ $= 2p\sqrt{1-(p)^2}$	$\checkmark \sin 72^\circ$ $\checkmark \sin 2(36^\circ)$ $\checkmark 2 \sin 36^\circ \cdot \cos 36^\circ$ $\checkmark 2p\sqrt{1-(p)^2}$ (4) OR/OR $\checkmark \cos 18^\circ = \cos (54^\circ - 36^\circ)$ $\checkmark \cos 54^\circ \cos 36^\circ + \sin 54^\circ \sin 36^\circ$ $\checkmark \sqrt{1-p^2} \times p + p \times \sqrt{1-p^2}$ $\checkmark 2p\sqrt{1-(p)^2}$ (4)

**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626

3.3	$= \frac{\cos(90^\circ + 50^\circ) - \cos \theta}{\sin 50^\circ + \cos \theta}$ $= \frac{-\sin 50^\circ - \cos \theta}{\sin 50^\circ + \cos \theta}$ $= \frac{-(\sin 50^\circ + \cos \theta)}{\sin 50^\circ + \cos \theta}$ $= -1$	✓ $-\sin 50^\circ$ ✓ $\cos \theta$ ✓ $\sin 50^\circ$ ✓ $\cos \theta$ ✓ common factor/ gemeenskaplike faktor ✓ answer/antwoord	(6)
-----	--	---	-----


SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626

3.4	$\cos(x + 65^\circ) \cdot \cos(x + 20^\circ) - \sin(x + 245^\circ) \cdot \sin(x + 20^\circ)$ $= \cos(x + 65^\circ) \cdot \cos(x + 20^\circ) + \sin(x + 65^\circ) \cdot \sin(x + 20^\circ)$ $= \cos[(x + 65^\circ) - (x + 20^\circ)]$ $= \cos 45^\circ$ $= \frac{1}{\sqrt{2}} \text{ or/of } \frac{\sqrt{2}}{2}$	✓ reduction/reduksie of herleiding ✓ compound angles/saamgestelde hoeke ✓ $\cos 45^\circ$ ✓ answer/antwoord (4)
3.5	$\cos^2 x - \sin^2 x = \frac{1}{2}$ $\cos 2x = \frac{1}{2}$ ref angle/verwysingshoek: $2x = \cos^{-1}\left(\frac{1}{2}\right)$ $\therefore 2x = 60^\circ + k \cdot 360^\circ$ 1 st Quad/1 ^{ste} Kwad: $2x = 60^\circ + k \cdot 360^\circ$ 4 th Quad/4 ^{de} Kwad: $2x = 360^\circ - 60^\circ + k \cdot 360^\circ$ $\therefore x = 30^\circ + k \cdot 180^\circ, k \in \mathbb{Z}$ $2x = 300^\circ + k \cdot 360^\circ$ $\therefore x = 150^\circ + k \cdot 180^\circ, k \in \mathbb{Z}$ OR/OF $\cos^2 x - \sin^2 x = \frac{1}{2}$ $2 \cos^2 x - 2 \sin^2 x = 1$ $2 \cos^2 x - 2 \sin^2 x = \sin^2 x + \cos^2 x$ $2 \sin^2 x + \sin^2 x = 2 \cos^2 x - \cos^2 x$ $3 \sin^2 x = \cos^2 x$ $\therefore \tan^2 x = \frac{1}{3}$ $\therefore \tan x = \frac{1}{\sqrt{3}}$ $\therefore x = 30^\circ$ #ref angle/ #verw hoek $\therefore x = 30^\circ + k \cdot 180^\circ; k \in \mathbb{Z}$ $x = 180^\circ + 30^\circ + k \cdot 180^\circ k \in \mathbb{Z}$ $x = 210^\circ + k \cdot 180^\circ k \in \mathbb{Z}$	✓ $\cos 2x = \frac{1}{2}$ ✓ $2x = 60^\circ + k \cdot 360^\circ$ ✓ $x = 30^\circ + k \cdot 180^\circ, k \in \mathbb{Z}$ (4) ✓ $x = 150^\circ + k \cdot 180^\circ, k \in \mathbb{Z}$ OR/OF ✓ multiplying by 2 and using identity/vermenigvuldig met 2 en gebruik identiteit ✓ $3 \sin^2 x = \cos^2 x$ ✓ $\tan x = \frac{1}{\sqrt{3}}$ ✓ answer/antwoord (4)


SA EXAM PAPERS

Proudly South African


**MARKING GUIDELINES/
NASIENRIGGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626

3.6	$\frac{\sin 2\theta \cdot \tan \theta}{\cos 2\theta + 1} = \tan^2 \theta$		
3.6.1	$\text{LHS} = \frac{\sin 2\theta \cdot \tan \theta}{\cos 2\theta + 1}$ $= \frac{2 \sin \theta \cos \theta \cdot \frac{\sin \theta}{\cos \theta}}{2 \cos^2 \theta - 1 + 1}$ $= \frac{2 \sin^2 \theta}{2 \cos^2 \theta}$ $= \tan^2 \theta$ $\therefore \text{LHS} = \text{RHS/LK} = \text{RK}$	$\checkmark 2 \sin \theta \cos \theta$ $\checkmark \frac{\sin \theta}{\cos \theta}$ $\checkmark 2 \cos^2 \theta - 1$ $\checkmark \frac{2 \sin^2 \theta}{2 \cos^2 \theta}$ (4)	
3.6.2	$\cos 2\theta + 1 = 0$ $\cos 2\theta = -1$ $2\theta = 180^\circ$ $\theta = 90^\circ$ If only $\theta = 90^\circ$ for $\tan \theta$ undefined ($\frac{1}{4}$) /as slegs $\theta = 90^\circ$ vir $\tan \theta$ ongedefinieërd $\tan \theta (\frac{1}{4})$	$\checkmark \text{denominator/noemer} = 0$ $\checkmark \cos 2\theta = -1$ $\checkmark 2\theta = 180^\circ$ $\checkmark \theta = 90^\circ$ (4)	
[35]			


SA EXAM PAPERS

Proudly South African

QUESTION/VRAAG 4



4.1	$\Delta ABC \equiv \Delta ABD$ (S.A.S)/(S $\angle$ S)	✓ R (1)
4.2	$\cos \alpha = \frac{x}{AC}$ $\therefore AC = \frac{x}{\cos \alpha}$	✓ correct ratio/korrekte verhouding ✓ AC simplified as subject/AC vereenvoudig as onderwerp (2)
4.3	$CD^2 = \left(\frac{x}{\cos \alpha}\right)^2 + \left(\frac{x}{\cos \alpha}\right)^2 - 2\left(\frac{x}{\cos \alpha}\right)\left(\frac{x}{\cos \alpha}\right)\cos(180^\circ - 2\beta)$ $CD^2 = \frac{2x^2}{\cos^2 \alpha} - 2\left(\frac{x^2}{\cos^2 \alpha}\right)\cos 2\beta$ $CD^2 = \frac{2x^2}{\cos^2 \alpha}(1 + \cos 2\beta)$ $CD^2 = \frac{2x^2}{\cos^2 \alpha}(1 + (1 - 2\sin^2 \beta))$ $CD^2 = \frac{2x^2}{\cos^2 \alpha}(1 + (1 - 2(1 - \cos^2 \beta)))$ $CD^2 = \frac{2x^2}{\cos^2 \alpha}(2\cos^2 \beta)$ $CD^2 = \frac{4x^2 \cos^2 \beta}{\cos^2 \alpha}$ $\therefore CD = \frac{2x \cos \beta}{\cos \alpha}$	✓ correct use of cos rule/korrekte gebruik van cos-reël ✓ simplification/vereenvoudig ✓ applying double angles/gebruik dubbelhoeke ✓ identity/identiteit (4)



**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626

	OR/OF $\widehat{ADC} = \beta$ $\widehat{CAD} = 180^\circ - 2\beta$ $\frac{CD}{\sin(180^\circ - 2\beta)} = \frac{AC}{\sin \beta}$ $CD = \frac{AC \cdot \sin 2\beta}{\sin \beta}$ but $AC = \frac{x}{\cos \alpha}$ $CD = \frac{\left(\frac{x}{\cos \alpha}\right) \cdot \sin 2\beta}{\sin \beta}$ $CD = \frac{x \cdot 2 \sin \beta \cdot \cos \beta}{\cos \alpha \cdot \sin \beta}$ $\therefore CD = \frac{2x \cdot \cos \beta}{\cos \alpha}$	OR/OF ✓ $\widehat{ADC} = \beta$ and/en $\widehat{CAD} = 180^\circ - 2\beta$ ✓ correct use of sin rule/ <i>korrekte gebruik van sin-reël</i> ✓ substituting AC/ <i>substitusie van AC</i> ✓ double angle/dubbelhoek (4)
4.4	$CD = \frac{2(25) \cos 65,62^\circ}{\cos 30^\circ}$ $CD = 23,83 \text{ cm}$ penalize 1 mark for incorrect rounding /penaliseer 1 punt vir foutiewe afronding	✓ substitution/substitusie ✓ answer/antwoord (2)
[9]		


SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626
QUESTION/VRAAG 5


5.1	$f(180^\circ) = \cos(180^\circ - 45^\circ)$ $= -0,71$ ANSWER ONLY: FULL MARKS SLEGS ANTWOORD: VOLPUNTE $\therefore -0,71 \leq y \leq 1$ OR/OF $\therefore y \in [-0,71; 1]$ OR/OF $-\frac{\sqrt{2}}{2} \leq y \leq 1$	$\checkmark f(180^\circ)$ $\checkmark$ answer/antwoord (2)
5.2	SEE THE GRAPH BELOW AND ALLOCATE MARKS ACCORDINGLY: SIEN DIE GRAFIEK HIERONDER EN KEN PUNTE TOE DAARVOLGENS: $\checkmark$ intercepts/afsnitte $\checkmark$ shape/vorm $\checkmark$ turning points/draaipunte (3)	
5.3	Period/periode = 180°	$\checkmark$ answer/antwoord (1)
5.4	$-45^\circ < x < 45^\circ$ OR/OF $x \in (-45^\circ; 45^\circ)$ ACCEPT/ AANVAAR: $-45^\circ \leq x \leq 45^\circ$	$\checkmark\checkmark$ answer/antwoord (2)
5.5	$x = -45^\circ$ or/of $x = 135^\circ$	$\checkmark -45^\circ$ $\checkmark 135^\circ$ (2)
5.6	$g(x) = \cos(x + 15^\circ)$	$\checkmark$ answer/antwoord (1)
SA EXAM PAPERS [11]		


SA EXAM PAPERS

Proudly South African

QUESTION/VRAAG 6

6.1




6.1.1	$\hat{B}_1 = 55^\circ$ ($\angle$ s opp equal radii)/($\angle$ e teenoor gelyke radiusse) OR/OF $OA = OB$ (radii) (radiusse) $\hat{B}_1 = 55^\circ$ ($\angle$ s opp equal sides)/($\angle$ e teenoor gelyke radiusse)	$\checkmark$ S $\checkmark$ R (2) $\checkmark$ $OA = OB$ (radii) (radiusse) $\checkmark$ S (2)
6.1.2	$\hat{O}_1 = 70^\circ$ (sum of/som van $\angle$ s in Δ)/($\angle$ e van Δ)	$\checkmark$ S $\checkmark$ R (2)
6.1.3	$\hat{D}_1 = 35^\circ$ ($\angle$ at centre = $2 \times \angle$ at circumference) (midpts $\angle$ = $2 \times$ omtreks $\angle$)	$\checkmark$ S $\checkmark$ R (2)
6.1.4	$\hat{C}_2 = 70^\circ$ (ext $\angle$ of a cyclic quad)/(buite $\angle$ van kvh) OR $\hat{C}_1 = 110^\circ$ (opp $\angle$ s of a cyclic quad)/(teenoorste $\angle$ e kvh) $\hat{C}_2 = 70^\circ$ ($\angle$ s on a st line)/($\angle$ e op reguitlyn)	$\checkmark$ S $\checkmark$ R (2) $\checkmark$ S/R $\checkmark$ S (2)



SA EXAM PAPERS

Proudly South African

6.2



6.2.1	$\hat{S}_1 = x$ (tan chord theorem)/($\angle$ tussen raaklyn en koord)	✓ S	✓ R	(2)
6.2.2	$\hat{R}_2 = 180^\circ - (\hat{S}_1 + 45^\circ)$ (sum of $\angle$ s in Δ)/($\angle$ e van Δ) $= 180^\circ - (x + 45^\circ)$ $\hat{R}_2 + \hat{P} = 180^\circ$ (opp $\angle$ s of cyclic quad)/(teenoorste $\angle$ e kvh) $\therefore 180^\circ - (x + 45^\circ) + \hat{P} = 180^\circ$ $\hat{P} = x + 45^\circ$ OR/OF $\hat{R}_1 = 45^\circ$ (tan chord theorem)($\angle$ tussen rlyn en koord) $\hat{R}_2 = 180^\circ - (x + 45^\circ)$ ($\angle$ s on st line)/($\angle$ e op reguitlyn) $\hat{R}_2 + \hat{P} = 180^\circ$ (opp $\angle$ s of cyclic quad)/teenoorste $\angle$ e kvh $\therefore 180^\circ - (x + 45^\circ) + \hat{P} = 180^\circ$ $\hat{P} = x + 45^\circ$	✓ S	✓ R	(4)
		✓ S	✓ R	(4)
[14]				



SA EXAM PAPERS



Proudly South African

MARKING GUIDELINES/
NASIENRIGLYNE

MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)

GR12 0626

QUESTION/VRAAG 7


7.1.1	SAS/(S<S)	✓ R	(1)
7.1.2	 		
	$\widehat{ADE} = \widehat{Q} \quad (\equiv \Delta s)/(\equiv \Delta e)$ $\widehat{ADE} = \widehat{B} \quad (\text{both} = \widehat{Q})/(\text{beide} = \widehat{Q})$ $\therefore DE \parallel BC \quad (\text{corresp } \angle s =)/(\text{ooreenk } \angle e =)$ $\frac{AB}{AD} = \frac{AC}{AE} \quad (\text{prop theorem; } DE \parallel BC)/(\text{lyn } \parallel \text{ een sy van } \Delta; DE \parallel BC)$ But/Maar $AD = PQ$ and/en $AE = PR$ (given)/(gegee) $\therefore \frac{AB}{PQ} = \frac{AC}{PR}$	✓ S ✓ S ✓ R ✓ S ✓ R	(5)
	OR/OF In $\triangle ABC$ and $\triangle PQR$: $\hat{A} = \hat{P} \quad (\text{given/gegee})$ $\hat{B} = \hat{Q} \quad (\text{given/gegee})$ $\hat{C} = \hat{R} \quad (\text{given/gegee})$ $\triangle ABC \parallel \triangle PQR \quad (<<<)$ $\frac{AB}{PQ} = \frac{AC}{PR}$	✓ S ✓ S ✓ R ✓ S ✓ R	(5)



SA EXAM PAPERS

Proudly South African

7.2



7.2.1	$\widehat{B} = 90^\circ$ (given)/(gegee) $\widehat{D}_1 = 90^\circ$ ($\angle$ in a semi – circle)/($\angle$ in halwe sirkel) $\therefore ABCD$ is a cyclic quadrilateral (converse: ext $\angle$ of a cyclic quad) / $ABCD$ is 'n kvh (buite $\angle$ van vierhoek = teenoorst binne $\angle$) OR/OF $\widehat{B} = 90^\circ$ (given) /(gegee) $\widehat{D}_1 = 90^\circ$ ($\angle$ in a semicircle)/($\angle$ in 'n halwe sirkel) $\widehat{D}_2 + \widehat{D}_3 = 90^\circ$ ($\angle$ s on a st line)/($\angle$ e op reguitlyn) $\therefore ABCD$ is a cyclic quadrilateral (converse: opp $\angle$ s of a cyclic quad)/ $ABCD$ is 'n kvh (buite $\angle$ van vierhoek = teenoorst binne $\angle$)	$\checkmark$ S $\checkmark$ R $\checkmark$ R (3) OR/OF $\checkmark$ S $\checkmark$ R $\checkmark$ R (3)
7.2.2	$\widehat{A}_1 = \widehat{D}_2$ ($\angle$ s in the same seg)/($\angle$ e in dies sirkel segm) $\widehat{D}_2 = \widehat{F}$ (tan chord theorem)/($\angle$ tussen raaklyn en koord) $\therefore \widehat{A}_1 = \widehat{F}$ (both = $\widehat{D}_2$) / (beide = $\widehat{D}_2$) $\therefore AB$ is a tangent/is 'n raaklyn (converse: tan – chord thrm OR $\angle$ between line and chord) (omg $\angle$ e in dies sirkel segm OR $\angle$ tussen lyn en koord = $\angle$ in teenoorst sirkel segm)	$\checkmark$ S $\checkmark$ R $\checkmark$ S/R $\checkmark$ R (4)

Proudly South African

[13]


MARKING GUIDELINES/
NASIENRIGLYNE

MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)

GR12 0626

QUESTION/VRAAG 8

8.1



$AQ = 4$ units/eenhede (line from centre $\perp$ chord) /
(loodlyn uit midpt O na koord)

$OC = OA = 8 - x$ (radii)/(radiusse)

$OA^2 = x^2 + AQ^2$ (Pythagoras)

$(8 - x)^2 = x^2 + 4^2$

$64 - 16x + x^2 = x^2 + 16$

$16x = 48$

$\therefore x = 3$

$\therefore OC = 5$ units/eenhede

OR/OF

$OC = OA$ (radii)/(radiusse)

$OQ = 8 - OC$

$AQ = 4$ units/eenhede (line from centre $\perp$ chord) /
(loodlyn uit midpt O na koord)

$OA^2 = OQ^2 + 4^2$ (Pythagoras)

$OC^2 = (8 - OC)^2 + 4^2$

$OC^2 = 64 - 16OC + OC^2 + 16$

$16OC = 80$

$\therefore OC = 5$ units/eenhede

✓ S ✓ R

✓ $OA = 8 - x$

✓ S

(5)

✓ $OC = 5$ units/
eenhede

OR/OF

✓ $OQ = 8 - OC$

✓ S ✓ R

✓ S


(5)

SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626

8.2



8.2.1	4 : 5	✓ S (1)
8.2.2	$\frac{AC}{AB} = \frac{9}{4}$ (given/gegee) $AC = \frac{9}{4} \times 6$ units/eenhede $AC = \frac{27}{2}$ units/eenhede $= 13,5$ units/eenhede	✓ substitution /substitusie ✓ answer/antwoord (2)
8.2.3	$\frac{\text{Area of } \triangle BDC}{\text{Area of } \triangle BED} = \frac{\frac{1}{2} \times CD \times BQ}{\frac{1}{2} \times DE \times BQ}$ $= \frac{CD}{DE}$ $= \frac{5}{4} \text{ (prop theorem; } BD \parallel AE \text{) / line one side of } \Delta$ $\text{(lyn een sy van } \Delta \text{)}$	✓ $\frac{\text{Area of } \triangle BDC}{\text{Area of } \triangle BED} = \frac{CD}{DE}$ $\frac{\text{Opp } \triangle BDC}{\text{Opp } \triangle BED} = \frac{CD}{DE}$ ✓ S/R (2)
8.2.4	$\frac{EC}{FC} = \frac{BC}{AC} = \frac{5}{9}$ (prop theorem; $BE \parallel AF$) / line one side of Δ $\text{(lyn een sy van } \Delta \text{)}$ $\frac{EC}{15} = \frac{5}{9}$ $EC = \frac{25}{3} = 8,33$ $\frac{DE}{EC} = \frac{4}{9}$ (prop theorem; $BD \parallel AE$) / line one side of Δ $\text{(lyn een sy van } \Delta \text{)}$ $DE = \frac{4}{9} \times EC$ $= \frac{4}{9} \times \frac{25}{3}$ $\therefore DE = \frac{100}{27}$ units/eenhede = 3,70 units/eenhede	✓ $\frac{EC}{FC} = \frac{5}{9}$ ✓ $EC = \frac{25}{3} = 8,33$ ✓ $\frac{DE}{EC} = \frac{4}{9}$ ✓ answer/antwoord (4)

SA EXAM PAPERS

Proudly South African


[14]

MARKING GUIDELINES/
NASIENRIGLYNE

MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)

GR12 0626

QUESTION/VRAAG 9




9.1	In $\triangle BEF$ and/en $\triangle CGF$, $\hat{E}_2 = \hat{G}_1$ (both = 90°, given)/(beide = 90°, gegee) $\hat{B}_1 = \hat{C}_2$ (tan chord theorem)/($\angle$ tussen raaklyn en koord) $\hat{E}\hat{F}\hat{B} = \hat{C}\hat{F}\hat{G}$ (sum of/som van $\angle$s in Δ)/($\angle$ e van Δ) $\therefore \triangle BEF \parallel \triangle CGF$ ($\angle \angle \angle$)	✓ $\hat{E}_2 = \hat{G}_1$ ✓ S/R ✓ $\angle \angle \angle$ OR/OF Sum of/som van $\angle$s in Δ/ ($\angle$ e van Δ) (3)
-----	--	---



SA EXAM PAPERS

Proudly South African

**MARKING GUIDELINES/
NASIENRIGLYNE**
**MATHEMATICS/WISKUNDE
(PAPER/VRAESTEL 2)**
GR12 0626


9.2	In $\triangle CDF$ and/en $\triangle BGF$, $\widehat{D}_2 = \widehat{G}_2$ (both = 90°, given)/(beide = 90°, gegee) $\widehat{C}_1 = \widehat{B}_2$ (tan chord theorem)/($\angle$ tussen raaklyn en koord) $\widehat{C}\widehat{F}\widehat{D} = \widehat{B}\widehat{F}\widehat{G}$ (sum of $\angle$ s in $\triangle$)/($\angle$ e van $\triangle$) $\therefore \triangle CDF \parallel \triangle BGF$ ($\angle \angle \angle$) $\triangle BEF \parallel \triangle CGF$ (proven) $\frac{FD}{FG} = \frac{FC}{FB}$ ($\triangle CDF \parallel \triangle BGF$) $\frac{FG}{FE} = \frac{FC}{FB}$ ($\triangle BEF \parallel \triangle CGF$) $\frac{FD}{FG} = \frac{FG}{FE}$ (both /beide = $\frac{FC}{FB}$) $\therefore FG^2 = FE \times FD$	✓ S ✓ R ✓ $\triangle CDF \parallel \triangle BGF$ ✓ S/R ✓ S ✓ $\frac{FD}{FG} = \frac{FG}{FE}$ (6)
		[9]

TOTAL/TOTAAL: 145

SA EXAM PAPERS

Proudly South African