

SA EXAM PAPERS This Paper was downloaded from SAEXAMPAPERS
SA's Leading Past Year

Exam Paper Portal



You have Downloaded, yet Another Great Resource to assist you with your Studies 😊

Thank You for Supporting SA Exam Papers

Your Leading Past Year Exam Paper Resource Portal

Visit us @ www.saexampapers.co.za



**SA EXAM
PAPERS**

SA EXAM PAPERS

Proudly South African



This Paper was downloaded from SAEXAMPAPERS

KWAZULU-NATAL PROVINCE

EDUCATION
REPUBLIC OF SOUTH AFRICA

PROVINCIAL STANDARDISED ASSESSMENT

GRADE 12

PHYSICAL SCIENCES P2 (CHEMISTRY)

JUNE 2026

MARKS: 150

TIME: 3 hours

**This question paper consists of 16 pages and
an 2 data sheets.**



SA EXAM PAPERS

Proudly South African



INSTRUCTIONS AND INFORMATION

- 1 Write your name in the appropriate spaces on the ANSWER BOOK.
- 2 This question paper consists of SEVEN questions. Answer ALL the questions in the ANSWER BOOK.
- 3 Start EACH question on a NEW page in the ANSWER BOOK.
- 4 Number the answers correctly according to the numbering system used in this question paper.
- 5 Leave ONE line between two sub-questions, for example between QUESTION 2.1 and 2.2.
- 6 You may use a non-programmable calculator.
- 7 You may use appropriate mathematical instruments.
- 8 Show ALL formulae and substitutions in ALL calculations.
- 9 Round off your final numerical answers to a minimum of TWO decimal places.
- 10 Give brief motivations, discussions et cetera where required.
- 11 You are advised to use the attached DATA SHEETS.
- 12 Write neatly and legibly.






QUESTION 1

Four possible options are provided as answers to the following questions. Each question has only one correct answer. Choose the correct answer and write the letter (A – D) next to the relevant question number (1.1 – 1.10) on the answer sheet.

1.1 The IUPAC name of the following organic compound is ...




- A 3-methylpent-3-ene
 B 3-ethylpent-3-ene
 C 2-ethylbut-1-ene
 D 2-ethylbut-2-ene (2)

1.2 Which ONE of the following is the EMPIRICAL formula of ethyl ethanoate?

- A C₂H₄O
 B C₂H₂O
 C C₄H₄O
 D C₄H₈O₂ (2)

1.3 Consider the reaction represented below:




The term that describes this reaction is ...

- A Hydrolysis
 B Hydrogenation
 C Combustion
 D Cracking (2)



- 1.4 Consider the vapour pressure against temperature curves for THREE STRAIGHT-CHAIN ISOMERS under standard atmospheric pressure.



Consider the statements regarding the curves for the THREE CHAIN ISOMERS.


- I Compound A has the strongest intermolecular forces
- II Compound B has a higher boiling point than compound C.
- III Compound B is in the gaseous phase at temperature T_3 .

Which of the above statement(s) is/are true?

- A I only
- B II only
- C III only
- D I and II only

(2)

- 1.5 Consider the DEHYDRATION reaction below:



The IUPAC names of the MAJOR and MINOR products of this reaction are ...


	MAJOR	MINOR
A	4-methylhex-3-ene	4-methylhex-2-ene
B	3-methylhex-3-ene	3-methylhex-4-ene
C	3-methylhex-3-ene	4-methylhex-2-ene
D	4-methylhex-2-ene	3-methylhex-3-ene

(2)





- 1.6 The Maxwell-Boltzmann distribution curve **P** represents the number of molecules against kinetic energy for a certain reaction.



Curve **Q** is obtained after a change was made to one reaction condition. Which ONE of the following changes resulted in curve **Q**?

- A Increase in activation energy
- B Increase in the concentration of the reactants
- C Increase in temperature
- D Addition of a catalyst


(2)

- 1.7 Which ONE of the following statements concerning equilibrium is NOT correct?

- A Equilibrium is reached in a closed system.
- B At equilibrium, the concentrations of reactants and products are constant.
- C At equilibrium, the concentration of reactants is equal to concentration of products.
- D At equilibrium, the concentration of reactants and products is not necessarily the same.

(2)

- 1.8 Consider the equation below for a hypothetical reaction.



If the equilibrium constant $K_c = 1 \times 10^{-4}$, then ...

- A $[\text{A}] > [\text{C}][\text{D}]$
- B $[\text{A}] < [\text{C}][\text{D}]$
- C $[\text{A}] = [\text{C}][\text{D}]$
- D $0 = [\text{C}][\text{D}]$

(2)





- 1.9 The following values represents the ionisation constant, k_b , for four different bases, that were obtained at a temperature of 25 °C:

65×10^{-8}	40×10^{-1}	28×10^{-5}	30×10^{-7}
---------------------	---------------------	---------------------	---------------------

Which ONE of these values represents the STRONGEST base?

- A 40×10^{-1}
 - B 28×10^{-5}
 - C 30×10^{-7}
 - D 65×10^{-8}
- (2)

- 1.10 The endpoint in a titration is the exact point where ...


- A equal number of moles of acid and base have reacted.
 - B the indicator changes colour.
 - C equal masses of base and acid have reacted.
 - D neutralisation occurs.
- (2)

[20]



**QUESTION 2 (Start on a new page)**

2.1 A certain compound **A**, has the following condensed structural formula:



Another compound **B** is a POSITIONAL ISOMER of compound **A**, while compound **C** is a CHAIN ISOMER of compound **A**.

2.1.1 Define the term *structural isomers*. (2)

2.1.2 Is compound **A** SATURATED or UNSATURATED?
Give a reason for your answer. (2)

2.1.3 Write down the IUPAC name of compound **B**. (2)

2.1.4 Draw the structural formula of compound **C**. (3)

2.2 Consider TWO organic compounds, **1** and **2**, shown in the table below.

Compound 1	Compound 2
	$\text{C}_x\text{H}_y\text{O}_z$

2.2.1 To which homologous series does compound **1** belong? (1)

2.2.2 Write down the IUPAC name of compound **1**. (3)

Compound **2** is a **carboxylic acid** with a relative molecular mass of 74.

2.2.3 Determine, by means of calculations, the values of X, Y, and Z. (5)

2.2.4 Draw the structural formula of compound **2**. (3)

[21]



QUESTION 3 (Start on a new page)

Two parallel investigations were conducted to investigate the effect of relative molecular mass on the boiling point of organic compounds. In one investigation, straight-chain alkanes were used; in the other, straight-chain primary alcohols. The results obtained are shown in the tables below.

ALKANES		
Compound	Relative Molecular mass	Boiling point (°C)
C ₂ H ₆	30	-89
C ₃ H ₈	44	-42
C ₄ H ₁₀	58	0
C ₅ H ₁₂	72	36

ALCOHOLS		
Compound	Relative Molecular mass	Boiling point (°C)
CH ₄ O	32	65
C ₂ H ₆ O	46	78
C ₃ H ₈ O	60	97
C ₄ H ₁₀ O	74	117

- 3.1 Write down a suitable conclusion for these investigations based on the data provided in the above tables. (2)
- 3.2 Consider the table containing Alkanes above:
- 3.2.1 Define the term *hydrocarbon*. (2)
- 3.2.2 Which ONE of these alkanes is a LIQUID at room temperature? Give a reason for your answer. (2)
- 3.2.3 Write down the balanced equation for the complete combustion of Propane. (3)
- 3.3 Consider the table containing Alcohols above.
- 3.3.1 Define the term *Primary alcohol*. (2)
- 3.3.2 Explain, by referring to the TYPE and STRENGTH of intermolecular forces involved, WHY the boiling point of C₄H₁₀O is higher than that of the other alcohols in the table. (3)
- 3.4 It is observed that, although C₅H₁₂ and C₄H₁₀O have comparable molecular masses, the boiling point of C₄H₁₀O is much higher than that of C₅H₁₂. Fully explain this observation. Refer to the TYPE and STRENGTH of the intermolecular forces present. (4)

[18]



**QUESTION 4 (Start on a new page)**

4.1 2-Bromobutane is mixed with NaOH in two different reactions, as shown below:



G and **Q** are organic compounds, and **M** is an inorganic compound.

4.1.1 Distinguish between concentrated base and dilute base. (2)

For **Reaction 1**, write down:

4.1.2 The type of reaction that takes place. (1)

4.1.3 The homologous series to which compound **G** belongs. (1)

4.1.4 The IUPAC name of compound **G**. (2)

Compound **Q** is the major product of **Reaction 2**:

4.1.5 Name the type of reaction that takes place in **Reaction 2**. (1)

4.1.6 Besides heat and concentrated NaOH, write down ONE other condition necessary for **Reaction 2**. (1)


4.1.7 Draw the structural formula of compound **Q**. (2)

4.1.8 Write down the chemical NAME or FORMULA of compound **M**. (1)





- 4.2 Propyl ethanoate can be prepared from prop-1-ene by a four-step process as shown in the flow diagram below.



Refer to the diagram above and write down: (1)

4.2.1 The NAME or FORMULA of the catalyst needed for reaction **A**.

4.2.2 The letter(s) that represent(s) Substitution reaction(s)? (2)

4.2.3 The IUPAC name of the Alcohol used in reaction **Q**. (2)

4.2.4 TWO reaction conditions for reaction **Q**. (2)

4.2.5 The NAME of the function group of Propyl ethanoate. (1)

4.2.6 The balanced equation of reaction **Q**, using MOLECULAR FORMULAE. (3)


[22]




QUESTION 5 (Start on a new page)

- 5.1 During the contact process, Sulphur dioxide reacts with Oxygen, in the presence of Vanadium oxide (V_2O_5), to form Sulphur trioxide.

The balanced chemical equation and the potential energy versus the course of reaction graph are given below.



Graph of Potential energy versus the course of the reaction.




- 5.1.1 Define the term *Activation energy*. (2)
- 5.1.2 Write down the value of the activation energy for the forward reaction in the PRESENCE of vanadium oxide. (1)
- 5.1.3 Calculate the energy (in kilojoules) released to make 24 g of SO_3 . (4)





- 5.2 Two investigations, **I** and **II**, have been performed to investigate factors that affect the rate of chemical reactions.

The balanced chemical equation for the reaction between Aluminium (Al) and sulfuric acid (H₂SO₄) solution is shown below.



In both investigations, 8 g of Aluminium and EXCESS dilute Sulphuric acid are used, while ensuring that the Aluminium is completely covered by the acid.

INVESTIGATION I

Two experiments are conducted using the conditions as shown in the table below.

	Mass of Al (g)	State of division of Al	Concentration of H ₂ SO ₄ (mol.dm ⁻³)	Temperature (°C)
Experiment 1	8	Lumps	0, 1	30
Experiment 2	8	Powder	0, 1	30

- 5.2.1 Write down an investigative question for this investigation. (2)
- 5.2.2 Which experiment, **1** or **2**, will reach completion within a shorter period of time? Give a reason. (2)

INVESTIGATION II

Two other experiments, **3** and **4**, are conducted using different conditions as shown in the table below.

	Mass of Al (g)	State of division of Al	Concentration of H ₂ SO ₄ (mol.dm ⁻³)	Temperature (°C)
Experiment 3	8	Powder	0, 1	30
Experiment 4	8	Powder	0, 1	40

As the reactions progress, it is observed that **Experiment 4** has a higher reaction rate than **Experiment 3**.

- 5.2.3 Write down the independent variable for this investigation. (1)
- 5.2.4 Use the collision theory to explain this observation. (3)
- 5.3 5 g of a sample of IMPURE Sodium carbonate (Na₂CO₃) is reacted with EXCESS dilute hydrochloric acid (HCl) according to the balanced equation below.



The average rate of CO₂ produced, at room temperature, is 0,005 dm³.s⁻¹. Take the molar volume of gas as 24 dm³.mol⁻¹, and assume that no impurities react. Calculate the percentage purity of Na₂CO₃ if the time for this reaction to reach completion is 3 minutes.


Proudly South African

(7)
[22]



QUESTION 6 (Start on a new page)


Ammonia (NH_3) is an important product in the fertiliser industry. The balanced equation below shows the reaction between Nitrogen and Hydrogen to produce Ammonia.



This reaction is reversible, and it will eventually reach a stage where the rate of the forward reaction equals the rate of the reverse reaction.

6.1 Write down the term that is described by the underlined sentence. (1)

Adjustments to the equilibrium mixture conditions are made to determine which condition results in the highest Ammonia yield. The graphs below represent the results obtained.



6.2 State *Le Chatelier's principle*. (2)


6.3 At time t₃, the temperature was adjusted.

Use *Le Chatelier's principle* to explain whether the temperature was INCREASED or DECREASED. (4)





- 6.4 At time t_2 , the pressure of the system was increased by decreasing the volume of the container. The INCOMPLETE graph, starting from the equilibrium after t_1 , of the reaction rate versus time is shown below.




- 6.4.1 Will the increase in pressure favour the FORWARD or the REVERSE reaction? Give a reason (2)

- 6.4.2 Sketch the reaction rate vs time graph showing:

- Immediate effect at t_2
- New equilibrium after t_3 . (4)

- 6.5 In a different experiment, Nitrogen monoxide is reacted with oxygen in a closed container at 100°C . The reaction reaches equilibrium according to the following chemical equation:



Initially 8 mol of NO and 8 mol of O_2 were placed in the reaction container. When equilibrium was established, it was found that 65% of oxygen was still present in the container. The equilibrium constant, K_c , for the reaction at 100°C is 3,15. (9)
Calculate the volume of the container used for this reaction.



[22]



**QUESTION 7 (Start on a new page)**

7.1 Sulphuric acid, H_2SO_4 , is a strong acid used in car batteries.

The equations below represent the ionisation of sulphuric acid.



7.1.1 Is Sulphuric acid a MONOPROTIC or DIPROTIC acid?
Give a reason for your answer. (2)

7.1.2 Write down the FORMULA of a species that acts as ampholyte in the
above reactions. (1)

7.1.3 Write down the NAME of the conjugate base of hydrogen sulphate ion. (1)

7.2 Sodium hydrogen carbonate, **NaHCO_3** , is a salt with a pH slightly greater than 7.

7.2.1 Define the term *Hydrolysis*. (2)

7.2.2 Write down a balanced equation that explains why a sodium hydrogen
carbonate solution has a pH greater than 7. (3)





- 7.3 During a Chemistry practical lesson, a group of Grade 12 learners notices three conical flasks, **A**, **B**, and **C**, labelled as shown below. The learners mix the contents of conical flasks **A** and **B**, assuming that both are sulphuric acid solutions. The total volume of the mixture obtained is now 500 ml.

Conical flask A	Conical flask B	Conical flask C
<p>Acid 200 ml</p>	<p>H₂SO₄ 300 ml</p>	<p>NaOH 300 ml</p>
$c = \text{Unknown}$	$c = 0,12 \text{ mol}\cdot\text{dm}^{-3}$	$c = 0,305 \text{ mol}\cdot\text{dm}^{-3}$

- 7.3.1 Calculate the pH of the H₂SO₄ solution in conical flask **B**, before the learners mixed it with the acid in conical flask **A**. (4)

Learners are now told that conical flask **A** contained a hydrochloric acid (HCl) solution of unknown concentration and are consequently tasked with performing a titration to neutralise the 500 ml acidic solution with the NaOH solution in conical flask **C**.

- 7.3.2 What indicator would be suitable for this titration?

Choose from METHYL ORANGE, PHENOLPHTHALEIN, or BROMOTHYMOLO BLUE.

Give a reason for your choice. (2)

After performing the titration, they find that 20 ml of the acidic solution is completely neutralised by 16 ml of the Sodium hydroxide solution (NaOH) of concentration 0,305 mol·dm⁻³.

- 7.3.3 Calculate the concentration of the HCl solution that was initially present in conical flask **A** before it was mixed with the content of conical flask **B**. Conical flask **B** contained 0,12 mol·dm⁻³ H₂SO₄. (10)

[25]

TOTAL MARKS: 150





**DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)**

**GEGEWENS VIR FISIESTE WETENSKAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)**

TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESTE KONSTANTES

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure <i>Standaarddruk</i>	p^\ominus	$1,013 \times 10^5 \text{ Pa}$
Molar gas volume at STP <i>Molêre gasvolume by STD</i>	V_m	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard temperature <i>Standaardtemperatuur</i>	T^\ominus	273 K
Charge on electron <i>Lading op elektron</i>	e	$-1,6 \times 10^{-19} \text{ C}$
Avogadro's constant <i>Avogadro-konstante</i>	N_A	$6,02 \times 10^{23} \text{ mol}^{-1}$

TABLE 2: FORMULAE/TABEL 2: FORMULES

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ or/of $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$
$\frac{C_a V_a}{C_b V_b} = \frac{n_a}{n_b}$	$\text{pH} = -\log[\text{H}_3\text{O}^+]$
$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14}$ at/by 298 K	
$E_{\text{cell}}^\ominus = E_{\text{cathode}}^\ominus - E_{\text{anode}}^\ominus$ / $E_{\text{sel}}^\ominus = E_{\text{katode}}^\ominus - E_{\text{anode}}^\ominus$ or/of $E_{\text{cell}}^\ominus = E_{\text{reduction}}^\ominus - E_{\text{oxidation}}^\ominus$ / $E_{\text{sel}}^\ominus = E_{\text{reduksie}}^\ominus - E_{\text{oksidasie}}^\ominus$ or/of $E_{\text{cell}}^\ominus = E_{\text{oxidisingagent}}^\ominus - E_{\text{reducingagent}}^\ominus$ / $E_{\text{sel}}^\ominus = E_{\text{oksideermiddel}}^\ominus - E_{\text{reduseermiddel}}^\ominus$	



